

From Chaos To Constructive Relativity

Has the Evidence Finally Reached a Threshold That Warrants Rejecting the Century-Old LCDM Hot Dense Center Origin Model?

DR JM NIPOK N.J.I.T.
orcid.org/0009-0006-3940-4450
osf.io/t8zny/overview

Copyright CC BY-NC-SA 4.0. June 2026
doi.org/10.13140/RG.2.2.23479.79528
Version 4.19

Abstract

The Lambda Cold Dark Matter (Λ CDM) model now confronts substantial empirical tension across 231 independent observational anomalies: the 5σ Hubble disagreement ($H_0 = 73.0 \pm 1.0$ vs 67.4 ± 0.5 km s⁻¹ Mpc⁻¹), the S8 growth-rate deficit, JWST's high-redshift mass and morphology problems, the SPT2349-56 ICM thermal energy excess at $z = 4.3$, gigaparsec-scale anomalous structures (Big Ring, Giant Arc), satellite-plane co-rotation across six independent hosts (joint $P \approx 2 \times 10^{-14}$), and the 10^{120} -magnitude vacuum-energy fine-tuning, yet retains its central commitment to a hot-dense-singular cosmic origin requiring inflation, cold-dark-matter particles, and a fundamental cosmological constant. We present Successive Collision Theory (SCT), a single-assumption-change framework grounded in 69 numbered premises (P1-P69) using only standard General Relativity, Special Relativity, the Standard Model, and lattice QCD applied to an eternal infinite universe. SCT replaces the singular origin with a superluminal collision between nested comoving structures, producing the thermalized debris field that became our visible universe.

The "From Chaos to Consilience" series develops the framework, organized as Series 1 (the cosmological papers, including this Paper 17) followed by Series 2 (the carrier-field / foundational papers, beginning with Series 2 Paper 1); the companion list in §XIV defines the boundary. All 231

cataloged tensions reduce to 11 Primary Concept Groups (M1-M11), each identifying one keystone premise whose removal would collapse that class of resolutions. Three low-parameter results stand out: $A^* = 6.173 = 1/f_b$ [DERIVED, using the virial/cluster baryon fraction $f_b = 0.162$; this is a consistency check, not a zero-free-parameter prediction], consistent at 0.6% with HIFLUGCS+CLASH (Paper 12); $R_b = 0.2545 \pm 0.032$ (dimensionless cascade baryon-loss/heating ratio from cascade geometry, agrees with observed 0.260 ± 0.002 at 0.17σ , Series 2 Paper 1); $N_{\text{eff}} = 2.514 \pm 0.050$ (effective relativistic species; if SCT is correct, CMB-S4 would separate 2.514 from the Standard-Model 3.046 by $\sim 17.7\sigma$ (forecast) in central values, though present Planck data already show a $\sim 2.8\sigma$ tension with 2.514, Series 2 Paper 1). The unified field equation [$0.08 \text{ fm} \leq r$]: $G_{\mu\nu} + \Lambda_{\text{eff}}(x,t) g_{\mu\nu} = (8\pi G/c^4)[T_{\mu\nu} + T^{\text{sup}}_{\mu\nu}(A)]$ is GR with three identified additions, dynamical Λ_{eff} , coherent superposition $T^{\text{sup}}_{\mu\nu}$, and a QCD lower-boundary domain restriction, not a replacement (Paper 8). The compiled predictions ledger (see Paper S1_18) makes SCT decidable. Primary near-term discriminant: CMB-S4 N_{eff} measurement (2030+). The accumulated evidence has reached a threshold where the inherited Λ CDM origin model warrants reconsideration on its own merits.

Keywords: successive collision theory; SCT; Λ CDM tensions; 231-tension catalog; 69 numbered premises; M1-M11 Primary Concept Groups; $A^* = 6.173$; $N_{\text{eff}} = 2.514$; $R_b = 0.2545$; CAR formula $c_s^2 = (1+R_b)/3$; polyquark cores; mesh dissipation; dynamical $\Lambda_{\text{eff}}(x,t)$; Hubble tension; S8 tension; gravitational superposition $T^{\text{sup}}_{\mu\nu}$; coherent amplification; hereditary proper time; frame-tree redshift; nested comoving frames; eternal infinite universe; superluminal pocket collision; collision-seeded structure formation; Big Bang alternative; dark matter alternative; dark

energy origin; SCT-MASTER unified field equation; QCD domain boundary 0.08 fm; CMB-S4 N_{eff} discriminant; 17.7σ forecast; DR JM NIPOK.

TOC I. Introduction II. Foundational Premises II.A. Category I: The Nature of the Universe (P1-P6) II.B. Category II: The Structure of the Universe (P7-P8) II.C. Category III: The Nature of Time (P9-P13) II.D. Category IV: The Nature of Dark Energy (P14-P19) II.E. Category V: Origin of Our Visible Universe (P20-P49) II.F. Category VI: The Nature of Dark Matter (P50-P56) II.G. Category VII: Our Place in the Universe and the Unified Field Equation (P57-P69) II.H. Theoretical Constraint Statement and Consistency Requirements

III. Geometric Structure of Nested Spacetime Pockets III.A. Nested Pocket Hierarchy: Definition and Structure III.B. Metric Structure Within and Between Pockets III.C. Collective Properties of Spacetime Pockets III.D. Boundary Conditions and Pocket Interfaces III.E. Consistency Requirements and Open Mathematical Tasks

IV. Relativistic Inheritance of Proper Time and Spacetime Perception IV.A. Conceptual Foundation: Time as Hereditary IV.B. Special Relativistic Time Dilation Across Nested Succession IV.C. Gravitational Time Dilation Across Nested Succession IV.D. Local Refinements Within Pockets IV.E. Proper-Time Recursion Formalism IV.F. Inherited Spatial and Causal Structure IV.G. Connection to Observational Cosmology IV.H. Required Derivations and Consistency Checks

V. Successive Superluminal Collisions: Replacing the Hot Dense Origin V.A. Conceptual Foundation: Superluminal Collisions Without SR Violation V.B. Collision Sequence and Thermalization V.C. Collision Geometry and Structure Formation V.D. Energy, Momentum, and Angular Momentum Conservation V.E. Entropy Production and the Second Law V.F.

Curvature Evolution and Gravitational Radiation V.G. Observational Signatures of Collision Origin V.H. Required Derivations and Consistency Checks

VI. Dark Energy as Gravitational Mesh Dissipation VI.A. Orbital Decay and Gravitational Binding Evolution VI.B. Effective Cosmological Term Λ_{eff} VI.C. Temporal and Spatial Variability of Λ_{eff} VI.D. Implications for the Hubble Tension VI.E. Observational Predictions and Falsifiability

VII. Cosmological Redshift from Hereditary Proper Time

Status update: The redshift formula $z = f(\text{pocket_depth}, \text{hereditary_time_path})$ is given in closed form by the hierarchical Lorentz frame-tree composition of Paper 2 (“From Chaos To Common Ancestry”). The single-boost approximation $1 + z = (1 + z_{\text{cosmo}})(1 + z_{\text{pec}})$ is replaced by the recursive product over the parent chain. Frame-tree corrections at the 10^{-5} - 10^{-4} level produce a ~ 0.5 -1% systematic between cluster-environment and field-environment supernovae, itself an observational discriminator testable with Pantheon+ + DESI subsamples. The Hubble law $H_0 \approx 67$ -73 km s⁻¹ Mpc⁻¹ is recovered statistically over a representative source-observer ensemble.

VII.A. Photon Propagation Through Nested Frames VII.B. Derivation of Redshift from Proper-Time Inheritance VII.C. Statistical Emergence of the Hubble Law VII.D. Predicted Deviations from Λ CDM

VIII. Dark-Matter-Like Phenomena from Gravitational Field Superposition VIII.A. Coherent Gravitational Superposition in Comoving Frames VIII.B. Effective Stress-Energy Tensor Modification VIII.C. Galaxy Rotation Curves and Lensing Predictions VIII.D. Cluster Dynamics and Large-Scale Structure

IX. Early Structure Formation and High-Redshift Observations IX.A. Collision-Seeded Density Perturbations IX.B. Rapid Galaxy and Black Hole Formation IX.C. Comparison with JWST Observations

X. Cosmic Microwave Background from Collision Thermalization X.A. Origin of the CMB Temperature X.B. Anisotropy Generation Mechanisms X.C. Power Spectrum and Non-Gaussian Signatures X.D. Comparison with Planck Observations

XI. Systematic Resolution of Observational Tensions XI.A. Expansion History and Distance Measurements XI.B. Structure Formation and Growth Rates XI.C. Large-Scale Anomalies and Alignments XI.D. Fine-Tuning and Initial Condition Problems

XII. Observational Predictions and Falsification Tests XII.A. Redshift-Dependent Dark Energy Signatures XII.B. Spatial Anisotropies and Bulk Flow Correlations XII.C. Rotation Curve and Lensing Discriminants XII.D. Gravitational Wave and Transient Signatures

XIII. Mathematical Framework and Outstanding Derivations XIII.A. Completed Derivations XIII.B. Explicit Mathematical Gaps XIII.C. Required Numerical Simulations

XIV. Discussion and Comparison with Alternative Models XIV.A. Comparison with Λ CDM XIV.B. Relation to Modified Gravity and Varying-Constant Models XIV.C. Conceptual and Philosophical Implications

XV. Conclusions and Future Work

Appendix A. Catalog of Λ CDM Tensions Addressed by SCT (231 Cases)

Introduction

For much of the past century, our theoretical commitments have been shaped as much by our discomfort with the unknown as by the empirical record itself. Finite temporal origins, bounded causal domains, and closed explanatory structures have often been preferred not because observations required them, but because the alternative, an eternal universe, an unbounded spacetime, or an infinite hierarchy of scales, was perceived as conceptually unwieldy or methodologically destabilizing. Yet scientific progress has never been advanced by constraining the universe to match our intuitions. It has advanced when we have allowed the universe to instruct us, even when its implications challenge our deepest assumptions.

The accumulating tensions within the standard framework, together with the growing body of observations that resist reconciliation within a finite, tightly delimited cosmology, now compel us to reconsider the foundational premises we have long taken for granted. A serious engagement with models that incorporate eternity and infinity is not an indulgence in metaphysics; it is a necessary step toward constructing theories capable of accommodating the full complexity revealed by contemporary data. Only by relinquishing our inherited fear of the unbounded can we hope to develop a cosmology that is empirically adequate, internally coherent, and conceptually complete.

The moment has therefore arrived for clarity, for methodological candor, and for the willingness to re-examine assumptions that have persisted more by tradition than by necessity. The time is now to stand up and declare that the emperor wears no clothes.

The Lambda Cold Dark Matter (Λ CDM) cosmological model has provided a remarkably successful phenomenological framework for describing large-scale structure, cosmic microwave background (CMB)

anisotropies, and the expansion history of the observable universe. Built upon the Friedmann-Lemaître-Robertson-Walker (FLRW) metric solution to Einstein's field equations, Λ CDM posits that approximately 95% of the universe's energy density consists of components, dark matter ($\sim 26\%$) and dark energy ($\sim 69\%$), for which no direct particle detection or fundamental physical explanation currently exists. The model extrapolates gravitational dynamics backward to a hot, dense initial singularity approximately 13.8 billion years ago, followed by an inflationary epoch that exponentially expanded spacetime by a factor exceeding 10^{26} within the first 10^{-32} seconds. Despite its observational successes, Λ CDM faces significant and growing empirical challenges. The Hubble constant H_0 exhibits a persistent tension at the 5σ level between measurements from the early universe ($H_0 = 67.4 \pm 0.5 \text{ km s}^{-1} \text{ Mpc}^{-1}$ from CMB acoustic peaks via Planck satellite data) and late-universe distance ladder calibrations ($H_0 = 73.0 \pm 1.0 \text{ km s}^{-1} \text{ Mpc}^{-1}$ from Cepheid-calibrated Type Ia supernovae), suggesting either unidentified systematic errors or new physics beyond the standard model. The James Webb Space Telescope (JWST) has revealed massive galaxies ($M_* > 10^{10} M_\odot$) and supermassive black holes ($M_{\text{BH}} > 10^6 M_\odot$) at redshifts $z > 10$, corresponding to cosmic ages less than 500 million years post-singularity, timescales apparently insufficient for hierarchical assembly via gravitational instability growth from primordial perturbations under standard Λ CDM assumptions.

A comprehensive analysis identifies 231 distinct observational tensions spanning cosmological parameters, structure formation timescales, large-scale anomalies, and measurements of fundamental constants, suggesting potential systematic issues with the Λ CDM framework rather than isolated measurement uncertainties. Theoretically, Λ CDM's

cosmological constant Λ , interpreted as vacuum energy density $\rho_\Lambda = \Lambda c^2 / (8\pi G) \approx 6 \times 10^{-27} \text{ kg m}^{-3}$, differs from quantum field theory predictions by approximately 120 orders of magnitude, constituting perhaps the most severe fine-tuning problem in physics. Dark matter remains undetected despite decades of direct detection experiments (XENON, LUX, PandaX), collider searches at the Large Hadron Collider, and indirect observation campaigns targeting proposed particle candidates including WIMPs, axions, and sterile neutrinos. Inflation, while solving horizon, flatness, and monopole problems, introduces its own fine-tuning requirements for inflaton potential shapes and initial conditions, and generates predictions (e.g., primordial tensor modes with specific spectral indices) not yet conclusively confirmed observationally. The initial singularity itself represents a breakdown of General Relativity at Planck scales ($\ell_P \approx 10^{-35} \text{ m}$, $t_P \approx 10^{-43} \text{ s}$), requiring unverified quantum gravity physics. This paper presents Successive Collision Theory (SCT), a cosmological framework that addresses these challenges through reinterpretation of standard General Relativity and Special Relativity applied to an eternal, infinite universe. SCT is not constructed as a phenomenological extension of Λ CDM with additional free parameters, but rather as a deductive framework built upon 69 explicit foundational premises (P1-P69) that invoke only well-established physics. These premises are organized into seven categories addressing: (I) the nature of the universe, (II) cosmic structure, (III) the nature of time, (IV) dark energy, (V) the origin of our visible universe, (VI) dark matter phenomena, and (VII) our cosmic location. The theory makes three core conceptual departures from Λ CDM, each tied directly to specific subsets of the 69 premises:

1. Rejection of Initial Singularity and Inflation (P1-P6, P20-P41) The universe is eternal in time (P1) and infinite in space (P2), containing pre-existing matter distributed throughout (P4-P5) in hierarchical structures governed by scale-invariant General Relativity (P7). Our observable universe emerged not from a singular hot dense state at $t = 0$, but from successive superluminal collisions between immense nested comoving structures (“spacetime pockets,” P11) whose relative velocities can exceed c without violating relativity. The speed-of-light constraint applies to local acceleration within nested frames (P20), not to relative velocities between independently formed structures at vastly larger scales (P21-P22). Collisions at extreme velocities (potentially $3c$, $7c$, or higher; P23) pulverize pre-existing matter into hot dense plasma (P25), providing the thermalized initial conditions that Λ CDM attributes to a Planck-scale singularity, but without requiring unverified quantum gravity or inflation.

2. Reinterpretation of Dark Energy, with a Proper-Time Reading of Redshift (P9-P19) SCT RETAINS standard FLRW metric expansion with scale factor $a(t)$; cosmological redshift is metric wavelength-stretching, $\lambda_{\text{obs}}/\lambda_{\text{emit}} = a_{\text{obs}}/a_{\text{emit}}$ (see §VI.B). The “hereditary time” construction (P10) is a proper-time bookkeeping WITHIN the expanding metric, not a replacement for metric expansion: cumulative proper-time differences accumulated across nested pocket succession provide a consistency reinterpretation of the same observed $z(d)$, with a small ($\sim 0.5\text{-}1\%$) frame-tree correction as the only distinguishing signature. Each comoving frame inherits baseline proper-time behavior from its parent frame and refines it through local velocities and gravitational trajectories (P12). The cosmological constant Λ is reinterpreted as a dynamical ratio $\Lambda_{\text{eff}} = f(U_{\text{local}}, U_{\text{parent}})$ (P17) between gravitational binding energy within a pocket and cumulative gravitational influence from the parent pocket succession. Orbital decay

(P14) causes dissipation of gravitational “mesh strength” (P16) that observers interpret as accelerating expansion (P15). Unlike Λ CDM’s constant vacuum energy, Λ_{eff} varies spatially and temporally (P19), exhibiting long-term exponential growth (P18) but permitting short-term fluctuations that provide a candidate partial mechanism for the Hubble tension. 3. Gravitational Field Superposition Replacing Dark Matter (P50-P54, P66) Dark-matter-like phenomena, flat rotation curves, enhanced gravitational lensing, cluster velocity dispersions, emerge from constructive interference of gravitational fields among coherently moving masses within pockets (P50, P52), modifying the effective stress-energy tensor (P53) without requiring new particles. This constitutes a proposed modification to Einstein’s field equations where multiple bodies sharing a comoving frame create amplified spacetime curvature through gravitational wave superposition. Structure formation proceeds through collision-seeded overdensities (P55) rather than dark matter-driven hierarchical clustering. Compact object cores are stabilized by quark degeneracy pressure at inter-quark separations ~ 0.08 fm (P66) rather than collapsing to singularities, representing a third modification to GR incorporating lattice QCD results.

SECTION II: FOUNDATIONAL PREMISES Successive Collision Theory is constructed from 69 explicit foundational premises organized into seven categories. All subsequent theoretical claims, mathematical derivations, and observational predictions can be traced to these premises combined with standard General Relativity and Special Relativity. No speculative physics, undiscovered particles, or modifications to fundamental interactions will be introduced beyond what these premises explicitly require.

CATEGORY I: THE NATURE OF THE UNIVERSE (P1-P6) P1: Eternal Time
Time has no beginning and no end; there exists no minimal or maximal measurable temporal intervals.

EXPLANATION: The temporal dimension extends infinitely in both the past and the future directions, eliminating the need for cosmological initial conditions or an origin event at $t = 0$.

MATHEMATICAL COMMITMENT: Temporal coordinate $t \in (-\infty, +\infty)$ with no singularities required at finite times; field equations must not impose temporal boundaries.

OBSERVATIONAL COMMITMENT: No observational evidence for a temporal origin is necessary within SCT; apparent “beginning” phenomena (CMB, nucleosynthesis) arise from local collision events, not cosmic inception.

P2: Infinite Space Space has no boundary or edge in any direction; there exists no minimal or maximal measurable spatial intervals.

EXPLANATION: The spatial manifold is unbounded and extends indefinitely in all directions, with no compactification, topology constraints, or edge effects.

MATHEMATICAL COMMITMENT: Spatial coordinates x^i span \mathbb{R}^3 without compactification or boundary conditions; metrics must accommodate infinite spatial extent.

OBSERVATIONAL COMMITMENT: Large-scale homogeneity must extend arbitrarily far beyond the observable universe; no observational horizon represents a fundamental boundary.

P3: Embedded Observable Universe Eternal time and infinite space together imply that our observable universe constitutes an infinitesimal patch embedded within an unbounded, larger reality.

EXPLANATION: The finite observable universe (radius ~ 46.5 Gly) represents a local neighborhood within infinite spacetime, not a privileged or unique region.

MATHEMATICAL COMMITMENT: Observable universe radius $r_{\text{obs}} \ll R_{\text{universe}} \rightarrow \infty$; local curvature and dynamics represent boundary-value problems within infinite manifold.

OBSERVATIONAL COMMITMENT: Phenomena at our observational horizon must be interpretable as local features of a larger structure, not fundamental cosmic boundaries.

P4: Ubiquitous Mass-Energy If our observable patch exists within infinite space, it is statistically inconsistent to assume that mass-energy exists only here and not elsewhere.

EXPLANATION: The Copernican principle extended to infinite space: our local concentration of matter cannot be unique; similar structures must exist throughout the infinite manifold.

MATHEMATICAL COMMITMENT: Mass-energy density $\rho(x,t)$ must be non-zero across arbitrarily large regions of \mathbb{R}^3 , not concentrated solely in our observable neighborhood.

OBSERVATIONAL COMMITMENT: No direct observational requirement; this is a logical consistency premise preventing anthropic fine-tuning of initial conditions.

P5: Infinite Total Mass-Energy Given eternal time and infinite space, the universe must contain effectively infinite total mass and energy distributed throughout.

EXPLANATION: Integration of non-zero mass-energy density over infinite spatial volume yields unbounded total mass-energy content.

MATHEMATICAL COMMITMENT: Global mass-energy integral $\int_{\mathbb{R}^3} \rho d^3x \rightarrow \infty$; local conservation laws ($\nabla_{\mu} T^{\mu\nu} = 0$) must hold, but no global energy accounting is required or possible.

OBSERVATIONAL COMMITMENT: No observational test; this follows logically from P2 + P4.

P6: Large-Scale Homogeneity At the largest scales, the cosmological principle of isotropic homogeneity holds when we consider an eternally infinite 4D Minkowski spacetime.

EXPLANATION: Statistical averaging over sufficiently large volumes (exceeding correlation length) yields homogeneous and isotropic distributions, consistent with cosmological principle.

MATHEMATICAL COMMITMENT: Two-point correlation function $\xi(r) \rightarrow 0$ for separations $r \gg L_{\text{correlation}}$; power spectrum $P(k)$ must exhibit isotropy for $k \ll k_{\text{min}}$.

OBSERVATIONAL COMMITMENT: Must reproduce observed homogeneity at scales ~ 100 Mpc while permitting larger-scale structures (superfilaments, giant arcs) as fluctuations about the mean.

CATEGORY II: THE STRUCTURE OF THE UNIVERSE (P7-P8)

P7: Scale-Invariant Structure Reality is an eternal, scale-invariant “follow-the-leader” process: an infinite number of celestial objects form larger and

larger structures by following the scale-independent field equations of General Relativity.

EXPLANATION: Gravitational clustering proceeds hierarchically at all scales; Einstein's field equations contain no preferred length scale, permitting self-similar structure formation indefinitely.

MATHEMATICAL COMMITMENT: Solutions to $G_{\mu\nu} + \Lambda g_{\mu\nu} = (8\pi G/c^4) T_{\mu\nu}$ must permit nested hierarchical bound systems at arbitrarily large scales without requiring inflation, topology change, or scale-dependent modifications.

OBSERVATIONAL COMMITMENT: Observed hierarchy (planets → stars → galaxies → clusters → superclusters → filaments) must extend to scales beyond current observations; no maximum structure scale exists.

P8: Comoving Frames When we apply the field equations of General Relativity and the time-dilation/ length-contraction formulas of Special Relativity to eternal time and infinite space, we do not obtain isolated inflating bubble universes. Instead, we obtain a nested succession of larger and larger comoving frames of reference, where multiple celestial objects share the relative trajectory and velocity of their most massive "leader" objects.

EXPLANATION: GR+SR applied to infinite spacetime yields hierarchical comoving frames (e.g., Solar System → Galaxy → Local Group → Virgo Supercluster → ...) rather than disconnected inflationary bubbles.

MATHEMATICAL COMMITMENT: Each nested level α has a metric $g_{\mu\nu}^{(\alpha)}$ related to parent metric $g_{\mu\nu}^{(\alpha+1)}$ through Lorentz transformations $\Lambda^{\mu\nu(\beta)}(\alpha)$ encoding relative motion and gravitational redshift factors $\exp[\Phi^{(\alpha)}/c^2]$.

OBSERVATIONAL COMMITMENT: Our observable universe must be identifiable as one such frame within a larger succession, with testable consequences (e.g., dipole anisotropies from parent frame motion, bulk flows).

CATEGORY III: THE NATURE OF TIME (P9-P13)

P9: Lorentz Boost Relations Between Frames

Each comoving frame is related to its parent by a Lorentz boost.

EXPLANATION: This establishes the SR-consistent kinematic relation between adjacent levels of the hierarchy. Objects in pocket α move with collective velocity $v^{\hat{(\alpha)}}$ relative to parent pocket $\alpha+1$, and the frame-to-frame transformation is the standard SR Lorentz boost $\Lambda^{\mu\nu\beta(\alpha)}$ with $\beta^{\hat{(\alpha)}} = v^{\hat{(\alpha)}}/c$.

MATHEMATICAL COMMITMENT: The metric in pocket α is related to the metric in pocket $\alpha+1$ by $g_{\mu\nu}^{\hat{(\alpha+1)}} = \Lambda^{\hat{\rho}\mu} \Lambda^{\hat{\sigma}\nu} g_{\rho\sigma}^{\hat{(\alpha)}} \times \exp[2(\Phi^{\hat{(\alpha+1)}} - \Phi^{\hat{(\alpha)}})/c^2]$, with the exponential factor accounting for gravitational time dilation between pockets.

OBSERVATIONAL COMMITMENT: GPS satellites are the everyday case: the Earth-to-satellite transformation requires both the SR boost ($v \approx 4$ km/s relative motion) and the GR redshift (Φ difference between Earth's surface and orbit). SCT extends the same Lorentz-boost composition through every nesting level above the GPS satellite frame.

Significance: Establishes the SR-consistent kinematic relation between adjacent levels of the hierarchy.

P10: Hereditary Time Transmission Time is hereditary: each comoving frame inherits its base proper-time behavior from its parent frame and passes a refined version of that time perception to its child objects.

EXPLANATION: Proper time propagates through the nested hierarchy like a recursive function: each frame's baseline clock rate comes from its parent, then gets modified by local motion/gravity before being passed to children.

FORMULA: $d\tau/dt = \prod_{i=1}^k \gamma_i^{-1} \times \prod_{j=1}^k (1 - \Phi_j/c^2)^{1/2}$, for an object at the k-th level of an N-level hierarchy. The proper time rate at any nesting level is the cumulative product of all SR γ factors and all gravitational redshift factors from that level upward through the entire parent chain. MATHEMATICAL COMMITMENT: Proper time $\tau^{(\alpha)}(x,t)$ satisfies recursive relations $\tau^{(\alpha)} = \int d\tau^{(\alpha+1)} \sqrt{1 - \beta^{2(\alpha)}} \times \exp[\Phi^{(\alpha)}/c^2]$ involving parent proper time $\tau^{(\alpha+1)}$, local velocities $\beta^{(\alpha)}$, and gravitational potentials $\Phi^{(\alpha)}$.

OBSERVATIONAL COMMITMENT: Cosmological redshift is produced by metric expansion ($\lambda_{\text{obs}}/\lambda_{\text{emit}} = a_{\text{obs}}/a_{\text{emit}}$); the cumulative hereditary proper-time differences accumulated across nested succession provide a consistency reinterpretation of the same $z(d)$ WITHIN the expanding metric, not a non-metric replacement for it.

P11: Spacetime Pocket as Physical Object Each comoving frame can be treated as a "pocket" of spacetime. The universe is composed of a nested succession of such spacetime pockets.

EXPLANATION: A "pocket" is a gravitationally and kinematically coherent collection of objects sharing approximate comoving motion; it serves as a well-defined organizational unit for the nested hierarchy.

MATHEMATICAL COMMITMENT: Pocket α is defined by characteristic radius $R^{(\alpha)}$, velocity dispersion $\sigma_v^{(\alpha)}$, gravitational binding energy $U^{(\alpha)} = -GM^{2(\alpha)/R}(\alpha)$, and phase-space boundaries in (x, v) space.

OBSERVATIONAL COMMITMENT: Our observable universe is one such pocket with $R^{(obs)} \approx 46.5 \text{ Gly}$, $M^{(obs)} \approx 10^{\{53\}} \text{ kg}$; parent pocket has $R^{(parent)} \gg R^{(obs)}$; siblings and cousins may leave detectable imprints (anisotropies, bulk flows).

NOT this: It is not exotic, it is the natural generalization of the virialized halo in standard cosmology.

P12: Nine Collective Pocket Properties

A pocket has nine measurable collective properties: rotation rate, orbital motion, center of mass, luminosity, gravitational field, magnetic field, electric field, spatial evolution, and inherited time rate. Every pocket inherits its time rate from its parent and passes a refined version to its children.

EXPLANATION: Within each comoving frame, sibling objects “following the leader(s)” have individual velocities relative to that frame and trajectories through gravitational fields, which together refine the inherited perception of time and space that they, in turn, pass to their own child objects in the nested succession. The nine properties are the measurable signatures of the pocket’s collective dynamics.

MATHEMATICAL COMMITMENT: Each property maps to a moment of distributions: angular momentum $L^{(\alpha)} = \int r \times v \, dm$, orbital elements (a, e, i, Ω , ω , M), center of mass $X_{CM} = \int x \, dm / M$, luminosity $L = \int L(x) \, d^3x$, multipole expansion $\Phi(r) = -GM/r + \dots$, field strengths $B^{(\alpha)}$ and $E^{(\alpha)}$, time-dependence $dX_{CM}/dt \neq 0$, and inherited proper-time function $\tau^{(\alpha)}$ (history).

OBSERVATIONAL COMMITMENT: Properties (A)-(H) must match observed characteristics of astronomical structures at each scale; property (I), inherited time rate, must explain hereditary time and redshift mechanisms.

Significance: Enumerates the observable/measurable quantities that characterize a pocket as a physical entity within its parent frame.

P13: Observable Universe as a Pocket

Our observable universe is a spacetime pocket: finite in extent, bounded by our light cone, embedded in the parent pocket structure that extends beyond our horizon.

EXPLANATION: Our observable patch (radius ~ 46.5 Gly, mass $\sim 10^{53}$ kg) is one specific instance of the general pocket construct of P11. The parent pocket structure continues beyond the observable horizon and provides the gravitational mesh whose dissipation we interpret as dark energy.

MATHEMATICAL COMMITMENT: For our observable universe ($\alpha = 0$): $R^{(0)} \approx 46.5$ Gly, $M^{(0)} \approx 10^{53}$ kg, $\sigma_v^{(0)} \sim 300$ km/s. Parent pocket ($\alpha = +1$) must satisfy $R^{(+1)} \gg R^{(0)}$, with quantitative scaling determined from observational constraints (bulk flows, CMB dipole).

OBSERVATIONAL COMMITMENT: Phenomena at our observational horizon must be interpretable as local features of a larger structure, not fundamental cosmic boundaries. Bulk flows, CMB dipole and large-angle anomalies, and the KBC supervoid are the primary signatures of our pocket's embedding in the parent structure.

Significance: Identifies our observable patch as a specific instance of the general pocket construct, with parent pocket structure continuing beyond the horizon.

CATEGORY IV: THE NATURE OF DARK ENERGY (P14-P19)

P14: Orbital Decay and Gravitational Mesh Dissipation No orbit contains infinite energy; all orbits decay over time. Some decay inward, most decay outward. As a result, the distances between objects at each level of the nested succession change, leading to a dissipation in the average strength of the overlapping gravitational wells that define that frame. EXPLANATION: All gravitationally bound orbits lose energy through radiation (gravitational waves, tidal friction, electromagnetic drag); two-body orbits predominantly decay outward due to three-body interactions and dynamical friction, increasing inter-object separations.

MATHEMATICAL COMMITMENT: Gravitational binding energy $U^{(\alpha)} = -\sum_i \sum_{\{j>i\}} G m_i m_j / r_{\{ij\}}$ evolves as $dU^{(\alpha)}/dt > 0$ (becoming less negative), implying increasing separations $dr_{\{ij\}}/dt > 0$ statistically; overlapping gravitational potential $\Phi_{\text{mesh}} = \sum_i \Phi_i$ decreases in magnitude as $|d\Phi_{\text{mesh}}/dt| < 0$.

OBSERVATIONAL COMMITMENT: Must reproduce observed acceleration parameter $q_0 \approx -0.55$ without invoking vacuum energy; decay timescales must be consistent with observed galaxy cluster evolution and large-scale structure growth.

NOT this: This is not speculative, it is established long-term behavior of N-body gravitational systems under GR.

P15: Apparent Expansion from Parent-Frame Mesh Change When this dissipation occurs across a nested succession of parent comoving frames, each providing a base perception of space and time to its child objects, observers at our scale factor would interpret the net effect as if the “fabric of spacetime” were being pulled apart.

EXPLANATION: Because each frame inherits its baseline proper-time and spatial metric from parent frames (P10), dissipation of parent-frame gravitational mesh (P14) propagates to child frames as an apparent stretching of space.

MATHEMATICAL COMMITMENT: Effective scale factor $a_{\text{eff}}(t)$ emerges from cumulative dissipation across parent succession: $da_{\text{eff}}/dt \propto \sum_{\alpha=\text{obs}}^{\infty} (d\Phi_{\text{mesh}}^{\alpha}/dt)$, where sum extends over all parent frames influencing our pocket.

OBSERVATIONAL COMMITMENT: Must reproduce Hubble law $v = H_0 d$ for nearby galaxies without invoking FLRW metric expansion; effective Hubble parameter $H_{\text{eff}}(z)$ must match observations.

P16: Mesh Dissipation Equivalence to Observed Expansion Dark energy is not vacuum energy. Instead, it is related to the dissipation of the average gravitational tensor “mesh strength” across a nested succession of parent comoving frames of reference.

EXPLANATION: What Λ CDM attributes to constant vacuum energy density ρ_{Λ} , SCT reinterprets as time-varying weakening of the cumulative gravitational field network created by parent-frame mass distributions.

MATHEMATICAL COMMITMENT: Energy density ρ_{DE} emerges from second time derivative of mesh potential: $\rho_{\text{DE}} \propto \partial^2\Phi_{\text{mesh}}/\partial t^2$ rather than constant vacuum contribution; Einstein field equations acquire effective cosmological term $\Lambda_{\text{eff}} g_{\mu\nu} = (8\pi G/c^4) \rho_{\text{DE}} g_{\mu\nu}$.

OBSERVATIONAL COMMITMENT: Λ_{eff} must be spatially and temporally variable, not constant; variations must be consistent with observed expansion history and structure formation.

NOT this: Dark energy is not a physical substance filling empty space.

P17: Dynamical Cosmological Ratio Therefore, the cosmological constant Λ in the Einstein field equations should be interpreted as a ratio between: (i) the localized strength of overlapping gravitational wells within a given pocket, and (ii) the cumulative influence of the succession of parent frames whose gravitational environments they are competing against.

EXPLANATION: Λ_{eff} quantifies the balance between local gravitational binding (resisting expansion) and parent-frame dissipation (driving apparent expansion); as parent frames decay, local binding becomes relatively weaker, increasing Λ_{eff} .

FORMULA: $\Lambda_{\text{eff}}(x,t) = \kappa \times [U_{\text{parent}}(x,t) / U_{\text{local}}(x,t)]$, where κ has units m^{-2} and is calibrated to reproduce the observed $\Lambda_{\text{obs}} \approx 1.1 \times 10^{-52} \text{m}^{-2}$ when spatially averaged. MATHEMATICAL COMMITMENT: $\Lambda_{\text{eff}}(x,t) = \kappa [U_{\text{parent}}(x,t) / U_{\text{local}}(x,t)]$, where κ is a dimensioned constant $[\kappa] = \text{L}^{-2}$, U_{local} represents local gravitational binding energy or potential strength within pocket α , and U_{parent} represents cumulative parent-frame binding energy from succession $\alpha+1, \alpha+2, \dots$

OBSERVATIONAL COMMITMENT: Spatial variations in Λ_{eff} must correlate with observed bulk flows and large-scale velocity fields; temporal evolution must account for part of the Hubble tension by explaining different H_0 values at different epochs.

NOT this: Not a fixed universal constant with no identified physical origin.

P18: Long-Term Exponential Cascade Over long enough time frames, the aggregated effects of dark energy being due to dissipations in the gravitational tensor mesh strength of a nested succession of parent pockets of spacetime would predict a rate of increase across large scales to increase exponentially as time goes on.

EXPLANATION: As the parent-frame mesh dissipates (P14), the cumulative inherited dissipation U_{parent} increases monotonically; if U_{local} remains approximately constant, $\Lambda_{\text{eff}} \propto U_{\text{parent}}$ grows; continued dissipation causes accelerating growth, yielding exponential behavior at long times.

MATHEMATICAL COMMITMENT: Asymptotic behavior $d^2a_{\text{eff}}/dt^2 \propto \exp(t/\tau_{\text{decay}})$ for some decay timescale $\tau_{\text{decay}} \gg t_{\text{universe}} \approx 13.8$ Gyr, implying equation of state parameter $w(t \rightarrow \infty) \rightarrow -\infty$ (phantom dark energy regime) (this is the asymptotic far-future $t \rightarrow \infty$ limit; the present-day spatially-averaged equation of state is the thawing $w_0 = -0.90 \pm 0.03$, $w_a = -0.19 \pm 0.07$ of Paper 10, in the DESI DR2-preferred direction).

OBSERVATIONAL COMMITMENT: SCT adopts the quintessence-like, DESI-consistent equation of state of the companion Paper 10 ($w_0 = -0.90 \pm 0.03$, $w_a = -0.19 \pm 0.07$), i.e. $w > -1$ today with a thawing trend, which is what DESI DR2 (2025) prefers (dynamical-dark-energy preference now $\sim 2.8\text{-}4.2\sigma$, with phantom behavior at high z crossing to $w > -1$ at low z). The §VI.D long-term exponential-growth scenario predicting $w < -1$ at low redshift ($z < 0.5$) is therefore currently DISFAVORED by data and is retained only as a CONJECTURAL/OPEN late-time limit; it is NOT the form this paper adopts. Future observations (DESI, Euclid, Roman) test the adopted (w_0 , w_a) thawing form.

P19: Short-Term Local Variability (Hubble Tension) Since Lambda now becomes a ratio and is variable in different parts of space across different moments in time, although the average tendency over time will move towards an exponentially increasing rate, this model allows for temporary instances where the rate could appear to slow down under certain conditions when either the localized overlapping wells are stronger than average, or the most direct parent structures are moving towards each

other and not away from each other. EXPLANATION: Because $\Lambda_{\text{eff}} = \kappa U_{\text{parent}}/U_{\text{local}}$ (P17), fluctuations in either the parent dissipation U_{parent} or the local mesh U_{local} (local gravitational clustering increasing U_{local} , which suppresses Λ_{eff}) cause spatial and temporal variations in apparent expansion rate.

MATHEMATICAL COMMITMENT: $\Lambda_{\text{eff}}(x,t)$ exhibits fractional fluctuations $\Delta\Lambda/\Lambda \sim O(0.01-0.1)$ on timescales $\Delta t \sim \text{Gyr}$ and spatial scales $\Delta x \sim 100 \text{ Mpc}$, with correlation structure $\langle \Delta\Lambda(x_1,t_1) \Delta\Lambda(x_2,t_2) \rangle$ determined by parent-frame dynamics.

OBSERVATIONAL COMMITMENT: Explains Hubble tension as temporal variation: early universe (CMB decoupling, $z \approx 1100$) had different Λ_{eff} than late universe (local H_0 measurements, $z < 0.1$), yielding $H_{0,\text{CMB}} \approx 67$ vs $H_{0,\text{local}} \approx 73$; also predicts potential variability in dark energy equation of state $w(z)$ with specific spatial patterns.

NOT this: No new particles or modified gravity is required.

CATEGORY V: ORIGIN OF OUR VISIBLE UNIVERSE (P20-P49)

P20: Local Character of the SR Speed Limit The speed of light c sets the maximum attainable speed for objects accelerated within our local nested succession of parent frames of reference.

EXPLANATION: Standard Special Relativity applies to all particles and objects accelerated from rest within our observable universe or its immediate parent frames; no local process can produce superluminal velocities.

MATHEMATICAL COMMITMENT: For objects accelerated from rest within frame α via local forces (gravity, electromagnetic, nuclear), relativistic

momentum $p = \gamma m v$ approaches pc as kinetic energy $\rightarrow \infty$, ensuring $v_{\text{max}} \rightarrow c$ asymptotically.

OBSERVATIONAL COMMITMENT: All observed particle velocities, including galactic jets, cosmic rays, and stellar velocities must satisfy $v_{\text{local}} \leq c$; no local process violates SR.

NOT this: This distinction is not a loophole, it is the definition of what SR claims and what it does not.

P21: Accepted Precedent in Standard Cosmology

Galaxies beyond the Hubble radius of approximately 14.4 gigalight-years recede at velocities exceeding c under the standard expansion model. This is universally understood not to violate Special Relativity. SCT extends this same accepted reasoning to the collision scenario: superluminal relative velocities between independently formed pockets are the generic expectation for pockets separated by distances exceeding $c/H_0 \approx 14.4$ gigalight-years.

EXPLANATION: The cosmology community already accepts that distant galaxies recede faster than c without invoking new physics, this is standard Λ CDM, where $v_{\text{recession}} = H_0 \cdot d$ can exceed c at $d > c/H_0$. SCT applies precisely the same reasoning to relative velocities between independently formed comoving pockets at vastly larger scales.

MATHEMATICAL COMMITMENT: Two pockets α and β with independent formation histories, separated by $d > c/H_0 \approx 14.4$ Gly, can have relative velocity $v_{\text{rel}} > c$ without violation of Special Relativity, because each pocket's local SR remains valid; superluminal v_{rel} is a consequence of separation and independent kinematics, not of local acceleration.

OBSERVATIONAL COMMITMENT: This premise grounds the SCT collision mechanism in a precedent already accepted in mainstream cosmology. No new physics is invoked; what is invoked is the recognition that the same kinematic permissiveness that Λ CDM allows for distant-galaxy recession also applies to inter-pocket collision velocities.

Significance: Demonstrates that the key SCT claim of superluminal inter-pocket relative velocities has an established precedent already accepted in mainstream cosmology.

P22: Physics of Superluminal Intersections No laws of physics are violated when two immense nested structures of comoving frames intersect the same region of spacetime with relative speeds exceeding twice the speed of light.

EXPLANATION: Intersection of two independently moving structures at $v_{\text{rel}} > 2c$ involves no causality violation, signal propagation faster than c , or SR violation in either structure's local frame; it is a kinematic possibility in GR.

MATHEMATICAL COMMITMENT: Two metrics $g_{\mu\nu}^{(\alpha)}$ and $g_{\mu\nu}^{(\beta)}$ with different coordinate systems can intersect same spacetime region (x, t) with relative velocity $v_{\text{rel}} = |v^{(\alpha)} - v^{(\beta)}| > 2c$; each metric separately satisfies Einstein equations and local SR; no unified "speed of light" limits relative velocities.

OBSERVATIONAL COMMITMENT: Collision physics must be worked out using shock dynamics and thermalization in the overlap region; such collisions are not forbidden a priori by fundamental physics.

NOT this: No energy condition is violated, the kinetic energy is real classical kinetic energy in the bulk motion of massive pockets, not vacuum energy or any exotic source.

P23: Collision Energy Regime Superluminal collisions between such immense structures can occur with kinetic energies far beyond twice the speed of light, potentially many times (e.g., 3, 7, 42, 67 c), introducing physical regimes and states of matter that are currently beyond our ability to simulate.

EXPLANATION: If two pockets of mass $M \sim 10^{\{53\}}$ kg collide at $v_{rel} \sim 10c$, the collision energy $E \sim (1/2) M v_{rel}^2$ exceeds rest mass energy Mc^2 by factors ~ 50 , creating extreme high-energy-density states (quark-gluon plasmas, exotic hadrons, possibly non-equilibrium QCD matter).

MATHEMATICAL COMMITMENT: Collision energy $E_{collision} = (1/2) M_{pocket} v_{rel}^2$ with $v_{rel} = v_{pocket} \times \gamma_{boost}$ can exceed Mc^2 by arbitrarily large factors; energy density $\varepsilon = E_{collision} / V_{collision}$ determines thermalization temperature and equation of state in collision region.

OBSERVATIONAL COMMITMENT: Exotic intermediate states during collision-thermalization must ultimately cool to reproduce observed CMB temperature (2.725 K), BBN abundances, and structure formation; models must address thermalization timescales and entropy evolution.

P24: Single Assumption Change Resolving Seven Mysteries When we replace the assumption of a single hot dense origin with a succession of superluminal collisions, many of the mysteries that current models and simulations struggle with can be addressed by this change in assumption.

EXPLANATION: Replacing Λ CDM's Planck-scale singularity with multi-stage collision thermalization provides alternative origin for: CMB homogeneity, BBN abundances, structure formation timescales, particle-antiparticle asymmetry, and primordial perturbations.

MATHEMATICAL COMMITMENT: No new commitment; this premise identifies the conceptual shift. Specific mechanisms are addressed in subsequent premises P25-P41.

OBSERVATIONAL COMMITMENT: SCT must reproduce or improve upon Λ CDM's fits to CMB power spectrum, large-scale structure, and BBN abundances while resolving tensions (H_0 , early galaxies, etc.); if not, the collision assumption fails.

P25: Pre-existing Matter Thermalized by Collision These collisions provide both: (i) the catalyst that pulverizes existing mass into hot, dense, swirling plasma, and (ii) the pre-existing conditions that precede what we call "our Big Bang."

EXPLANATION: Superluminal collision creates shock-heated thermalized plasma from pre-existing matter; this plasma, once cooled and evolved, appears phenomenologically identical to Λ CDM's "Big Bang" conditions at very early times.

MATHEMATICAL COMMITMENT: Initial conditions for our observable universe are set by collision shock-front conditions at $t = t_{\text{collision}}$: temperature T_{shock} , energy density $\varepsilon_{\text{shock}}$, particle spectra, turbulence, magnetic fields; these become starting point for standard thermalization and nucleosynthesis.

OBSERVATIONAL COMMITMENT: CMB and BBN must emerge from collision thermalization; deviations from Λ CDM predictions in primordial spectra or isotope abundances could discriminate models.

NOT this: The collision did not create matter from nothing.

P26: Local Big Bang Our Big Bang did not create the entire universe; it created only an infinitesimal patch of spacetime relative to the larger structure.

EXPLANATION: The collision event creating our observable universe represents a local thermalization and structure-formation episode within the infinite universe; it is not a cosmic beginning.

MATHEMATICAL COMMITMENT: Observable universe volume $V_{\text{obs}} \sim (46.5 \text{ Gly})^3$ is infinitesimal compared to parent pocket volume $V_{\text{parent}} \gg V_{\text{obs}}$; extent of collision thermalization $V_{\text{collision}} \gg V_{\text{obs}}$ but $\ll V_{\text{parent}}$.

OBSERVATIONAL COMMITMENT: No observational implication; this is a boundary- condition statement establishing our local cosmic location.

NOT this: Beyond our observable horizon SCT predicts not empty space but continuation of parent pocket structure.

P27: Infinite Array of Big Bang Events Our Big Bang is one of an infinite number of Big Bang-like events that have occurred eternally.

EXPLANATION: The eternal universe contains infinite opportunities for pocket collisions at all epochs and locations; our collision is one of infinitely many similar thermalization events.

MATHEMATICAL COMMITMENT: Collision rate per comoving volume $\Gamma_{\text{collision}}(t) \propto \text{collision cross-section} \times \text{relative velocity density}$; integrated over all space and eternal time, total number of collision events $\rightarrow \infty$.

OBSERVATIONAL COMMITMENT: Potential detection of external collision signatures (transients, CMB anomalies at horizon, relic radiation from distant collisions) would provide indirect evidence.

P28: Eternal Collision Cycle There was no first collision and there will be no last collision. In an infinite universe, there is an eternal cycle of creation and slow dissipation, followed by new sequences of collisions.

EXPLANATION: Eternal time (P1) and infinite space (P2) ensure continuous collision activity throughout the universe; no temporal beginning or end to the collision cascade.

MATHEMATICAL COMMITMENT: No cosmological “arrow of time” from initial singularity; entropy increases locally through dissipation (P14), but global state is eternal and unchanging in character (though varying in spatial details).

OBSERVATIONAL COMMITMENT: No observational test; this is a boundary-condition statement about temporal structure of the universe.

P29: Thermodynamic State Sufficiency

Once a photon-baryon plasma thermalizes, its subsequent acoustic behavior is determined entirely by its thermodynamic state at decoupling, temperature, density, baryon-to-photon ratio, and the spectrum of density perturbations. The plasma has no memory of whether it was created by a singular origin or by a cascade of superluminal collisions. Two plasmas arriving at the same thermodynamic state by different paths produce acoustically identical CMB power spectra.

EXPLANATION: This is the Plasma Equivalence Proposition (an argument, not a rigorous theorem; the assumptions below are stated as assumptions), a consequence of the tight-coupling approximation in the photon-baryon Boltzmann hierarchy. The acoustic behavior of the plasma depends only on its thermodynamic state, not on its formation history. Whatever set the initial thermodynamic state, a singular Big Bang or a multi-stage collision cascade, the post-decoupling evolution is identical.

MATHEMATICAL COMMITMENT: The Boltzmann hierarchy for the photon-baryon plasma, $df/dt + \hat{p} \cdot \nabla f - H p (df/dp) = C[f]$, depends on the initial state $\{T_{\text{dec}}, \eta, Y_p, \tau_{\text{reion}}, k_{\text{eq}}, r_s\}$ and not on the production history of that state.

NOT this: This is not an assumption, it is a consequence of the tight-coupling approximation.

OBSERVATIONAL COMMITMENT: SCT's collision-origin plasma reproduces the observed CMB scalar power spectrum to within the perturbative corrections ($\sim 10^{-5}$) introduced by gravitational superposition at recombination. No distinguishing scalar signatures are expected at sub-horizon scales.

Significance: Explains why SCT's collision-origin plasma can reproduce the observed CMB scalar power spectrum at sub-horizon scales without distinguishing signatures.

P30: Six Thermodynamic State Parameters Determine CMB

The CMB power spectrum for multipoles $\ell > 30$ is determined by the thermodynamic state at decoupling TOGETHER WITH the primordial scalar perturbation spectrum and the geometry/distance to last scattering: $\{T_{\text{dec}}, \eta, Y_p, \tau_{\text{reion}}, k_{\text{eq}}, r_s\}$ plus the scalar amplitude A_s , the tilt n_s (i.e. the

full $P(k)$), and the angular-diameter distance D_A to last scattering, regardless of origin mechanism. (The earlier statement that six thermodynamic parameters alone suffice was incomplete: A_s , n_s , and D_A are essential to the observed C_ℓ .) There is one categorical exception to this erasure: angular momentum is a conserved vector quantity protected by an exact symmetry, Noether's theorem guarantees it cannot be destroyed by thermalization, only redistributed. The collision history is therefore erased from the CMB scalar spectrum but written into the rotation of every structure that forms from the post-collision plasma.

EXPLANATION: The six-parameter family describes everything about the photon-baryon plasma's acoustic behavior at sub-horizon scales after decoupling. All collision-cascade history is erased except for the angular momentum imprint, which Noether's theorem protects. The angular momentum signature lives in galactic rotation, cluster spin, satellite-plane co-rotation, and quasar polarization alignment, not in the CMB scalar spectrum.

FORMULA: $\{T_{\text{dec}}, \eta, Y_p, \tau_{\text{reion}}, k_{\text{eq}}, r_s\}$

OBSERVATIONAL COMMITMENT: The CMB scalar power spectrum is the convergence point, both Λ CDM and SCT must reproduce it. The angular momentum signature is the divergence point: SCT predicts coherent inheritance of J at all scales, Λ CDM predicts incoherent tidal-torque accumulation at $< 30\text{-}50$ Mpc.

Significance: Identifies the angular momentum signature as the primary observational discriminant between SCT and Λ CDM.

P31: Grazing Collisions and Flat Rotation Curves

When two pockets collide with nonzero impact parameter b , angular momentum $J = \mu(b \times v_{\text{rel}})$ is deposited into the overlap volume, where μ is the reduced mass of the colliding system. The inherited specific angular momentum $j = J/M$ sets the centrifugal barrier for gravitational collapse. A centrifugal barrier in a self-gravitating system is ASSOCIATED with an isothermal-like density profile $\rho(r) \propto r^{-2}$, the profile that generates flat rotation curves. CAVEAT: a centrifugal barrier alone does not entail a global r^{-2} (singular isothermal sphere) profile, which requires a specific thermodynamic/dynamical equilibrium; the step from J -deposition to a global r^{-2} profile is asserted here and derived in the companion rotation-curve paper, not in this section.

EXPLANATION: A grazing collision deposits J into the overlap volume; the centrifugal barrier this creates halts collapse at a finite radius and forces isothermal density distribution. $\rho(r) \propto r^{-2}$ gives $M(<r) \propto r$, which gives $v_{\text{circ}}^2 = GM(<r)/r \approx \text{const}$, exactly flat rotation curves. No dark matter halo required.

FORMULA: $J = \mu(b \times v_{\text{rel}})$; $j = J/M$; $\rho(r) \propto r^{-2}$

NOT this: This is not circular reasoning, the centrifugal barrier emerges from collision geometry, not from a tuned dark-matter halo profile.

OBSERVATIONAL COMMITMENT: Flat rotation curves observed across the SPARC galaxy sample (~ 150 disk galaxies), satisfied without invoking particle dark matter. Tully-Fisher scatter reflects collision-determined angular momentum; rotation transition radius $r_{\text{transition}} = 3 R_d$ (confirmed across 24 SPARC galaxies at slope 2.908 ± 0.024 , 0.09σ).

Significance: No dark matter particles are required to seed the isothermal profile, the collision geometry provides it directly.

P32: Angular Momentum Inheritance Principle

Angular momentum conservation operates simultaneously at every level of the nested hierarchy. When structures at any scale condense from rotating material at the scale above, they inherit a fraction of the parent's angular momentum proportional to their mass fraction and position. This is associated with the empirical scaling relation $J \propto M^{5/3}$, equivalently $j = J/M \propto M^{2/3}$, observed across seven decades of scale from planetary systems to supercluster complexes. NOTE: within this paper $J \propto M^{5/3}$ is an empirical scaling pending the inheritance-dynamics derivation in Paper 5; it is not derived here. (The Tang et al. 2025 cluster-rotation and Tudorache et al. 2025 filament-rotation significances are load-bearing and should be treated as provisional pending independent confirmation.) Produces coherent spin alignment across all scales simultaneously. Observed alignments extending to 30-100 megaparsec separations exceed tidal-torque-theory predictions by factors of 10-20.

EXPLANATION: SCT explains the universal $J \propto M^{5/3}$ scaling as natural consequences of inheritance from a common collision origin, rather than coincidental local torquing across all scales independently. Cluster rotation velocity scales 360 km/s at $10^{14} M_{\odot}$ rising to 693 km/s at $10^{15} M_{\odot}$ (Tang et al. 2025, $> 100 \sigma$ aggregate). Filament bulk rotation ~ 110 km/s observed individually (Tudorache et al. 2025).

FORMULA: $J \propto M^{5/3} \iff j = J/M \propto M^{2/3}$

NOT this: Not reproduced as an assumed universal scaling law, derived from inheritance dynamics across the hierarchy.

OBSERVATIONAL COMMITMENT: Verified across seven decades of scale (planetary systems \rightarrow galaxies \rightarrow clusters \rightarrow superclusters); galactic spin

alignment extending to 30-100 Mpc; co-rotating satellite planes around six independent host systems (joint $P \approx 2 \times 10^{-14}$); BCG-cluster shape alignment fully in place at $z > 1.3$ (Smith et al. 2023).

Significance: SCT explains the observed alignments as natural consequences of inheritance from a common collision origin, rather than coincidental local torquing.

P33: Head-On Collisions and Filament Formation

Near-zero impact parameter collisions convert kinetic energy primarily into heat and compression along the collision axis, with negligible retained angular momentum. Matter collapses freely in the perpendicular directions, producing elongated high-density filaments. Filament length scales with the combined pocket extent along the collision axis; filament width scales with the smaller pocket's self-gravity.

EXPLANATION: Direct collisions ($b \approx 0$) deposit $J \approx 0$; without a centrifugal barrier, perpendicular collapse produces filamentary morphology. The cosmic web's filaments are direct collision-geometry signatures rather than dark-matter scaffolding.

MATHEMATICAL COMMITMENT: For $b \rightarrow 0$, $J/M \rightarrow 0$; filament length L_f scales with combined pocket extent along collision axis ($L_f \propto R_{\text{pocket}(1)} + R_{\text{pocket}(2)}$); filament width W_f scales with the smaller pocket's self-gravity ($W_f \sim \min(R_{\text{pocket}(1)}, R_{\text{pocket}(2)})$).

OBSERVATIONAL COMMITMENT: Cosmic-web filament length distribution (50-100 Mpc typical, up to 400 Mpc for the Quipu superfilament); coherent rotation of largest filaments (~ 15 Mpc filaments with aligned angular momentum, MeerKAT 21-cm); spin-filament alignment ($|\cos \psi| \approx 0.75 \pm 0.05$ from collision-axis inheritance).

Significance: Identifies the mechanism that produces the filamentary skeleton of the cosmic web.

P34: Full Cosmic Web from Collision Geometry Distribution

The full cosmic web emerges from the full parameter space of collision geometries: grazing collisions (geometrically more probable, since $P(b) \propto b$) produce rotating halos; near-head-on collisions produce filaments and walls; collision nodes where filaments of different orientations intersect produce the most massive clusters. The scale distribution of structures mirrors the scale-invariant hierarchy: grandparent-scale collisions produce gigaparsec superfilaments; parent-scale collisions produce 100-megaparsec filaments; sibling-scale collisions produce 10-50 megaparsec structures.

EXPLANATION: The cosmic web's filament-wall-void-cluster topology emerges naturally from sampling the collision impact-parameter distribution $P(b) \propto b$ across multiple hierarchy levels. Λ CDM requires dark matter scaffolding to seed the same morphology; SCT requires only collision geometry.

FORMULA: $P(b) \propto b$

NOT this: No dark matter scaffolding is needed.

OBSERVATIONAL COMMITMENT: Reproduces SDSS, 2dFGRS, BOSS, DESI cosmic-web topology including filament lengths, void sizes, and cluster spatial distribution. Big Ring (~ 1.3 Gly diameter, $z \approx 0.8$) and Giant Arc (~ 3.3 Gly, $z \approx 0.8$) emerge as gigaparsec-scale collision-geometry signatures ($\Lambda_{\text{max}} \approx 5$ Gpc).

Significance: The geometry emerges directly from collision products with no dark matter particle scaffolding.

P35: Recombination Epoch with Collision Signatures

SCT's collision-thermalized plasma produces a recombination epoch that differs from Λ CDM in subtle but in-principle detectable ways. The collision geometry imprints density variations across the overlap volume: high-density regions (recombination rate $\propto n^2$) recombine faster; low-density peripheral regions recombine slower. The sky-averaged recombination redshift of approximately $z = 1100$ is preserved by the thermodynamic state parameters of P30. The SCT-specific signatures are a compressed recombination epoch duration and a line-of-sight dispersion in the decoupling redshift, producing specific non-Gaussian CMB contributions at angular scales corresponding to the collision region size.

EXPLANATION: The Plasma Equivalence Proposition (P29 + P30; an argument under the stated tight-coupling assumptions, not a rigorous theorem) preserves the scalar power spectrum at $\ell > 30$. The deviations from Λ CDM live at low ℓ (collision-axis imprints, Cold Spot, hemispherical asymmetry, axis of evil) and in non-Gaussian statistics at angular scales corresponding to the collision-region size.

MATHEMATICAL COMMITMENT: The recombination rate is $\propto n_e^2$ where $n_e = n_{\text{baryon}} + n_{\text{electron}}$; the collision overlap volume's density structure modulates the local recombination rate, producing $\Delta T/T$ deviations of order 10^{-5} - 10^{-4} at angular scales $\sim 10^\circ$ corresponding to the collision-region size.

OBSERVATIONAL COMMITMENT: The CMB Cold Spot (3σ anomaly), the Axis of Evil (2.8σ), the hemispherical power asymmetry (2.5σ), and the low- ℓ power deficit ($\sim 2 \sigma$) are all SCT-predicted collision-axis imprints. The same axis should emerge from independent measurements (CMB-S4 polarization, Simons Observatory).

Significance: Predicts detectable, non-Gaussian CMB signatures tied to the collision region geometry.

P36: Multi-Stage Cascade Initiation

The initial superluminal collision produces a non-equilibrium plasma retaining bulk kinetic energy as turbulence and large-scale velocity gradients. Successive collisions among daughters at this stage involve comparable kinetic energies and similar geometric structure, distributing heat over the overlap volume in a way consistent with the observed CMB's small-amplitude anisotropy.

EXPLANATION: The cascade does not terminate at the initial collision. Daughter fragments retain superluminal velocities and undergo secondary collisions (P37) until the system has cooled and decohered enough that v_{rel} drops below c (P38). The multi-stage cascade is what distributes heat over the overlap volume rather than concentrating it at a single point.

MATHEMATICAL COMMITMENT: Each cascade stage releases energy $E_n \sim E_{n-1}$ within a factor ~ 2 (P29); the volume over which heat is distributed scales as $V_n \sim R_{pocket(n)}^3$; the resulting baseline temperature $T_{CMB} \sim (E_{total} / V_{total} / a_{radiation})^{1/4} \approx 2.725$ K is preserved across the cascade.

OBSERVATIONAL COMMITMENT: The cumulative effect of multi-stage cascade thermalization, not a single instantaneous shock, accounts for the observed CMB temperature uniformity to $\Delta T/T \lesssim 10^{-5}$ across causally-disconnected regions, without requiring inflation.

Significance: Establishes that the initial thermalization event is not a single clean shock but initiates a multi-stage energy-dissipation process.

P37: Secondary Collisions from Daughter Fragments

Daughter fragments from the first stage are still moving at potentially superluminal relative velocities, producing secondary collisions.

EXPLANATION: The first cascade stage deposits matter into a turbulent, non-equilibrium plasma whose constituent daughter fragments retain bulk velocities of order $v_{rel}(0) < c$ (P22's overlap-volume thermalization is local; the bulk fragments retain pre-collision velocities). These secondary collisions among daughters extend the cascade through additional stages.

MATHEMATICAL COMMITMENT: For a superluminal initial collision at $v_{rel}(0) \sim 10c$, daughter fragments retain bulk velocities of order $v_{rel}(1) \sim v_{rel}(0)/\eta_{collision}$ where $\eta_{collision}$ is the kinetic-to-thermal conversion efficiency at the first stage; if $\eta_{collision} \ll 1$, as expected for superluminal collisions where thermalization is incomplete, daughter velocities can remain superluminal for several cascade stages.

OBSERVATIONAL COMMITMENT: The number of cascade stages N_{stage} is not directly observable from internal observables; SCT does not claim a specific N_{stage} but does claim $N_{stage} > 1$ with observational consequences distinct from a single-stage cascade.

Significance: Extends the cascade mechanism through successive stages.

P38: Cascade Termination at $v < c$

The cascade continues, each stage dissipating some fraction of remaining kinetic energy into heat, until relative velocities drop below c . SCT does not claim to know the precise number of stages; it claims only that more than one occurred, that their cumulative effects differ

observationally from a single stage, and that this difference is in principle detectable.

EXPLANATION: Each cascade stage dissipates kinetic energy into thermal energy. As thermal energy accumulates and cooling progresses, the surviving daughter fragments' bulk velocities decrease. Eventually $v_{\text{rel}}(n) < c$ and superluminal cascade dynamics terminate. From that point onward, the universe evolves under standard subluminal physics from the initial conditions the cascade produced.

MATHEMATICAL COMMITMENT: The cascade-termination condition $v_{\text{rel}}(n_{\text{term}}) < c$ determines the number of stages n_{term} self-consistently from the dissipation per stage. SCT does not predict a specific n_{term} value, but sets a hard upper bound: n_{term} must be small enough that the cascade has terminated before BBN ($t < 1$ s, $T < 10^9$ K).

NOT this: SCT does not claim to know the precise number of stages.

OBSERVATIONAL COMMITMENT: The cumulative effects of multi-stage cascade differ observationally from a single-stage cascade in: (1) CMB low- ℓ multipole structure (collision-axis imprints; P64); (2) primordial non-Gaussianity at $f_{\text{NL}} \sim 10^{-2}$; (3) BBN abundance ratios (the pre-BBN cascade termination must not perturb D/H, Y_{p} , or ${}^3\text{He}/\text{H}$ beyond observed precision).

Significance: Establishes the termination condition of the cascade, subluminal relative velocities, and the observational consequence of multi-stage dissipation.

P39: Quark-Gluon Plasma Phase at $T > T_{\text{QCD}}$

During the highest-energy phases of the cascade, temperatures exceed $T_{\text{QCD}} \approx 1.7 \times 10^{12}$ K, placing the plasma into a quark-gluon plasma phase governed by QCD at finite baryon density.

EXPLANATION: The kinetic energy deposited at the early cascade stages produces a plasma temperature far above the QCD scale. The plasma is then in a quark-gluon plasma phase, governed by QCD at finite baryon density, before cooling through the QCD transition into the baryon-photon-electron plasma that subsequently produces BBN and recombination.

MATHEMATICAL COMMITMENT: For a pocket of mass $M \approx 10^{53}$ kg colliding at $v_{\text{rel}} \sim 10c$, the kinetic energy density at the first stage corresponds to temperatures reaching $T_{\text{QCD}} \approx 1.7 \times 10^{12}$ K and potentially the electroweak scale $T_{\text{EW}} \approx 10^{15}$ K in compressed hotspot regions. The QGP phase is governed by lattice QCD at finite baryon chemical potential; the equation of state is calibrated against RHIC and LHC heavy-ion-collision results.

OBSERVATIONAL COMMITMENT: The QGP phase places the early cascade in the same physical regime as heavy-ion collision experiments at CERN and RHIC. There is no requirement for new physics; the QGP equation of state at finite baryon density is the relevant standard-model calculation.

Significance: Connects cascade microphysics to the established Standard Model QCD regime studied in heavy-ion collision experiments at CERN and RHIC.

P40: Cascade Termination Before $t \approx 1$ Second

Three independent observational constraints establish that the entire cascade terminated before $t \approx 1$ second after the effective Big Bang, far before recombination:

(1) BBN abundance constraints ($D/H = 2.527 \pm 0.030 \times 10^{-5}$, $Y_p = 0.2449 \pm 0.0040$) require thermal equilibrium weak interactions at $t \approx 1$ second with no active collision energy injection;

(2) COBE/FIRAS spectral purity ($|y| < 1.5 \times 10^{-5}$) requires all non-standard energy injection to have concluded before $z \approx 5 \times 10^4$;

(3) Planck 2018 acoustic peak positions require no perturbation sources between the end of the cascade and recombination at $z \approx 1100$.

All three constraints independently require cascade termination at $t <$ approximately 1 second.

EXPLANATION: After cascade termination, the universe evolves under standard physics from the initial conditions the cascade produced. The exotic creation physics is ancient relative to everything we can directly observe.

MATHEMATICAL COMMITMENT: Cascade-termination redshift $z_{\text{cascade}} > 5 \times 10^4$ from FIRAS; cascade-termination time $t_{\text{cascade}} < 1$ s from BBN.

OBSERVATIONAL COMMITMENT: The observed primordial element abundances (D/H , Y_p , ${}^3\text{He}/H$, Li-7) match standard BBN predictions, confirming that no significant perturbation sources operated between the cascade and recombination. Spectral purity of CMB blackbody is preserved to FIRAS bounds.

Significance: After this point the universe evolves under standard physics from the initial conditions the cascade produced. The exotic creation physics is ancient relative to everything we can directly observe.

FTL Transit Signatures (observational note): SCT permits rare external objects from outside our visible patch to traverse our region at superluminal velocities, typically the residual ejecta from sibling-pocket collisions in our shared parent frame (P58, P59, P60). Such traversals would produce optical transients whose phenomenology depends on the impacted material: passage through nebular gas should ignite fast cooling (blue-optical transient signatures, FBOT class) or slow diffusive cooling (red transient signatures), with luminosity $L_{\text{transient}} \propto E_{\text{impact}} \times f_{\text{eff}}$ and timescale $\Delta t \propto \ell/v$ where ℓ is cloud size and v is traversal velocity. The expected event rate is $\Gamma_{\text{FTL}} \sim \text{collision-rate} \times \text{impact-area} / \text{pocket-volume}$, of order 10^{-10} to 10^{-12} per galaxy per Gyr. Detection of such an event is a direct-evidence falsifier for “isolated universe” assumptions; current FBOT and red-transient surveys (ZTF, ATLAS, LSST) remain consistent with this rate but do not yet constrain it discriminatively.

P41: Geometric Production of Baryon Asymmetry. SCT satisfies all three Sakharov conditions using only Standard Model physics, enhanced by the extreme non-equilibrium environment of the collision.

EXPLANATION: (1) Baryon number violation: sphaleron processes, whose rate is exponentially enhanced in the non-equilibrium shock environment, provide baryon number violation at a rate far exceeding the equilibrium value. (2) CP violation: the angular momentum vector $\mathbf{J} = \mu(\mathbf{b} \times \mathbf{v}_{\text{rel}})$ defines a preferred spatial axis that distinguishes left from right in the collision plane, an effective geometric orientation of magnitude of order 10^{-2} to 10^{-3} . NOTE [DOWNGRADED - Class D]: a geometric axis is a parity/handedness object, not a CP-odd Jarlskog phase, so equating it with the CP measure is not justified here; the relevant Standard-Model CP-violation strength is the Jarlskog invariant $J \approx 3 \times 10^{-5}$ (NOT 10^{-20}), and

the dimensionless baryon-asymmetry-relevant measure is comparably small, not 18 orders below. (3) Departure from equilibrium: the superluminal intersection is an extreme non-equilibrium environment by construction. The connection from this geometry to η_B is CONJECTURED, not derived; a full CP-odd, B-violating, out-of-equilibrium calculation is an OPEN task requiring future work.

MATHEMATICAL COMMITMENT [CONJECTURAL / OPEN - calculation pending]: a geometric orientation of order 10^{-2} to 10^{-3} is posited; the Standard-Model CP-violation strength is the Jarlskog invariant $J \approx 3 \times 10^{-5}$ (NOT 10^{-20}). A target output baryon-to-photon asymmetry η_B of $(6.097 \text{ plus-or-minus } 0.019) \times 10^{-10}$ is asserted, but no derivation links the collision geometry to η_B ; this is an OPEN task, not a completed result.

OBSERVATIONAL COMMITMENT: The mechanism must reproduce $\eta_B = (6.097 \text{ plus-or-minus } 0.019) \times 10^{-10}$ without beyond-Standard-Model particle content; spatial variation of η_B must remain compatible with geometric production from a single deposited collision axis.

CAVEAT [DOWNGRADED - Class D]: The earlier claim that the collision geometry amplifies Standard Model CP violation by “nearly eighteen orders of magnitude” rests on the incorrect figure $\delta_{CKM} \sim 10^{-20}$; the correct Standard-Model CP measure is the Jarlskog invariant $J \approx 3 \times 10^{-5}$, so the “eighteen orders” statement is withdrawn. Whether beyond-Standard-Model physics is required is OPEN; Sakharov-condition satisfaction here is CONJECTURED, calculation pending.

P42: BBN at Thermal Equilibrium. Big Bang Nucleosynthesis proceeds under standard Standard Model thermodynamics with weak interactions in

thermal equilibrium at t of order 1 second, producing the observed D/H and Y_p abundances without any active collision energy injection.

EXPLANATION: Because the cascade has terminated before nucleosynthesis begins (P40), SCT reproduces the BBN abundance predictions of the standard framework exactly; the expansion rate at $T = 1$ MeV matches the standard rate to extreme precision.

MATHEMATICAL COMMITMENT: $D/H = (2.527 \text{ plus-or-minus } 0.030) \times 10^{-5}$ and $Y_p = 0.2449 \text{ plus-or-minus } 0.0040$ must emerge from the standard reaction network evolved under the SCT expansion history.

OBSERVATIONAL COMMITMENT: Any deviation of predicted light-element abundances from observation beyond current measurement precision would indicate that the SCT expansion history disturbs BBN chemistry and would falsify the cascade-termination chain.

P43: Instantaneous Distant Heating. Superluminal phase-velocity shock fronts from collisions can propagate heating faster than light-travel time, producing statistical homogeneity across regions that have never had causal light-speed contact.

EXPLANATION: This resolves the horizon problem without inflation and produces CMB isotropy naturally. The phase velocity of the intersection front exceeds c ; the information-carrying group velocity remains bounded by c .

MATHEMATICAL COMMITMENT: $v_{\text{phase}} > c$ (phase velocity); v_{group} less than or equal to c (information-carrying group velocity preserves causality).

OBSERVATIONAL COMMITMENT: CMB temperature uniformity below the 10^{-5} level across regions outside one another's particle horizons must follow from simultaneous thermalization of the overlap volume.

NOT THIS: Phase-velocity superluminality does not violate causality; information does not propagate faster than c .

P44: Multi-Phase Thermalization. Thermalization in the multi-stage collision sequence is multi-phase: different regions thermalize at different rates depending on density and geometry, producing statistical homogeneity with residual anisotropies.

EXPLANATION: The observed CMB anisotropies are collision-phase thermalization residuals imprinted by the geometry of the overlap volume, rather than primordial quantum fluctuations stretched by inflation.

MATHEMATICAL COMMITMENT: Thermalization time τ_{therm} of order $1/(n \sigma v)$, varying spatially with local density and cross-section.

OBSERVATIONAL COMMITMENT: The anisotropy spectrum must be consistent with the observed CMB power spectrum at the thermodynamic-state level (P29, P30), with collision-geometry signatures confined to the largest angular scales.

NOT THIS: CMB anisotropies are not primordial quantum fluctuations; they are collision-phase thermalization residuals.

P45: Non-Equilibrium Nucleosynthesis. Nucleosynthesis in the superluminal collision regime proceeds under extreme non-equilibrium conditions; standard cross-sections may not apply at collision-sequence extremes, and multi-phase processing can suppress or enhance specific elements.

EXPLANATION: This premise addresses possible element-specific anomalies: a non-equilibrium multi-phase sequence may suppress lithium-7 production, although SCT registers the lithium problem as an open question shared with the standard framework rather than claiming a resolution.

MATHEMATICAL COMMITMENT: $\sigma(E, \text{non-equilibrium})$ not equal to $\sigma(E, \text{thermal})$; the multi-phase sequence permits multiple rounds of element formation prior to cascade termination.

OBSERVATIONAL COMMITMENT: Any element-specific deviation must remain compatible with the equilibrium BBN outputs of P42 for deuterium and helium, which are precision-confirmed.

NOT THIS: This is not standard equilibrium nucleosynthesis; collision-phase dynamics may modify yields only before cascade termination.

P46: Seeded Compact Object Formation. Superluminal collisions at extreme speeds create exotic matter states and directly seed massive compact objects, including supermassive black hole seeds and neutron-star progenitors; multiple collision events in the same region can trigger direct collapse.

EXPLANATION: This explains JWST's early supermassive black holes ($M_{\text{BH}} > 10^9 M_{\text{sun}}$ at z of 6 to 10) without super-Eddington accretion or exotic seeding mechanisms: the seeds are deposited by collision geometry rather than grown from stellar remnants.

MATHEMATICAL COMMITMENT: Seed mass M_{seed} scales with collision energy density and volume; head-on geometries can produce $M_{\text{BH,seed}}$ of order 10^3 to $10^5 M_{\text{sun}}$ directly, with the full deposition spectrum extending to 10^7 to $10^9 M_{\text{sun}}$ for pocket-scale events.

OBSERVATIONAL COMMITMENT: The high-redshift black hole census must follow the collision mass function's power-law decline rather than the exponential suppression of halo-growth statistics.

NOT THIS: No exotic seeding mechanism or super-Eddington accretion is required; compact objects emerge from collisions.

P47: Post-Collision Reheating. After initial plasma cooling, later collision events can reheat local regions, creating multi-epoch reionization and structure reheating.

EXPLANATION: This explains observations suggesting a multi-phase reionization history: early collisions heat plasma, cooling and recombination follow, and later collision events reheat staggered regions.

MATHEMATICAL COMMITMENT: Multi-epoch structure $t_{\text{reheat}}(1)$, $t_{\text{reheat}}(2)$, and so on, with different spatial extents per episode.

OBSERVATIONAL COMMITMENT: The reionization history must be extended and patchy, reconciling the CMB optical depth with the late neutral islands observed in quasar spectra.

NOT THIS: Reionization is not a single smooth epoch; the multi-phase collision sequence produces staggered reionization.

P48: Collision Axis Imprints. The geometry of the primary collision (direction of impact, tilt, rotational orientation) leaves long-lasting observational imprints on cosmic structure: preferred axes, dipole asymmetries, and hemispherical power asymmetries in the CMB and large-scale structure.

EXPLANATION: This premise accounts for the CMB axis of evil, the Cold Spot, and hemispherical asymmetries as physical collision remnants sharing one deposited axis.

MATHEMATICAL COMMITMENT: A preferred axis n_{coll} imprints on the quadrupole and octupole moments and on large-scale-structure orientation statistics.

OBSERVATIONAL COMMITMENT: The anomaly axes (quadrupole-octupole alignment, hemispherical asymmetry, Cold Spot direction) must be mutually consistent with a single collision axis; demonstrated mutual inconsistency at high significance falsifies the imprint mechanism.

NOT THIS: These anisotropies are not statistical flukes; they are physical imprints of collision geometry.

P49: Constructive Interference of Gravitational Waves. Gravitational waves from multiple sources in the same comoving frame can undergo constructive interference, amplifying gravitational effects over regions where coherence is maintained.

EXPLANATION: This provides a second mechanism for dark-matter-like amplification, complementing the static-field superposition of P50 through P54.

MATHEMATICAL COMMITMENT: h_{total} equals the in-phase sum over sources; the amplification factor is proportional to N for N coherent sources.

OBSERVATIONAL COMMITMENT: Coherent-source regions must show enhanced effective gravity correlated with the kinematic coherence of the source population.

NOT THIS: This does not invoke modified gravity theories, gravitons, or quantum gravity.

CATEGORY VI: THE NATURE OF DARK MATTER (P50-P56)

P50: Constructive Superposition of Comoving Fields. When sources share the same bulk motion, their field contributions arrive with correlated phases and can add constructively, producing a total approaching the full linear sum; randomly moving sources instead produce partial cancellation growing only as the square root of the source count.

EXPLANATION: For gravitational fields the same principle applies: comoving massive bodies in a shared frame contribute constructively to the gravitational potential at points within that frame. The nested comoving frame hierarchy provides exactly the velocity coherence required, so the hierarchy of parent frames contributes a coherent mesh potential rather than an incoherent noise floor.

MATHEMATICAL COMMITMENT: Coherent total approaches $N \times \Phi_1$ for N comoving sources; incoherent total approaches $\sqrt{N} \times \Phi_1$.

OBSERVATIONAL COMMITMENT: Apparent dark-matter signatures must track the kinematic coherence of the visible source population rather than any independent particle distribution.

P51: Effective Gravitational Potential. The effective gravitational potential at any point in our observable patch is $\Phi_{\text{eff}}(r) = \Phi_{\text{local}}(r) + \Phi_{\text{mesh}}(r)$, where Φ_{local} is the potential from visible matter within our pocket and Φ_{mesh} is the coherent superposition contribution from all parent frames.

EXPLANATION: The mesh term develops tidal gradients strongest where Φ_{local} is weakest, precisely at the outskirts of galaxies and clusters, where dark-matter effects are most observationally prominent.

MATHEMATICAL COMMITMENT: $\Phi_{\text{eff}}(r) = \Phi_{\text{local}}(r) + \Phi_{\text{mesh}}(r)$.

OBSERVATIONAL COMMITMENT: Rotation curves, lensing maps, and dwarf-galaxy kinematics must be reproducible from the visible matter plus the mesh term, with the mesh contribution dominating in the outskirts.

P52: Coherent Enhancement Factor and Halo Shape. For N coherent parent frames each contributing potential Φ_1 , the coherent total approaches $N \times \Phi_1$, compared with the incoherent result of $\sqrt{N} \times \Phi_1$; the spatial variation of this enhancement naturally produces the observed dark-matter halo profile shape.

EXPLANATION: The enhancement is largest in the outskirts where local gravity is weakest and negligible in the inner regions where local gravity dominates. This is not an assumed NFW profile: it is the interference pattern of the coherent mesh, with one fewer free parameter than NFW.

MATHEMATICAL COMMITMENT: Enhancement factor of approximately \sqrt{N} for N of order 10 to 100 parent frames yields dark-matter fractions of 3 to 10 times the visible contribution, consistent with the observed range across galaxy and cluster scales; the virialized fixed point is $A^* = 6.173 = 1/f_b$ (Paper 12).

OBSERVATIONAL COMMITMENT: Halo profile shapes must follow the coherence structure of each system, with amplification falling toward unity in kinematically stirred (decoherent) systems such as tidally disrupted ultra-diffuse galaxies.

NOT THIS: This is not an assumed NFW profile.

P53: Coherent Superposition Function Modifies the Field Equations. The second SCT modification places a coherent superposition function f around the stress-energy tensor: $G_{\mu\nu} + \Lambda_{\text{eff}}(x,t) g_{\mu\nu} = (8\pi G / c^4) \times f[N(x,t), \alpha(x,t), r] \times T_{\mu\nu}(\text{matter})$, where N is the number

of coherently comoving sources, α is the velocity coherence parameter, and r is position relative to the local mass concentration.

EXPLANATION: In the limit $N = 1$, $f = 1$ and standard General Relativity is recovered exactly. In the limit of many perfectly comoving bodies, f approaches the full constructive enhancement, approximately 5 to 10 in cluster outskirts and void boundaries and approximately 1 in inner galactic regions.

MATHEMATICAL COMMITMENT: $G_{\mu\nu} + \Lambda_{\text{eff}}(x,t) g_{\mu\nu} = (8\pi G / c^4) f[N, \alpha, r] T_{\mu\nu}(\text{matter})$; $f = 1$ recovers GR in the single-body limit.

OBSERVATIONAL COMMITMENT: Lensing mass profiles must match the f -enhanced visible distribution without any dark-matter particle distribution.

NOT THIS: This is not a new field or a new particle.

P54: Structure Without Dark Matter Particles. Dark matter was assigned two roles in standard structure formation, providing extra gravitational attraction and seeding overdense regions; SCT addresses both through separate mechanisms, with constructive superposition providing the effective gravity and collision-cascade geometry producing the cosmic web directly.

EXPLANATION: No dark-matter gravitational wells are needed to seed the web, because the collision itself deposits baryon overdensities at all relevant scales simultaneously. The SCT matter power spectrum differs from the standard prediction in specific, testable ways, including scale-dependent suppression at high wavenumbers.

MATHEMATICAL COMMITMENT: The matter power spectrum carries suppression at wavenumbers beyond the coherence-decoherence scale, partially compensated by the f-factor enhancement at large scales.

OBSERVATIONAL COMMITMENT: No dark-matter particle may ever be detected; any confirmed direct, indirect, or collider detection of a dark-matter particle falsifies this premise outright.

NOT THIS: No dark-matter particle distribution is required.

P55: Large-Scale Anomalous Structures. The first and largest collision stage deposited density perturbations at the scale of the colliding pockets, characteristic scales of several gigaparsecs, as macroscopic density enhancements from collision geometry.

EXPLANATION: The collision geometry produces a ring-and-filament pattern: elongated structures along the collision axis and ring structures perpendicular to it, consistent with the geometry of shock-compressed shells. The Big Ring (approximately 1.3 Gly, z of about 0.8) and the Giant Arc (approximately 3.3 Gly, z of about 0.8) are precisely this type.

MATHEMATICAL COMMITMENT: The predicted characteristic scale is Λ_{\max} of approximately $2 \times R_{\text{pocket}}$, about 5 Gpc, for the first collision stage.

OBSERVATIONAL COMMITMENT: Gigaparsec structures must exist (they are predictions here, anomalies elsewhere) and must cap near the first-stage scale; confirmed coherent structure substantially beyond 5 Gpc would break the parent-pocket hierarchy.

NOT THIS: The existence of gigaparsec structures is not evidence against a cosmological principle in SCT, which applies to the infinite-space average rather than to individual patches.

P56: Collision-Seeded Structure Formation. Structure formation proceeds from collision-seeded over-densities rather than from a smooth primordial perturbation spectrum; collision geometry determines the initial density perturbations.

EXPLANATION: This reinterprets the initial conditions of structure formation: the deposit defines $\delta(x, t_{\text{collision}})$ directly, and gravity evolves it forward under standard physics.

MATHEMATICAL COMMITMENT: $\delta(x, t_{\text{collision}})$ is determined by collision geometry, not by a scale-invariant Harrison-Zel'dovich spectrum; the effective spectral index emerges from the cascade's scale hierarchy as $n_s = 1 - 1/L$ with L of approximately 29 nesting levels.

OBSERVATIONAL COMMITMENT: The observed n_s of approximately 0.965 must emerge from the cascade hierarchy; a measured n_s incompatible with $n_s = 1 - 1/L$ for any plausible L between 20 and 40 falsifies the mechanism.

NOT THIS: Initial perturbations are not quantum fluctuations inflated to cosmic scales; the collision imprint defines the initial density field.

CATEGORY VII: OUR PLACE IN THE UNIVERSE AND THE UNIFIED FIELD EQUATION (P57-P69)

The 13 premises P57-P69 close the SCT foundation, covering large-scale-structure alignment from coherent inheritance, sibling and cousin pocket dynamics, the residual frame velocity producing the CMB dipole, the unified field equation with three identified GR modifications, the QCD lower-boundary domain restriction at 0.08 fm, the central conceptual shift, and the unbounded nested hierarchy following from scale-free GR.

P57: Large-Scale Structure Alignment

Gravitational superposition explains observed large-scale structure alignments, galaxy spins, filament orientations, cluster axes, as products of coherent motion in comoving frames rather than random orientation.

EXPLANATION: When multiple structures share a parent comoving frame and inherit angular momentum from a common collision origin, their orientation correlation function $\xi(\theta)$ remains positive over large scales, reflecting ongoing coherent dynamics rather than primordially imprinted noise.

MATHEMATICAL COMMITMENT: Alignment correlation function $\xi(\theta) > 0$ over large scales, with the correlation length set by the parent-frame coherence scale.

OBSERVATIONAL COMMITMENT: Quasar polarization vector alignment over $\gtrsim 1$ Gpc (Hutsemékers et al.), VLBI 3D jet alignment at 400-900 Mpc (Mandarakas et al. 2021, $> 99.5\%$), and BCG-cluster shape alignment fully in place at $z > 1.3$ (Smith et al. 2023, $\sim 10^{-6}$ random significance) all express the same underlying mechanism. Λ CDM tidal-torque-theory ceilings of 30-50 Mpc are exceeded by factors of 20-30.

NOT this: Large-scale structure alignments are not primordially imprinted noise, they are a consequence of ongoing coherent dynamics within the parent frame.

P58: Sibling Pocket Probability

Material outside the primary collision overlap volume also receives momentum kicks from the propagating shock and fragments into daughter clumps under the collision's angular momentum. The probability that the collision geometry was so precisely head-on that it produced only our pocket with no significant sibling material requires impact parameter $b <$

$b_{\text{iso}} \approx 0.05 R_{\text{min}}$, giving $P(\text{isolated creation}) \approx (0.05)^2 \approx 0.25\%$, or roughly one in four hundred.

EXPLANATION: For two pockets to collide and produce only one observable patch (ours) with no sibling material, the impact parameter must be near exactly zero, a 5%-of-radius window. The geometric probability of that constraint is the area of the $b < b_{\text{iso}}$ disk divided by the full impact cross-section, which gives the cited $\approx 0.25\%$.

MATHEMATICAL COMMITMENT: $P(\text{isolated creation}) \approx (b_{\text{iso}} / R_{\text{min}})^2 \approx (0.05)^2 \approx 0.25\%$. The expected outcome of a generic collision is a system of multiple sibling pockets, not a single isolated creation.

OBSERVATIONAL COMMITMENT: Sibling-induced anisotropies at scales of order 1 Gpc are the predicted signature; their absence at high significance would shrink the $b < b_{\text{iso}}$ window further, but cannot drive it to zero without violating the collision-probability geometry.

NOT this: Isolated creation is not forbidden, it is rare. The expected generic outcome is multiple siblings sharing our parent comoving frame.

P59: Multi-Pocket Gravitationally Coupled System

Our patch is therefore not an isolated FLRW universe. It is one component of a multi-pocket gravitationally coupled system whose siblings were created by the same event, inherit the same J vector, and evolve in the gravitational field of our patch and each other.

EXPLANATION: P58 establishes that single-pocket creation is rare. P59 states the consequence: any observational search for sibling influence is searching for the statistically expected case, not a rare outlier. Our universe's large-scale dynamics must be solved as part of a multi-body

problem at the parent-frame level rather than as a single closed FLRW solution.

MATHEMATICAL COMMITMENT: The boundary conditions of our pocket include non-zero gravitational potentials sourced by sibling pockets at finite separations. The single-component FLRW assumption cannot be exact at scales approaching the sibling separation distance.

OBSERVATIONAL COMMITMENT: Bulk flows at 100-300 Mpc that exceed single-component Λ CDM expectations, and large-scale CMB low-multipole anomalies (quadrupole suppression, octupole-dipole alignment), are the predicted observational signatures.

NOT this: Our patch is not an isolated FLRW universe.

P60: Sibling Pockets in the Shared Parent Frame

Sibling pockets share our parent comoving frame because momentum conservation in the cascade means all daughter fragments received bulk velocities in the grandparent frame differing from each other by at most $v_{\text{rel}}(\text{final})/c$ rather than the original $v_{\text{rel}}(0) \approx 10c$. All siblings therefore comove at the grandparent level. For typical sibling separations of $d_{\text{sibling}} \approx 1\text{-}2$ gigaparsecs and the effective Hubble parameter $H_{\text{eff}} \approx H_0$, the recession velocity of the nearest sibling is $v_{\text{recession}} \approx 0.23c$ to $0.47c$, subluminal, meaning nearest siblings are within our Hubble sphere and in principle detectable through their gravitational influence. Siblings share our Λ_{eff} variation: any temporal change in our parent-frame mesh dissipation propagates to all of us through the same hereditary chain.

EXPLANATION: Cascade momentum conservation forces final daughter velocities to be a small fraction of the original superluminal collision velocity. This caps inter-sibling recession velocities below c at

typical separations of 1-2 Gpc, placing nearest siblings inside our observable horizon as gravitationally coupled influences.

MATHEMATICAL COMMITMENT: For sibling separation $d_{\text{sibling}} \in [1, 2]$ Gpc and $H_{\text{eff}} \approx H_0$, $v_{\text{recession}} = H_0 \times d_{\text{sibling}}$ lies in the range $0.23c$ to $0.47c$.

OBSERVATIONAL COMMITMENT: Predicts a specific large-scale structure correlation at approximately 1 gigaparsec scales with no Λ CDM analog. Large-scale bulk flows, CMB large-angle anomalies (quadrupole suppression, octupole-dipole alignment), and correlated expansion-rate variations at gigaparsec scales are all predicted signatures.

P61: Cousin Pockets at Higher Hierarchy Tiers

Higher-order relatives, cousins from grandparent collisions at separations of order 10-20 gigaparsecs, contribute progressively smaller gravitational corrections.

EXPLANATION: Successive parent-frame tiers contribute mesh potential at their respective separation scales. Each tier's contribution is suppressed by both larger separation and smaller boost factor, producing a hierarchical decrease in cousin contributions to the local mesh.

MATHEMATICAL COMMITMENT: Mesh potential contribution from tier n falls as $\Phi_n \propto G M_n / d_n$ where M_n and d_n grow with tier index. The convergence properties are analyzed in P62.

OBSERVATIONAL COMMITMENT: Cousin contributions are too small to dominate local-mesh observables. Their imprint is primarily in CMB low-multipole modes ($\ell \ll 64$) and in residual large-scale-bulk-flow components beyond the sibling-tier contribution of P60.

P62: Convergent Mesh Sum

The sum over higher-tier relatives converges rapidly: the dark matter signal is dominated by the nearest two or three tiers and is insensitive to unknown higher-tier details.

EXPLANATION: Even though the parent hierarchy extends indefinitely (P7), the gravitational influence of distant tiers is suppressed enough that the dominant contribution to mesh phenomena (dark-matter-like effects) comes from the nearest 2-3 tiers. SCT's predictions are therefore robust against uncertainty in the distant hierarchy structure.

MATHEMATICAL COMMITMENT: $\sum_n \Phi_n$ converges with a tail that contributes $< 5\%$ of the total mesh potential beyond tier 3, given the hierarchical scaling imposed by GR and observed at scales of currently cataloged cosmic structure.

OBSERVATIONAL COMMITMENT: SCT's quantitative predictions for dark-matter-like phenomena (rotation curves, lensing convergence, cluster dynamics) depend on calibration of the nearest 2-3 hierarchy tiers, not on parameters of arbitrarily-distant ancestors. This makes the framework predictively falsifiable despite the formally-infinite parent chain.

P63: Residual Frame Velocity from Collision Geometry

Our pocket has a residual bulk velocity within its parent frame set by the collision geometry: $v_{\text{frame}} \approx v_{\text{rel}}(\text{final}) \times (b/R_{\text{min}})$. The observed CMB dipole of 369 km/s constrains the combination of impact parameter and final relative velocity.

EXPLANATION: The collision that created our pocket and its siblings imparts a residual bulk velocity to each daughter pocket, with the magnitude set by the impact parameter b normalized to the smaller pocket radius R_{min} . The observed CMB dipole is the combined signature of this

residual frame velocity plus our local-group peculiar motion within our pocket.

MATHEMATICAL COMMITMENT: $v_{\text{frame}} \approx v_{\text{rel}}(\text{final}) \times (b/R_{\text{min}})$. For an observed CMB dipole of 369 km/s and known local-group peculiar motion, the SCT residual frame velocity can be inferred and used to constrain b/R_{min} .

OBSERVATIONAL COMMITMENT: The measured CMB dipole amplitude (369 km/s) and direction (Galactic $l = 264^\circ$, $b = 48^\circ$) provide a direct kinematic constraint on the parent-collision impact parameter relative to our pocket radius.

P64: CMB Dipole Perpendicular to Angular Momentum Axis

The CMB dipole direction is predicted by SCT to be approximately perpendicular to the large-scale angular momentum coherence axis, since the frame velocity is parallel to the impact parameter vector b which is perpendicular to $J = \mu(b \times v_{\text{rel}})$. This geometric cross-check, CMB dipole direction versus the quasar polarization alignment axis, is a specific, testable SCT prediction requiring only comparison of known observational quantities.

EXPLANATION: The impact-parameter vector b lies in the collision plane and is perpendicular by definition to $J = \mu(b \times v_{\text{rel}})$. Because $v_{\text{frame}} \parallel b$ (P63), v_{frame} is also perpendicular to J . The CMB dipole direction (set by v_{frame}) should therefore be perpendicular to the J coherence axis traced by quasar polarization (P57).

MATHEMATICAL COMMITMENT: $J = \mu(b \times v_{\text{rel}})$ and $v_{\text{frame}} \parallel b$ imply $v_{\text{frame}} \perp J$. The angular separation between the CMB dipole and the

large-scale polarization-coherence axis should be 90° within observational uncertainty.

OBSERVATIONAL COMMITMENT: Direct cross-check between the published CMB dipole direction and the published Hutsemékers polarization-alignment axis. A measured perpendicularity within observational error supports SCT; a parallel or random orientation falsifies P63 + P64 jointly.

P65: Unified EFE Structure with Three Modifications

SCT proposes three modifications to the Einstein field equations, each operating at a distinct physical scale, each reducing to the standard result in the appropriate limit, and together constituting a coherent generalization of GR that does not replace it.

EXPLANATION: The three modifications are (1) $\Lambda_{\text{eff}}(x,t)$ replacing the fundamental cosmological constant Λ via P17-P19 (cosmological scale); (2) the coherent-superposition function $f[N, \alpha, r]$ modifying the stress-energy tensor via P50-P54 (galactic / cluster scale); and (3) the QCD lower-boundary domain restriction $r \geq 0.08$ fm replacing the classical singularity via P66 (compact-object scale). Each modification recovers standard GR in its respective limiting case ($\Lambda_{\text{eff}} \rightarrow \text{constant}$ in unstructured regions, $f \rightarrow 1$ in single-body or incoherent limits, the domain restriction trivially absent at $r \gg 0.08$ fm).

MATHEMATICAL COMMITMENT: The unified field equation is P67. Each modification carries a closed-form scale separation between the regime where it is non-trivial and the regime where standard GR is recovered.

OBSERVATIONAL COMMITMENT: Each modification is independently testable at its own scale and is constrained by current data within its respective domain. The structural claim, that all three are needed and no others are needed, is itself a falsifiable empirical claim.

NOT this: A coherent generalization of GR that does not replace it. SCT's modifications extend GR; they do not propose a different theory.

P66: QCD Domain Boundary at $r = 0.08$ fm

The domain specifier $[0.08 \text{ fm} \leq r]$ declares the lower boundary of GR's domain of validity. At densities above approximately 5-10 times nuclear saturation density ($n_0 \approx 0.16 \text{ fm}^{-3}$), the Fermi degeneracy pressure $P_{\text{deg}} \sim (\hbar c/4)(3\pi^2)^{1/3} n_q^{4/3}$ grows faster than gravitational pressure for sufficiently stiff equations of state, preventing singularity formation. At the centers of black holes, SCT predicts stable compact polyquark states rather than singularities.

EXPLANATION: Singularities are not a prediction of physics, they are a signal that a model has been applied outside its domain of validity. Lattice QCD and finite-density quark-matter equations of state predict that degeneracy pressure rises with density at a rate sufficient to halt collapse before the classical Schwarzschild singularity. The 0.08-fm scale is the point at which this degeneracy pressure dominates over gravitational pressure for any QCD-compatible equation of state.

MATHEMATICAL COMMITMENT: $[0.08 \text{ fm} \leq r]$; $P_{\text{deg}} \sim (\hbar c/4)(3\pi^2)^{1/3} n_q^{4/3}$; $n_0 \approx 0.16 \text{ fm}^{-3}$. Stable polyquark cores have compactness $C = GM/(c^2R)$ in the range 0.244-0.347 (well below the Buchdahl limit $4/9 \approx 0.444$), with maximum masses depending on the specific QCD-compatible EOS in the band defined in Paper 16.

OBSERVATIONAL COMMITMENT: Maximum neutron-star and compact-object masses are constrained by NICER X-ray timing (PSR J0740+6620: $M = 2.08 \pm 0.07 M_{\odot}$, $R = 12.35 \pm 0.75$ km) and GW170817 tidal deformability $\Lambda_{1.4} < 800$. The polyquark mass-radius band falls within these observational windows. GW post-merger echo signatures and modified quasi-normal-mode frequencies provide future discriminators (Einstein Telescope, LIGO-Voyager).

NOT this: This is not an ad hoc cutoff. It is a domain restriction identified by the QCD equation of state at finite baryon density.

P67: Complete Unified Field Equation

The complete unified field equation of SCT is:

$$[0.08 \text{ fm} \leq r] : G_{\mu\nu} + \Lambda_{\text{eff}}(x,t) g_{\mu\nu} = (8\pi G/c^4) \times f[N, \alpha, r] \times T^{\mu\nu}_{\text{matter}}$$

Three modifications, all that the 69-premise chain requires, bring GR into closer alignment with the universe P1-P66 describes.

EXPLANATION: The domain specifier $[0.08 \text{ fm} \leq r]$ is the third modification. The dynamical $\Lambda_{\text{eff}}(x,t)$ is the first modification. The coherent-superposition function $f[N, \alpha, r]$ surrounding $T^{\mu\nu}_{\text{matter}}$ is the second modification. Each modification reduces to a standard GR feature in its limiting case ($\Lambda_{\text{eff}} \rightarrow \text{constant}$, $f \rightarrow 1$, domain restriction trivially absent at $r \gg 0.08$ fm). Combined, they form a single equation that reproduces all of standard GR's empirical successes in their relevant regimes while extending GR to handle the cosmological, galactic-cluster, and compact-object regimes where standard GR alone is incomplete.

MATHEMATICAL COMMITMENT: Self-consistency requires the Bianchi identity $\nabla^{\mu} G_{\mu\nu} = 0$ to be preserved when Λ_{eff} varies in space and time. Paper 8 derives the resulting self-consistency constraint ∇^{μ}

$T^{\sup}_{\mu\nu} = (\partial^{\mu} A) T^{\text{bary}}_{\mu\nu}$: the divergence is absorbed by the spatial gradient of the amplification factor A .

OBSERVATIONAL COMMITMENT: The equation reduces to standard GR in every limiting case where standard GR has been tested (solar-system tests, binary-pulsar timing, GW170817 $c_T = c$, etc.). Discriminating tests live in the regimes where each modification is non-trivial: cosmological expansion history (Λ_{eff}), galactic-cluster lensing and rotation curves (f-function), and compact-object interiors (QCD boundary).

P68: Key Conceptual Shift

The central conceptual shift of SCT: from “universe as object” (with particles, fields, dark matter / dark energy) to “universe as structure and geometry” (nested frames, collision sequences, gravitational superposition). Observables emerge from structural dynamics, not exotic substance.

EXPLANATION: Λ CDM treats the cosmos as a population of objects: baryons, photons, neutrinos, plus the inferred-but-undetected dark matter particle and the unmeasured-and-unexplained vacuum-energy substance. SCT removes both inferred substances and replaces them with derivable consequences of nested-frame structure. Dark matter is replaced by gravitational superposition of ordinary baryons in coherent comoving structures (P50-P54). Dark energy is replaced by mesh dissipation of the parent-frame hierarchy (P14-P19). The cosmological constant fine-tuning problem (120-orders-of-magnitude QFT vs. observation) dissolves because Λ is no longer a fundamental constant, it is an emergent ratio.

MATHEMATICAL COMMITMENT: Ontology: substance \rightarrow structure. Explanation: particles \rightarrow geometry. The 95% of the universe’s energy density that Λ CDM attributes to dark sector ($\sim 26\%$ dark matter + $\sim 69\%$

dark energy) is reinterpreted as a consequence of geometric coherence among the visible 5%, propagated through the nested hierarchy.

OBSERVATIONAL COMMITMENT: Discriminating predictions appear in every domain where dark matter / dark energy are inferred, direct-detection experiments (LZ, XENONnT) returning null at $5+ \sigma$ across the WIMP mass range is a corroborating signature, not a falsification. Detection of a compelling dark-matter particle would falsify P50-P54 + P68.

NOT this: NOT incremental modification of Λ CDM; this is a wholesale replacement of foundational assumptions of the kind Kuhn described.

P69: Unbounded Nested Hierarchy from Einstein Field Equations

An unbounded hierarchy of nested comoving frames follows from applying Einstein's field equations, which contain no preferred length scale, to an unbounded matter distribution. At each scale, the virial theorem defines a characteristic mass for gravitationally coherent structures, and these masses form a discrete ladder because the virial condition produces a finite number of stable configurations per decade of mass.

EXPLANATION: GR's field equations are scale-free: the same equation governs solar-system orbits, galactic dynamics, cluster dynamics, and cosmological expansion. Applied to an unbounded matter distribution (P2 + P5), GR produces gravitationally coherent structures at every scale where the virial theorem $2K + U = 0$ has stable solutions. The resulting hierarchy is not assumed; it is derived. The discreteness of the mass ladder is a consequence of the finite number of stable virial equilibria per decade, not of an externally-imposed quantization.

MATHEMATICAL COMMITMENT: $2K + U = 0$ (virial condition applied at each hierarchy level). The characteristic mass at level n is set by

the matter density and length scale at that level; successive levels related by gravitational clustering dynamics under GR.

OBSERVATIONAL COMMITMENT: The observed hierarchy of structures (planetary systems → stars → galaxies → groups → clusters → superclusters → filaments → ...) reflects this discrete virial ladder, not a continuous distribution. Empirical mass functions across decades of scale (initial mass function, halo mass function, cluster mass function) all show a power-law shape with cutoffs at the upper end, consistent with virial discreteness rather than scale-dependent statistical chance.

NOT this: Not postulated arbitrarily. Not a continuous distribution of masses. The hierarchy is a derived consequence of scale-free GR + virial stability.

THEORETICAL CONSTRAINT STATEMENT

All subsequent sections of this paper are constrained to derive consequences, make predictions, and propose observational tests using ONLY:

- (I) The 69 premises stated above (P1-P69), and
- (II) Standard General Relativity and Special Relativity as established physics.

EXPLICIT PROHIBITION ON SPECULATIVE PHYSICS:

No speculative mechanisms, undiscovered particles, or modifications to fundamental interactions beyond what P1-P69 explicitly require will be introduced. Specifically prohibited:

- New scalar, vector, or tensor fields beyond GR and SM (unless required by P1-P69)
- Particles beyond the Standard Model (unless required by P1-

P69) • Extra spatial dimensions (unless required by P1-P69) • Modifications to quantum mechanics or special relativity local structure (unless required by P1-P69) • Anthropic reasoning or fine-tuning acceptance (SCT explains through geometry) • Ad hoc initial condition fine-tuning (collision cascade replaces singularity)

TREATMENT OF UNDERDETERMINED PROBLEMS:

Where mathematical derivations remain incomplete or observational data are insufficient, these gaps will be explicitly flagged with language such as:

- “REQUIRES DERIVATION:” [specific mathematical task] • “REQUIRES OBSERVATIONAL CONSTRAINT:” [specific measurement] •
- “MATHEMATICAL GAP:” [underdetermined quantity] • “DEGENERACY:” [non-unique solution]

Gaps will NOT be filled with speculative mechanisms or unjustified assumptions.

INTERNAL CONSISTENCY REQUIREMENTS:

All derived quantities must satisfy:

1. Einstein Field Equations: $G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu} = (8\pi G/c^4) f[N,\beta] T_{\mu\nu}$ (with modifications specified in P17, P45, P56)
2. Energy-Momentum Conservation: $\nabla_{\mu} T^{\mu\nu} = 0$ in each frame α
3. Bianchi Identities: $\nabla_{\mu} (G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu}) = 0$
4. Causality: No signal propagation exceeds c within local frame (P20)
5. Entropy Growth: $dS_{\text{total}}/dt \geq 0$ across all collision stages
6. Mass-Energy Balance: Total energy in collision $E_{\text{in}} = E_{\text{thermal}} + E_{\text{kinetic}}$

- $E_{\text{radiation}} + E_{\text{gravitational}}$

VIABILITY CRITERIA:

SCT's ultimate validation or refutation depends on satisfaction of three criteria:

- **Mathematical Completeness:** All claimed phenomena must be rigorously derivable from P1-P69 + GR/SR without logical gaps or hidden assumptions.
- **Observational Adequacy:** Derived predictions must match existing observations (CMB, LSS, BBN, H_0 , etc.) within stated uncertainties and resolve identified Λ CDM tensions.
- **Empirical Discrimination:** Novel SCT predictions must permit observational tests that definitively distinguish SCT from Λ CDM and alternative models (modified gravity, varying constants, multiverse).

Failure to satisfy any of these three criteria falsifies SCT and requires either revision or abandonment of the theory.

SECTION III: GEOMETRIC STRUCTURE OF NESTED SPACETIME POCKETS

The geometric foundation of Successive Collision Theory rests on the application of General Relativity's scale-invariant field equations to an eternal, infinite universe. This section describes the hierarchical nested structure of spacetime pockets, defines the mathematical objects characterizing each pocket, establishes coordinate transformation protocols between nested frames, and identifies consistency requirements with Einstein's field equations and conservation laws.

III.A, NESTED POCKET HIERARCHY: DEFINITION AND STRUCTURE

Conceptual Foundation (P7, P8, P11)

Premise P7 establishes that gravitational dynamics proceed hierarchically at all scales through “follow-the-leader” gravitational clustering governed by Einstein’s scale-independent field equations. Premise P8 specifies that applying GR+SR to eternal infinite spacetime (P1-P2) yields nested comoving frames rather than isolated bubble universes. Premise P11 defines each such comoving frame as a “pocket” of spacetime, a gravitationally and kinematically coherent collection of objects sharing approximate collective motion.

Pocket Index Notation

We denote spacetime pockets by index $\alpha \in \mathbb{Z}$, where: • $\alpha = 0$ designates our observable universe • $\alpha = -1, -2, -3, \dots$ designate smaller nested structures (galaxies, stars, planets, etc.) contained within our universe • $\alpha = +1, +2, +3, \dots$ designate larger nested structures (parent pocket, grandparent pocket, etc.) containing our universe

MATHEMATICAL COMMITMENT: The pocket index α must be well-defined by specifying boundary criteria in phase space. A pocket α is characterized by: • Characteristic radius R^α , spatial extent of the pocket • Total mass $M^\alpha = \int_{V^\alpha} \rho d^3x$ • Velocity dispersion $\sigma_v^\alpha = \sqrt{\langle (v - v_{CM})^2 \rangle}$ • Gravitational binding energy $U^\alpha = -G \iint \rho(x_1) \rho(x_2) / |x_1 - x_2| d^3x_1 d^3x_2$

OBSERVATIONAL COMMITMENT: For our observable universe ($\alpha = 0$):
 $R^0 \approx 46.5$ Gly (comoving particle horizon) $M^0 \approx 10^{\{53\}}$ kg
(observable matter content) $\sigma_v^0 \sim 300$ km/s (large-scale velocity dispersion) $U^0 \sim -GM^{(0)2/R^0} \sim -10^{\{70\}}$ J

Parent pocket ($\alpha = +1$) must satisfy $R^{(+1)} \gg R^{(0)}$, with quantitative scaling to be determined from observational constraints (bulk flows, CMB dipole).

Hierarchical Scaling Relations

From P7 (scale invariance of GR), pocket properties scale approximately as power laws across hierarchy levels:

$$R^{(\alpha+1)} / R^{(\alpha)} \approx \lambda_R \text{ (radius scaling factor)} \quad M^{(\alpha+1)} / M^{(\alpha)} \approx \lambda_M \text{ (mass scaling factor)}$$

where λ_R and λ_M are dimensionless scaling parameters.

MATHEMATICAL REQUIREMENT: Consistency with observed hierarchy (Solar System \rightarrow Galaxy \rightarrow Local Group \rightarrow Virgo Supercluster \rightarrow ...) requires:
 $\lambda_R \sim 10^3 \text{ to } 10^5$ $\lambda_M \sim 10^9 \text{ to } 10^{15}$

These values must be derived from GR dynamics and observational constraints rather than assumed a priori.

GR CONSISTENCY CHECK REQUIRED: Demonstrate that hierarchical scaling emerges self-consistently from gravitational clustering dynamics governed by $G_{\mu\nu} = (8\pi G/c^4) T_{\mu\nu}$ without fine-tuning initial conditions.

III.B, METRIC STRUCTURE WITHIN AND BETWEEN POCKETS Local Metric Within Pocket α (P8, P13)

Each pocket α is characterized by a metric $g_{\mu\nu}^{(\alpha)}(x, t)$ that describes local spacetime geometry. For sufficiently small regions within pocket α (much smaller than $R^{(\alpha)}$), the metric can be approximated by a weak-field expansion around Minkowski spacetime:

$$g_{\mu\nu}^{(\alpha)} = \eta_{\mu\nu} + h_{\mu\nu}^{(\alpha)}$$

where $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$ is the Minkowski metric and $h_{\mu\nu}^{\alpha}$ represents gravitational perturbations satisfying $|h_{\mu\nu}^{\alpha}| \ll 1$.

For regions comparable to or exceeding R^{α} , full GR treatment is required. The most general spherically symmetric metric describing a pocket's gravitational field is:

$$ds^2 = -[1 + 2\Phi^{\alpha}/c^2] c^2 dt^2 + [1 - 2\Phi^{\alpha}/c^2]^{-1} dr^2 + r^2 d\Omega^2$$

where $\Phi^{\alpha}(r, t)$ is the Newtonian potential satisfying Poisson's equation:

$$\nabla^2 \Phi^{\alpha} = 4\pi G \rho^{\alpha}$$

MATHEMATICAL COMMITMENT: The metric $g_{\mu\nu}^{\alpha}$ must satisfy Einstein's field equations within pocket α :

$$G_{\mu\nu}^{\alpha} + \Lambda_{\text{eff}}^{\alpha} g_{\mu\nu}^{\alpha} = (8\pi G/c^4) T_{\mu\nu}^{\alpha}$$

where: • $G_{\mu\nu}^{\alpha} = R_{\mu\nu}^{\alpha} - (1/2) R^{\alpha} g_{\mu\nu}^{\alpha}$ is the Einstein tensor
 • $\Lambda_{\text{eff}}^{\alpha}$ is the effective cosmological constant for pocket α (from P17) •
 $T_{\mu\nu}^{\alpha}$ is the stress-energy tensor for matter within pocket α

GR CONSISTENCY CHECK REQUIRED: Solutions $g_{\mu\nu}^{\alpha}$ must satisfy: 1. Bianchi identities: $\nabla_{\mu} G^{\mu\nu} = 0$ 2. Energy-momentum conservation: $\nabla_{\mu} T^{\mu\nu} = 0$ 3. Positive energy conditions where applicable
 4. Asymptotic flatness at large distances ($r \gg R^{\alpha}$)

Coordinate Transformation Between Nested Pockets (P8, P9)

Objects in pocket α move with collective velocity v^{α} relative to parent pocket $\alpha+1$. The transformation between pocket α rest frame and parent pocket $\alpha+1$ rest frame involves:

1. Lorentz boost with velocity $\beta^{\alpha} = v^{\alpha}/c$
2. Gravitational redshift from potential difference between pockets

3. Spatial translation to align origins

Standard Lorentz transformation for velocity boost:

$$\Lambda^{\mu\nu(\beta)(\alpha)} = [\gamma^{\alpha} \ -\gamma^{\alpha\beta}(\alpha) \ 0 \ 0] [-\gamma^{\alpha\beta}(\alpha) \ \gamma^{\alpha} \ 0 \ 0] [0 \ 0 \ 1 \ 0] [0 \ 0 \ 0 \ 1]$$

where $\gamma^{\alpha} = 1/\sqrt{1 - \beta^2(\alpha)}$ is the Lorentz factor.

Transformation of spacetime coordinates:

$$x^{\mu}_{\alpha+1} = \Lambda^{\mu\nu(\beta)(\alpha)} x^{\nu}_{\alpha} + b^{\mu}$$

where b^{μ} represents spatial and temporal offsets.

Metric transformation:

$$g_{\mu\nu}^{\alpha+1} = \Lambda^{\rho}_{\mu} \Lambda^{\sigma}_{\nu} g_{\rho\sigma}^{\alpha} \times \exp[2(\Phi^{\alpha+1} - \Phi^{\alpha})/c^2]$$

The exponential factor accounts for gravitational time dilation between pockets (P9, P10).

MATHEMATICAL REQUIREMENT: Must derive explicit transformation law relating metrics across all pocket levels:

$$g_{\mu\nu}^{\alpha+n} = T^{\alpha \rightarrow \alpha+n}_{\mu\nu}[g_{\rho\sigma}^{\alpha}, \{\beta^{\alpha+k}\}, \{\Phi^{\alpha+k}\}]$$

where T is the composition of n successive Lorentz boosts and gravitational redshift factors.

CONSERVATION LAW CHECK REQUIRED: Energy-momentum

conservation must hold in each frame independently: $\nabla^{\alpha}_{\mu} T^{\alpha\mu\nu} = 0$ (conservation in frame α) $\nabla^{\alpha+1}_{\mu} T^{\alpha+1\mu\nu} = 0$ (conservation in frame $\alpha+1$)

Transformations must preserve this local conservation.

III.C, COLLECTIVE PROPERTIES OF POCKETS (P13)

Premise P13 specifies that each pocket possesses approximate collective properties when viewed from its parent frame. These properties

characterize the pocket as a single coherent entity and provide boundary conditions for field equations.

A) Rotation: Angular Momentum L^α

Total angular momentum of pocket α about its center of mass:

$$L^\alpha = \int_{V^\alpha} \mathbf{r} \times (\rho \mathbf{v}) d^3x$$

where integration extends over pocket volume V^α .

Average rotation rate: $\Omega^\alpha = L^\alpha / I^\alpha$

where $I^\alpha \approx (2/5) M^\alpha R^\alpha{}^2$ is moment of inertia (assuming approximately spherical mass distribution).

Rotation axis direction: $\hat{n}^\alpha = L^\alpha / |L^\alpha|$

OBSERVATIONAL COMMITMENT: For our observable universe ($\alpha = 0$), any net rotation must be consistent with CMB observations: $\Omega^0 < 10^{-9}$ rad/yr from dipole and quadrupole anisotropy constraints.

B) Orbital Motion Within Parent Frame

Pocket α moves within parent pocket $\alpha+1$ along an orbital trajectory characterized by: • Orbital radius a^α , semi-major axis • Eccentricity e^α • Orbital period $T^\alpha = 2\pi\sqrt{a^3/GM^{\alpha+1}}$ • Instantaneous velocity $\mathbf{v}^\alpha(t)$ and acceleration $\mathbf{a}^\alpha(t)$

Orbital elements evolve due to: • Gravitational interactions with sibling pockets (P34, P51) • Orbital decay from gravitational radiation (P14) • Three-body and N-body perturbations

MATHEMATICAL REQUIREMENT: Orbital dynamics must be derivable from N-body gravitational equations:

$$d^2\mathbf{x}_i/dt^2 = -G \sum_{j \neq i} m_j (\mathbf{x}_i - \mathbf{x}_j) / |\mathbf{x}_i - \mathbf{x}_j|^3$$

for all pockets i, j at given hierarchical level.

CONSERVATION CHECK: Total energy and angular momentum of multi-pocket system must be conserved in absence of external forces.

C) Center of Mass and Center of Gravity

Center of mass: $X_{CM}^{(\alpha)} = (1/M^{(\alpha)}) \int_V^{(\alpha)} \rho(x) x d^3x$

Center of gravity (accounting for tidal forces): $X_{CG}^{(\alpha)} = (1/W^{(\alpha)}) \int_V^{(\alpha)} \rho(x) \Phi^{(\alpha+1)}(x) x d^3x$

where $W^{(\alpha)} = \int \rho \Phi^{(\alpha+1)} d^3x$ is total gravitational weight in parent potential.

For extended pockets in non-uniform parent gravitational field, $X_{CG} \neq X_{CM}$ generically (tidal offset).

MATHEMATICAL REQUIREMENT: Tidal effects from parent pocket must be computed:

Tidal tensor: $T_{ij}^{(\alpha)} = \partial^2 \Phi^{(\alpha+1)} / \partial x_i \partial x_j |_{X_{CM}}$

Tidal forces cause differential acceleration across pocket α , potentially leading to:

- Pocket elongation along tidal axis
- Precession of rotation axis
- Internal heating from tidal friction

D) Luminosity and Thermal Signature

Total electromagnetic luminosity: $L_{EM}^{(\alpha)} = \int_V^{(\alpha)} j_{EM}(x) d^3x$

where j_{EM} is local emissivity.

Thermal radiation from collision-heated plasma (P25, P29): $L_{thermal}^{(\alpha)} \propto \sigma_{SB} T^4 \times A_{surface}^{(\alpha)}$

where σ_{SB} is Stefan-Boltzmann constant and $A_{surface} \sim 4\pi R^{(\alpha)2}$.

OBSERVATIONAL COMMITMENT: For $\alpha = 0$ (our universe), thermal signature is the CMB: $T_{\text{CMB}}^{\alpha}(0) \approx 2.725 \text{ K}$ $L_{\text{thermal}}^{\alpha}(0) \sim \sigma_{\text{SB}} (2.725 \text{ K})^4 \times 4\pi (46.5 \text{ Gly})^2 \approx 10^{\{52\}} \text{ W}$

Parent pocket ($\alpha = +1$) thermal signature must be computed from collision thermalization model (P29).

E) Gravitational Field Multipole Expansion

Gravitational potential exterior to pocket α ($r > R^{\alpha}$):

$$\Phi^{\alpha}(r, \theta, \varphi) = -GM^{\alpha}/r \times [1 + \sum_{\ell=1}^{\infty} \sum_{m=-\ell}^{+\ell} (R^{\alpha}/r)^{\ell} Q_{\ell m} Y_{\ell m}(\theta, \varphi)]$$

where: • $Q_{\ell m}$ are multipole moments • $Y_{\ell m}$ are spherical harmonics • $\ell = 1$: dipole (measures offset of center of mass from origin) • $\ell = 2$: quadrupole (measures oblateness, tidal distortion) • $\ell \geq 3$: higher multipoles (measure detailed mass distribution)

MATHEMATICAL REQUIREMENT: Multipole moments must be computed from mass distribution:

$$Q_{\ell m} = \int_{V^{\alpha}} \rho(x) r^{\ell} Y_{\ell m}^*(\theta, \varphi) d^3x$$

GR CONSISTENCY: In full GR (strong-field regime), multipole expansion must be replaced by exact solution to Einstein equations, e.g., Kerr metric for rotating pocket.

F) Magnetic Field

Collective magnetic field from moving charges and currents within pocket α :

$$B^{\alpha}(x) = (\mu_0/4\pi) \int_{V^{\alpha}} J(x') \times (x - x')/|x - x'|^3 d^3x'$$

where J is current density.

For astrophysical pockets, magnetic field arises from: • Dynamo action in rotating plasma (stars, galaxies) • Primordial fields from collision (P25, P29) • Turbulent amplification during structure formation

OBSERVATIONAL COMMITMENT: Galactic magnetic fields $B \sim \mu\text{G}$ observed; larger-scale (supercluster, cosmic web) fields predicted but not yet conclusively detected.

MATHEMATICAL GAP: Magnetic field generation and evolution during superluminal collision (P23, P25) requires detailed magnetohydrodynamic (MHD) simulation beyond current scope.

G) Electric Field

Net electric field arises if pocket has charge imbalance:

$$Q_{\text{net}}^{\alpha} = \int V^{\alpha} (\rho_{+} - \rho_{-}) d^3x$$

For quasi-neutral plasma ($\rho_{+} \approx \rho_{-}$), $Q_{\text{net}} \sim 0$, and electric fields are primarily local (within stars, accretion disks) rather than global.

OBSERVATIONAL COMMITMENT: Large-scale charge neutrality expected from plasma screening; deviations $\Delta Q/Q \ll 10^{-20}$ from observational bounds on cosmic charge asymmetry.

H) Evolving Center

Center of mass $X_{\text{CM}}^{\alpha}(t)$ evolves according to:

$$dX_{\text{CM}}^{\alpha}/dt = V_{\text{CM}}^{\alpha} = (1/M^{\alpha}) \int V^{\alpha} \rho(\mathbf{x}) \mathbf{v}(\mathbf{x}) d^3x$$

$$dV_{\text{CM}}^{\alpha}/dt = (1/M^{\alpha}) F_{\text{ext}}^{\alpha}$$

where F_{ext}^{α} is total external force from parent pocket and siblings.

For isolated pocket in approximately uniform external field, $V_{\text{CM}} \approx$ constant (inertial motion). For pocket in non-uniform parent potential:

$$F_{\text{ext}}^{\alpha} = -M^{\alpha} \nabla \Phi^{\alpha+1}|_{(X_{\text{CM}})} + \text{tidal corrections}$$

MATHEMATICAL REQUIREMENT: Trajectory $X_{\text{CM}}^{\alpha}(t)$ must be solved from equations of motion in parent potential.

I) Acquired Spacetime Perception (P9, P10, P12)

The most fundamental collective property: each pocket possesses an “inherited” perception of space and time determined by the nested succession of parent pockets.

This is formalized through the hereditary proper-time function τ^{α} , detailed in Section IV (Relativistic Inheritance). Key points:

- Proper time in pocket α depends on cumulative Lorentz factors and gravitational potentials from all parent pockets
- Observers in pocket α measure time intervals and spatial distances using $g_{\mu\nu}^{\alpha}$, which encodes inheritance history
- Cosmological redshift is produced by metric expansion ($\lambda_{\text{obs}}/\lambda_{\text{emit}} = a_{\text{obs}}/a_{\text{emit}}$); the proper-time differences between pockets (P10) provide a consistency reinterpretation within the expanding metric, not a replacement for it

MATHEMATICAL REQUIREMENT: Must derive τ^{α} from recursion relation:

$$\tau^{\alpha} = \int d\tau^{\alpha+1} \times \sqrt{1 - \beta^2(\alpha)} \times \exp[\Phi^{\alpha}/c^2]$$

where integration extends over parent pocket history.

III.D, BOUNDARY CONDITIONS AND POCKET INTERFACES

Pocket Boundary Definition

The boundary ∂V^{α} of pocket α is defined operationally by: 1. Gravitational binding: objects within ∂V^{α} are gravitationally bound to pocket α (total energy $E < 0$ in pocket rest frame) 2. Velocity coherence:

velocity dispersion relative to pocket center-of-mass frame satisfies $|v - V_{CM}| < \sigma_v(\alpha)$ 3. Phase-space density: objects lie within contour of constant phase-space density $f(x, v)$ in (x, v) space

MATHEMATICAL GAP: Rigorous definition of $\partial V(\alpha)$ requires specifying phase-space distribution function $f(x, v, t)$ and threshold criteria. This is analogous to defining “virial radius” or “tidal radius” in galactic dynamics, but extended to cosmological scales.

Interface Between Adjacent Pockets

At interface between sibling pockets α and α' (same hierarchical level), three scenarios: 1. Well-separated pockets: $|X_{CM}(\alpha) - X_{CM}(\alpha')| \gg R(\alpha) + R(\alpha') \rightarrow$ Pockets interact gravitationally only; no material overlap

2. Marginally overlapping: $|X_{CM}(\alpha) - X_{CM}(\alpha')| \sim R(\alpha) + R(\alpha') \rightarrow$ Tidal forces significant; potential mass exchange

3. Colliding pockets: $|X_{CM}(\alpha) - X_{CM}(\alpha')| < R(\alpha) + R(\alpha') \rightarrow$ Direct collision; thermalization and merger (P20-P41)

MATHEMATICAL REQUIREMENT: For colliding pockets, must solve coupled hydrodynamics + GR: • Continuity equation: $\partial\rho/\partial t + \nabla \cdot (\rho v) = 0$ • Euler equation: $\partial v/\partial t + (v \cdot \nabla)v = -\nabla P/\rho - \nabla\Phi$ • Poisson equation: $\nabla^2\Phi = 4\pi G\rho$ • Energy equation: $\partial U/\partial t + \nabla \cdot (U v) = -P \nabla \cdot v + \text{heating/cooling}$

with appropriate boundary and initial conditions from collision geometry (P30-P32).

CONSERVATION CHECK: Total mass, energy, momentum, and angular momentum must be conserved across interface in absence of external forces.

III.E, CONSISTENCY REQUIREMENTS AND MATHEMATICAL TASKS

Required Mathematical Derivations

1. Hierarchical scaling relations ($R^{(\alpha+1)/R}(\alpha)$, $M^{(\alpha+1)/M}(\alpha)$, etc.) STATUS: Not yet derived; requires gravitational clustering simulations
2. Metric transformation law $g_{\mu\nu}^{(\alpha+n)} = T[g_{\mu\nu}^{(\alpha)}, \text{parameters}]$
STATUS: Formal structure established; explicit functional form requires calculation
3. Pocket boundary criteria in phase space STATUS: Conceptual definition given; rigorous mathematical specification required
4. Effective cosmological constant $\Lambda_{\text{eff}}^{(\alpha)}$ from P14-P19 STATUS: Deferred to Section VI (Dark Energy)
5. Tidal tensor effects on pocket structure and evolution STATUS: Formalism established; numerical solutions required for realistic configurations

Required GR Consistency Checks

1. Einstein equations satisfied within each pocket α CHECK: Solutions $g_{\mu\nu}^{(\alpha)}$ must satisfy $G_{\mu\nu}^{(\alpha)} + \Lambda_{\text{eff}}^{(\alpha)} g_{\mu\nu}^{(\alpha)} = (8\pi G/c^4) T_{\mu\nu}^{(\alpha)}$
2. Bianchi identities hold: $\nabla_{\mu} (G^{\mu\nu} + \Lambda_{\text{eff}} g^{\mu\nu}) = 0$ CHECK: Automatic if $G_{\mu\nu}$ computed correctly from $g_{\mu\nu}$; Λ_{eff} variation must be consistent
3. Energy-momentum conservation in each frame: $\nabla_{\mu} T^{\mu\nu} = 0$ CHECK: Must verify for matter distribution in each pocket
4. Positive energy conditions where applicable CHECK: Dominant energy condition (DEC), weak energy condition (WEC) must hold for ordinary matter; may be violated during exotic collision states (P37)

5. Asymptotic flatness at large distances CHECK: $g_{\mu\nu}^{\alpha} \rightarrow \eta_{\mu\nu}$ as $r \rightarrow \infty$
in each pocket's frame

Required Conservation Law Checks

1. Local energy-momentum conservation in each pocket frame α
2. Angular momentum conservation during pocket interactions and collisions
3. Mass conservation (baryon number conservation) during thermalization
4. Charge conservation (if charge asymmetries present)
5. Entropy evolution during collisions: $dS_{\text{total}}/dt \geq 0$ (addressed in Section V)

SECTION IV: GEOMETRIC STRUCTURE OF NESTED SPACETIME POCKETS

This section describes the geometric framework underlying Successive Collision Theory using standard General Relativity concepts: worldlines, reference frames, proper time, and metric structure. We identify what must be mathematically proven and what consistency checks are required.

SUBSECTION IV.1: DEFINING SPACETIME POCKETS

From Premises P7-P8, P11:

A spacetime pocket is a comoving frame, a collection of gravitationally bound or kinematically associated objects sharing approximate collective motion, considered as a spacetime region with shared metric properties.

FORMAL DEFINITION:

Let $\{x_i^\mu(\tau)\}$ be worldlines of N objects (galaxies, stars, gas clouds) in 4D spacetime with coordinates $x^\mu = (ct, x, y, z)$.

Pocket α is defined as the set: $\alpha = \{x_i^\mu : |u_i^\mu - \langle u^\mu \rangle_\alpha| < \varepsilon_v\}$

where: • $u_i^\mu = dx_i^\mu/d\tau_i$ is four-velocity of object i • $\langle u^\mu \rangle_\alpha = (1/N) \sum_i u_i^\mu$ is average four-velocity in pocket α • ε_v is velocity dispersion threshold (defines “comoving” criterion)

TIED TO PREMISES: • P7: Objects “follow the leader” → small velocity dispersion around leader’s motion defines pocket membership • P8: Nested frames emerge from GR applied to eternal infinite spacetime • P11: Each comoving frame treated as “pocket” of spacetime

MATHEMATICAL REQUIREMENTS: \triangle Must specify ε_v criterion: -

Physical: $\varepsilon_v \sim$ virial velocity $v_{\text{virial}} = \sqrt{GM/R}$ for gravitationally bound systems - Kinematic: $\varepsilon_v \sim 1\sigma$ velocity dispersion for kinematic groups

\triangle Must define pocket boundaries: - Sharp boundary (ε_v threshold) or smooth transition (weight function)? - Overlapping pockets allowed?

(Objects can belong to multiple pockets at different scales)

\triangle Must verify pocket definition is coordinate-independent (covariant): - $u^\mu u_\mu = -c^2$ maintained for all objects - Pocket membership invariant under Lorentz transformations

SUBSECTION IV.2: NESTED SUCCESSION HIERARCHY

From Premises P7-P8, P10-P11:

Pockets organize hierarchically: smaller pockets nest within larger pockets, forming a succession extending from smallest gravitationally bound systems to arbitrarily large scales.

HIERARCHICAL STRUCTURE:

Define nesting relation: $\alpha \subset \beta$ (pocket α nested within pocket β) if: 1. Spatial containment: All objects in α lie within spatial extent of β 2. Velocity hierarchy: $\varepsilon_v(\alpha) < \varepsilon_v(\beta)$ (tighter comoving criterion in child) 3. Scale hierarchy: $R_\alpha < R_\beta$ (α smaller than β)

Example hierarchy (our observable location): Solar System \subset Milky Way \subset Local Group \subset Virgo Supercluster \subset Laniakea \subset larger structure $\subset \dots \rightarrow \infty$

POCKET DEPTH INDEX:

Define n_α = number of parent pockets above α in succession.

For observational convenience, set $n = 0$ at some reference scale (e.g., our observable universe). Then: • $n < 0$: child pockets (galaxies, stars, planets within our observable universe) • $n = 0$: our observable universe pocket • $n > 0$: parent pockets (larger structures containing our observable universe)

TIED TO PREMISES: • P7: Scale-invariant follow-the-leader \rightarrow nesting extends indefinitely • P8: Nested succession of larger and larger comoving frames • P10: Time inheritance through succession \rightarrow depth n matters • P2-P3: Infinite space \rightarrow no maximum nesting level ($n \rightarrow \infty$ allowed)

MATHEMATICAL REQUIREMENTS: \triangle Must prove nesting relation is transitive: If $\alpha \subset \beta$ and $\beta \subset \gamma$, then $\alpha \subset \gamma$

\triangle Must verify scale hierarchy consistency: $R_\alpha < R_\beta < R_\gamma$ for $\alpha \subset \beta \subset \gamma$
 $M_\alpha < M_\beta < M_\gamma$ (typically, with exceptions for compact objects)

\triangle Must address infinite regress (P1-P3): - Is there a "root" pocket (n_{\max}) or does succession extend infinitely upward? - If infinite: do observables converge as $n \rightarrow \infty$? - Answer from P2-P3: succession extends infinitely; observables must have well-defined statistical properties (P6)

CONSISTENCY CHECKS: ✓ Verify nesting preserves causality: light cones don't close ✓ Verify hierarchical structure consistent with observed large-scale structure (galaxy groups → clusters → superclusters → filaments)

SUBSECTION IV.3: METRIC STRUCTURE IN POCKETS

From Premises P8, P11:

Each pocket α has associated metric tensor $g_{\mu\nu}^{\alpha}$ describing spacetime geometry within that pocket.

METRIC ANSATZ:

For pocket α , the metric can be written (in appropriate coordinates):

$$g_{\mu\nu}^{\alpha} = \eta_{\mu\nu} + h_{\mu\nu}^{\alpha}$$

where: • $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$ is Minkowski metric (flat spacetime background per P6 at large scales) • $h_{\mu\nu}^{\alpha}$ represents curvature from matter/energy in pocket α and contributions from parent pockets

PERTURBATION STRUCTURE:

The perturbation $h_{\mu\nu}^{\alpha}$ includes: 1. Local curvature: $h_{\mu\nu}^{\alpha}(\text{local})$ from matter distribution within α 2. Parent influence: $h_{\mu\nu}^{\alpha}(\text{parent})$ ($\beta \rightarrow \alpha$) from embedding in parent pocket β 3. Sibling influence: $h_{\mu\nu}^{\alpha}(\text{sibling})$ (α_i) from nearby sibling pockets (typically small)

Total: $h_{\mu\nu}^{\alpha} = h_{\mu\nu}^{\alpha}(\text{local}) + h_{\mu\nu}^{\alpha}(\text{parent}) + h_{\mu\nu}^{\alpha}(\text{sibling}) + \dots$

TIED TO PREMISES: • P8: Nested frames → metric includes parent influence • P11: Each pocket is spacetime region → has metric • P6: Large-scale limit → Minkowski ($\eta_{\mu\nu}$) • P13E: Each pocket has gravitational field → encoded in $h_{\mu\nu}$

MATHEMATICAL REQUIREMENTS: \triangle CRITICAL DERIVATION NEEDED (Section XV Tier 2): Must specify functional form of $h_{\mu\nu}^{\wedge}(\text{parent } \beta \rightarrow \alpha)$: - How does parent pocket β 's curvature, matter distribution, and motion influence metric in child pocket α ? - Is influence tidal (quadrupole field)? Monopole correction? Both? - How does influence scale with distance from parent center?

\triangle Must solve Einstein field equations in pocket α : $G_{\mu\nu}^{\wedge}(\alpha) + \Lambda_{\text{eff}}^{\wedge}(\alpha) g_{\mu\nu}^{\wedge}(\alpha) = (8\pi G/c^4) F(T_{\mu\nu}^{\wedge}(\alpha), M_{\text{bodies}}^{\wedge}(\alpha))$

where: - $G_{\mu\nu}^{\wedge}(\alpha) =$ Einstein tensor computed from $g_{\mu\nu}^{\wedge}(\alpha) - \Lambda_{\text{eff}}^{\wedge}(\alpha)$ from P14-P17 (Tier 1 derivation required) - $F(T_{\mu\nu}, M_{\text{bodies}})$ from P40-P43 (Tier 1 derivation required)

\triangle Must verify weak-field limit: For $h_{\mu\nu} \ll 1$: Einstein equations \rightarrow Poisson equation $\nabla^2 \Phi = 4\pi G \rho$ (Newtonian gravity recovered)

CONSISTENCY CHECKS: \checkmark Signature: $g_{\mu\nu}^{\wedge}(\alpha)$ must have Lorentzian signature $(-, +, +, +)$ everywhere \checkmark Regularity: $g_{\mu\nu}^{\wedge}(\alpha)$ must be smooth (C^2 or better) except at singularities (which P54 prevents) \checkmark Asymptotic flatness: $\lim(r \rightarrow \infty) g_{\mu\nu}^{\wedge}(\alpha) \rightarrow \eta_{\mu\nu}$ (far from matter) \checkmark Energy conditions: Which does $T_{\mu\nu}^{\wedge}(\alpha)$ satisfy? - Weak energy condition: $T_{\mu\nu} u^{\wedge\mu} u^{\wedge\nu} \geq 0$ for all timelike $u^{\wedge\mu}$ - Dominant energy condition: $T_{\mu\nu} u^{\wedge\mu}$ is non-spacelike - Strong energy condition: $(T_{\mu\nu} - \frac{1}{2}T g_{\mu\nu}) u^{\wedge\mu} u^{\wedge\nu} \geq 0$ Answer depends on whether Λ_{eff} and $F(T_{\mu\nu}, M_{\text{bodies}})$ preserve conditions

SUBSECTION IV.4: TRANSFORMATIONS BETWEEN POCKETS

From Premises P8, P9, P10, P12:

Observers in different pockets measure different proper times and spatial intervals due to relative motion and gravitational potentials. Transformations $\Lambda^{\mu}_{\nu}(\alpha \rightarrow \beta)$ relate coordinates between pockets.

TRANSFORMATION TENSOR:

For pockets α nested in β , define transformation:

$$x^{\mu}(\beta) = \Lambda^{\mu}_{\nu}(\alpha \rightarrow \beta) x^{\nu}(\alpha) + b^{\mu}(\alpha \rightarrow \beta)$$

where: • $\Lambda^{\mu}_{\nu}(\alpha \rightarrow \beta)$ is 4×4 transformation matrix • $b^{\mu}(\alpha \rightarrow \beta)$ is offset vector (centers of mass typically don't coincide)

COMPONENTS OF Λ^{μ}_{ν} :

The transformation includes: 1. Lorentz boost: from relative velocity $v_{\alpha\beta}$ between pocket centers 2. Spatial rotation: if pocket axes not aligned 3. Gravitational redshift factor: from potential difference $\Delta\phi = \phi_{\beta} - \phi_{\alpha}$

Explicitly (to leading order):

$$\Lambda^0_0 = \gamma_{\alpha\beta} \sqrt{g_{00}(\beta) / g_{00}(\alpha)} \quad \Lambda^i_j = R^i_j \text{ (rotation matrix)} \quad \Lambda^0_i = -\gamma_{\alpha\beta} (v_{\alpha\beta})^i / c$$

$$\Lambda^i_0 = -\gamma_{\alpha\beta} (v_{\alpha\beta})_i / c$$

where $\gamma_{\alpha\beta} = 1/\sqrt{1 - v_{\alpha\beta}^2/c^2}$ is Lorentz factor.

TIED TO PREMISES: • P8: Nested frames \rightarrow transformations connect them
 • P9: Each frame has own time/space perception $\rightarrow \Lambda^{\mu}_{\nu}$ needed • P10: Time is hereditary $\rightarrow \Lambda^0_0$ component critical • P12: Individual velocities and potentials refine transformation

MATHEMATICAL REQUIREMENTS: \triangle Must specify $\Lambda^{\mu}_{\nu}(\alpha \rightarrow \beta)$ explicitly:

- Full functional form including gravitational contributions - How does $g_{\mu\nu}(\alpha)$ and $g_{\mu\nu}(\beta)$ enter? - Weak-field approximation sufficient or need exact form?

△ Must verify composition rule: $\Lambda^{\mu}_{\nu}(\alpha \rightarrow \gamma) = \Lambda^{\mu}_{\rho}(\beta \rightarrow \gamma) \Lambda^{\rho}_{\nu}(\alpha \rightarrow \beta)$ for $\alpha \subset \beta \subset \gamma$ (Transformations compose associatively through succession)

△ Must verify metric preservation or controlled modification: $g_{\mu\nu}^{\beta} = \Lambda^{\rho}_{\mu}(\alpha \rightarrow \beta) \Lambda^{\sigma}_{\nu}(\alpha \rightarrow \beta) g_{\rho\sigma}^{\alpha} + \text{corrections}$ What are corrections? (Parent influence $h_{\mu\nu}^{\text{parent}}$)

△ Must verify inverse exists: $\Lambda^{\mu}_{\nu}(\beta \rightarrow \alpha) = [\Lambda^{\nu}_{\mu}(\alpha \rightarrow \beta)]^{-1}$

CONSISTENCY CHECKS: ✓ Lorentz invariance preserved within each pocket ✓ Four-velocity magnitude preserved: $u^{\mu} u_{\mu} = -c^2$ after transformation ✓ Causality: timelike curves remain timelike, null remain null, spacelike remain spacelike ✓ No closed timelike curves created by composition $\Lambda(\alpha \rightarrow \beta \rightarrow \gamma \rightarrow \dots \rightarrow \alpha)$

SUBSECTION IV.5: WORLDLINES AND GEODESICS IN NESTED GEOMETRY

From Premises P7, P12:

Objects follow geodesics (or nearly geodesics with local forces) in their local pocket metric $g_{\mu\nu}^{\alpha}$. “Follow-the-leader” means smaller objects approximately follow geodesics in the effective spacetime curved by larger objects.

GEODESIC EQUATION IN POCKET α :

$$d^2x^{\mu}/d\tau^2 + \Gamma^{\mu}_{\nu\rho}(\alpha) (dx^{\nu}/d\tau)(dx^{\rho}/d\tau) = f^{\mu}_{\text{external}} / m$$

where: • $\Gamma^{\mu}_{\nu\rho}(\alpha)$ are Christoffel symbols computed from $g_{\mu\nu}^{\alpha}$ • $f^{\mu}_{\text{external}}$ = non-gravitational forces (electromagnetic, pressure, etc.) • For “free-fall” objects: $f^{\mu}_{\text{external}} = 0$ (pure geodesic motion)

LEADER-FOLLOWER DYNAMICS:

In pocket β containing leader mass M_{leader} and follower objects $\{m_i\}$: • Leader's worldline $x^\mu_{\text{leader}}(\tau)$ approximately defines pocket β 's trajectory • Follower geodesics in effective metric: $g_{\mu\nu}^{\text{(eff)}} \approx$ Schwarzschild metric of M_{leader} + tidal corrections • Followers orbit leader or stream toward it (gravitational attraction)

TIED TO PREMISES: • P7: Follow-the-leader process \rightarrow smaller masses follow geodesics in leader's field • P12: Individual trajectories through gravitational fields refine inherited time perception • P8: GR field equations govern at all scales

MATHEMATICAL REQUIREMENTS: Δ Must show "follow-the-leader" emerges from geodesic equation: - Derive effective potential for followers in leader's field - Show stable orbits exist (Kepler problem in GR) - Verify hierarchical stability: perturbations from parent pocket don't disrupt child pocket structure

Δ Must compute Christoffel symbols: $\Gamma^\mu_{\nu\rho}(\alpha) = \frac{1}{2} g^{\mu\sigma}(\alpha) [\partial_\nu g_{\sigma\rho}(\alpha) + \partial_\rho g_{\nu\sigma}(\alpha) - \partial_\sigma g_{\nu\rho}(\alpha)]$ from $g_{\mu\nu}^{\text{(eff)}}(\alpha)$ specified in Subsection IV.3

CONSISTENCY CHECKS: \checkmark Equivalence principle: freely falling observers in pocket α experience local inertial frames \checkmark Geodesic deviation: tidal forces from parent pocket appear as spacetime curvature in child pocket \checkmark Energy-momentum conservation along worldlines: $\nabla_\mu T^{\mu\nu} = 0$

SUBSECTION IV.6: POCKET BOUNDARIES AND TRANSITION REGIONS

From Premises P11, P8:

Pockets don't have sharp boundaries. Transition regions exist where objects belong partially to pocket α and partially to parent β .

BOUNDARY CHARACTERIZATION:

Define pocket α 's core region: $R_{\text{core}}(\alpha) =$ region where $\geq 90\%$ of α 's mass resides and velocity dispersion $\sigma_v < \varepsilon_v$

Define transition region: $R_{\text{transition}}(\alpha \leftrightarrow \beta) =$ region where objects have velocities intermediate between $\langle u^\mu \rangle_\alpha$ and $\langle u^\mu \rangle_\beta$

Example: Milky Way (α) embedded in Local Group (β):

- Core: $r < 30$ kpc (Milky Way disk and halo)
- Transition: $30 \text{ kpc} < r < 500 \text{ kpc}$ (Magellanic Stream, diffuse satellites)
- Parent domain: $r > 500 \text{ kpc}$ (Local Group medium)

TIED TO PREMISES:

- P11: Pockets are spacetime regions with approximate properties
- P8: Nested frames blend smoothly (no discontinuities)

MATHEMATICAL REQUIREMENTS: Δ Must define membership function: $w_\alpha(x^\mu) =$ weight of object at x^μ belonging to pocket α Properties: $0 \leq w_\alpha \leq 1$, $\sum_{\text{pockets}} w_\alpha = 1$ at each point

Δ Must show metric transitions smoothly: $g_{\mu\nu}(x)$ blends from $g_{\mu\nu}^\wedge(\alpha)$ in core to $g_{\mu\nu}^\wedge(\beta)$ in parent No discontinuities in $g_{\mu\nu}$ or derivatives $\partial_\rho g_{\mu\nu}$

CONSISTENCY CHECKS: \checkmark No singularities at boundaries (contrary to "bubble universe" models with domain walls) \checkmark Observable properties (density, velocity dispersion) vary smoothly across transitions

SUBSECTION IV.7: OBSERVABLE UNIVERSE AS POCKET AT SPECIFIC DEPTH

From Premises P3, P11, P24, P32, P47-P52:

Our observable universe (particle horizon ~ 46 Gly, comoving) is one pocket at specific nesting level $n = n_{\text{obs}}$ in the infinite succession.

OBSERVATIONAL POCKET PROPERTIES:

Set $n_{\text{obs}} = 0$ for convenience. Then: • Pocket radius: $R_{\text{obs}} \approx 46$ Gly (comoving horizon) • Mass: $M_{\text{obs}} \approx 10^{23} M_{\odot} \approx 2 \times 10^{53}$ kg (observable baryonic + inferred “dark” matter; consistent with the $\sim 10^{53}$ kg value quoted in P11/P13, expressed here in solar masses for one consistent convention) • Velocity dispersion: $\sigma_v(\text{obs}) \approx 600$ km/s (large-scale flows, Hubble flow residual) • Age: $t_{\text{obs}} \approx 13.8$ Gyr (time since collision sequence creating this pocket, per P23-P24)

PARENT POCKET ($n = +1$):

Our parent pocket β contains: • Our observable universe α plus siblings (N_{siblings} per P32, P48-P52) • Estimated radius: $R_{\text{parent}} \sim 10^3$ to $10^5 R_{\text{obs}}$ (speculative, observationally inaccessible) • Estimated mass: $M_{\text{parent}} \sim 10^3$ to $10^5 M_{\text{obs}}$

TIED TO PREMISES: • P24: Our Big Bang created infinitesimal patch relative to larger structure • P32: Siblings exist in same parent pocket • P47: Not created in isolation • P48-P52: Low probability of being unique; likely siblings and cousins

MATHEMATICAL REQUIREMENTS: Δ Must estimate parent pocket properties from observables: - CMB dipole ($v_{\text{CMB}} \approx 370$ km/s) \rightarrow our motion in what frame? Parent pocket? Grandparent? - Large-scale flows ($v_{\text{bulk}} \approx 600$ km/s toward Great Attractor) \rightarrow parent motion signature? - CMB low-multipole alignments \rightarrow parent frame rotation axis (P13A)?

Δ Must relate observables to pocket depth: - Redshift z as function of $\Delta n = n_{\text{source}} - n_{\text{obs}}$ (Section VIII) - Λ_{eff} dependence on n_{obs} (Section VII) - Anisotropies from parent frame properties (P63)

OBSERVATIONAL COMMITMENTS: • CMB rest frame defines approximate parent pocket frame • Large-scale structure alignments trace parent/sibling geometry (Section XII) • Hubble tension may reflect calibration differences between n-levels (Section VII)

SUBSECTION IV.8: GEOMETRY SUMMARY AND OPEN MATHEMATICAL TASKS

SUMMARY:

SCT's geometric structure consists of: 1. Spacetime pockets: comoving frames with shared metric properties 2. Nested succession: hierarchical embedding $\alpha \subset \beta \subset \gamma \subset \dots$ extending infinitely 3. Pocket metrics: $g_{\mu\nu}^{\alpha} = \eta_{\mu\nu} + h_{\mu\nu}^{\text{local}} + h_{\mu\nu}^{\text{parent}} + \dots$ 4. Transformations: $\Lambda^{\mu}_{\nu}(\alpha \rightarrow \beta)$ including Lorentz boost + gravitational redshift 5. Geodesic motion: follow-the-leader via GR equations 6. Smooth transitions: no sharp boundaries between pockets 7. Observable universe: one pocket at depth $n_{\text{obs}} = 0$

CRITICAL MATHEMATICAL TASKS (must complete for geometric foundation):

TIER 2, POCKET GEOMETRY (Section XV): 1. Specify $g_{\mu\nu}^{\alpha}$ including parent influence $h_{\mu\nu}^{\text{parent}}(\beta \rightarrow \alpha)$ Dependencies: Matter distribution $T_{\mu\nu}^{\alpha}$, parent properties $(M_{\beta}, R_{\beta}, v_{\beta})$ Output: Explicit metric in suitable coordinates

2. Specify $\Lambda^{\mu}_{\nu}(\alpha \rightarrow \beta)$ transformation tensor Dependencies: Relative velocity $v_{\alpha\beta}$, potential difference $\Delta\phi_{\alpha\beta}$ Output: 4×4 matrix with gravitational and kinematic components Verification: Composition rule, metric compatibility, inverse exists

3. Solve modified Einstein equations in pocket α : $G_{\mu\nu}^{\alpha} + \Lambda_{\text{eff}}^{\alpha}$
 $g_{\mu\nu}^{\alpha} = (8\pi G/c^4) F(T_{\mu\nu}^{\alpha}, M_{\text{bodies}}^{\alpha})$ Dependencies: Λ_{eff} from
P14-P17 (Tier 1), F from P40-P43 (Tier 1) Output: Self-consistent solution
for $g_{\mu\nu}^{\alpha}$

SECTION V: RELATIVISTIC INHERITANCE OF TIME AND SPACE PERCEPTION

This section derives how proper time and spatial measurement standards propagate through the nested pocket succession via Special Relativity's time dilation and length contraction combined with General Relativity's gravitational redshift. We show what is derivable from premises and identify gaps requiring mathematical completion.

SUBSECTION V.1: PROPER TIME IN A SINGLE POCKET

From Premises P9, P11:

Observers at rest in pocket α measure proper time τ_{α} along their worldlines according to the metric $g_{\mu\nu}^{\alpha}$ defined in Section IV.

STANDARD GR PROPER TIME:

For worldline $x^{\mu}(\lambda)$ in pocket α with metric $g_{\mu\nu}^{\alpha}$:

$$d\tau_{\alpha} = \sqrt{-g_{\mu\nu}^{\alpha} dx^{\mu} dx^{\nu}}$$

For observer at rest in pocket α 's rest frame ($dx^i = 0$):

$$d\tau_{\alpha} = \sqrt{-g_{00}^{\alpha}} dt$$

where t is coordinate time in pocket α .

GRAVITATIONAL TIME DILATION:

For weak-field metric $g_{00}^{\alpha} \approx -(1 + 2\Phi_{\alpha}/c^2)$:

$$d\tau_{\alpha}/dt = \sqrt{1 + 2\Phi_{\alpha}/c^2} \approx 1 + \Phi_{\alpha}/c^2$$

where $\Phi_\alpha < 0$ is gravitational potential (negative for attractive gravity).

Observer deep in gravitational well (more negative Φ_α) experiences slower proper time relative to coordinate time.

TIED TO PREMISES: • P9: Each frame has own perception of time • P11: Each pocket has metric $g_{\mu\nu}(\alpha)$ • Standard GR: No new physics

MATHEMATICAL STATUS: ✓ DERIVABLE from GR given $g_{\mu\nu}(\alpha)$ Δ

Requires $g_{\mu\nu}(\alpha)$ from Section IV.3 (Tier 2 task)

SUBSECTION V.2: KINEMATIC TIME DILATION BETWEEN POCKETS

From Premises P9, P12:

Objects moving with velocity v relative to pocket α 's rest frame experience Special Relativity time dilation.

LORENTZ TIME DILATION:

For object moving with velocity v^i in pocket α :

$$d\tau_{\text{moving}}/d\tau_{\text{rest}} = 1/\gamma = \sqrt{1 - v^2/c^2}$$

where $\gamma = 1/\sqrt{1 - v^2/c^2}$ is Lorentz factor.

Moving observer's clock runs slower: $d\tau_{\text{moving}} < d\tau_{\text{rest}}$ for $v > 0$.

POCKET-TO-POCKET KINEMATIC EFFECT:

Consider pocket α (child) moving with velocity $v_{\alpha\beta}$ relative to pocket β (parent) rest frame. Observer at rest in α has velocity $v_{\alpha\beta}$ in β 's frame:

$$d\tau_\alpha/d\tau_\beta|_{\text{kinematic}} = \sqrt{1 - v_{\alpha\beta}^2/c^2}$$

TYPICAL VALUES:

- Solar System (α) moving in Milky Way (β): $v_{\alpha\beta} \sim 220 \text{ km/s} \rightarrow \gamma \approx 1.0000003 \rightarrow \Delta\tau/\tau \sim 10^{-7}$ (tiny effect)
- Galaxy (α) in cluster (β): $v_{\alpha\beta} \sim 500\text{-}1000 \text{ km/s} \rightarrow \gamma \approx 1.000001$ to $1.000006 \rightarrow \Delta\tau/\tau \sim 10^{-6}$ to 10^{-5}
- Cluster (α) in supercluster (β): $v_{\alpha\beta} \sim 300\text{-}600 \text{ km/s} \rightarrow$ similar magnitude

CUMULATIVE EFFECT:

Through N nesting levels with velocities v_1, v_2, \dots, v_N :

$$d\tau_{\text{child}}/d\tau_{\text{root}} = \prod_{i=1}^N \sqrt{1 - v_i^2/c^2}$$

For small $v_i \ll c$:

$$\ln(d\tau_{\text{child}}/d\tau_{\text{root}}) \approx -\frac{1}{2} \sum_i v_i^2/c^2$$

Even for $N = 10$ levels with $v_i \sim 500 \text{ km/s}$: $\Delta\tau/\tau \sim 10 \times (500 \text{ km/s} / c)^2 \sim 10 \times 10^{-6} \sim 10^{-5}$

TIED TO PREMISES: • P9: Motion through space slows motion through time (explicit SR citation) • P12: Individual velocities refine inherited time perception • Standard SR: No new physics

MATHEMATICAL STATUS: ✓ DERIVABLE from SR given relative velocities $v_{\alpha\beta}$ Δ Requires specification of $v_{\alpha\beta}$ from pocket dynamics (Section IV.4)

SUBSECTION V.3: GRAVITATIONAL TIME DILATION BETWEEN POCKETS

From Premises P9, P10, P12, P13E:

Observer in child pocket α sits deeper in parent pocket β 's gravitational potential well, experiencing additional gravitational time dilation beyond local effects.

GRAVITATIONAL REDSHIFT FACTOR:

For observer at gravitational potential Φ_α in pocket α and Φ_β in parent pocket β :

$$d\tau_\alpha/d\tau_\beta|_{\text{gravitational}} = \sqrt{g_{00}^{(\alpha)}/g_{00}^{(\beta)}}$$

$$\text{In weak-field limit: } g_{00}^{(\alpha)} \approx -(1 + 2\Phi_\alpha/c^2) \quad g_{00}^{(\beta)} \approx -(1 + 2\Phi_\beta/c^2)$$

$$\text{Therefore: } d\tau_\alpha/d\tau_\beta|_{\text{gravitational}} \approx \sqrt{[(1 + 2\Phi_\alpha/c^2)/(1 + 2\Phi_\beta/c^2)]} \approx 1 + (\Phi_\alpha - \Phi_\beta)/c^2 = 1 + \Delta\Phi_{\alpha\beta}/c^2$$

where $\Delta\Phi_{\alpha\beta} = \Phi_\alpha - \Phi_\beta$ is potential difference.

INTERPRETATION:

If child pocket α sits in deeper potential well (Φ_α more negative), then $\Delta\Phi_{\alpha\beta} < 0$ and $d\tau_\alpha < d\tau_\beta \rightarrow$ time runs slower in child pocket.

CUMULATIVE GRAVITATIONAL EFFECT:

Through nested succession with potential differences $\Delta\Phi_1, \Delta\Phi_2, \dots, \Delta\Phi_N$:

$$d\tau_{\text{child}}/d\tau_{\text{root}}|_{\text{gravitational}} = \prod_i \exp(\Delta\Phi_i/c^2) \approx \exp(\sum_i \Delta\Phi_i/c^2)$$

$$\text{Total potential difference: } \Delta\Phi_{\text{total}} = \sum_i \Delta\Phi_i$$

ESTIMATION:

For pocket at distance R from parent's center of mass M_{parent} : $\Phi_{\text{pocket}} \approx -GM_{\text{parent}}/R$

Potential difference between child at R_{child} and parent edge at R_{parent} :

$$\Delta\Phi \approx GM_{\text{parent}}(1/R_{\text{child}} - 1/R_{\text{parent}})$$

Example: Observable universe ($M_{\text{obs}} \sim 10^{23} M_\odot$, $R_{\text{obs}} \sim 46 \text{ Gly}$)

embedded at distance $R \sim 1000 R_{\text{obs}}$ in parent pocket ($M_{\text{parent}} \sim 10^{26} M_\odot$):

$$\Delta\Phi \sim G(10^{26} M_\odot)(1/1000 R_{\text{obs}} - 1/R_{\text{parent}})$$

If $R_{\text{parent}} \gg R_{\text{obs}}$: $\Delta\Phi \sim G(10^{26} M_\odot)/(10^3 \times 46 \text{ Gly})$

Converting: $\Delta\Phi/c^2 \sim (G M_{\text{parent}})/(c^2 R) \sim (\text{Schwarzschild radius})/(2R) \sim (10^{26} M_{\odot} \times 3 \text{ km}/M_{\odot})/(2 \times 10^3 \times 46 \text{ Gly}) \sim (3 \times 10^{29} \text{ km})/(10^2 \times 46 \times 10^9 \text{ ly} \times 10^{13} \text{ km/ly}) \sim 10^{-8} \text{ to } 10^{-7}$ (small effect per level)

Through $N \sim 5\text{-}10$ parent levels: cumulative $\Delta\Phi_{\text{total}}/c^2 \sim 10^{-7} \text{ to } 10^{-5}$

TIED TO PREMISES: • P9: Each frame has own time perception • P10: Time is hereditary, child inherits from parent modified by environment • P12: Trajectories through gravitational fields refine time • P13E: Each pocket has gravitational field • Standard GR: gravitational time dilation

MATHEMATICAL STATUS: ✓ DERIVABLE from GR given $g_{\mu\nu}^{\alpha}$, $g_{\mu\nu}^{\beta}$ Δ Requires pocket metrics from Section IV.3 Δ Requires specification of pocket positions within parent ($R_{\text{child}}/R_{\text{parent}}$)

SUBSECTION V.4: HEREDITARY TIME, COMBINING EFFECTS

From Premise P10 (critical premise for time inheritance):

Time is hereditary: each pocket inherits base proper time from parent, modified by kinematic and gravitational effects.

COMBINED TIME DILATION FORMULA:

For pocket α nested in parent pocket β :

$$d\tau_{\alpha}/d\tau_{\beta} = \sqrt{(1 - v_{\alpha}^2/c^2)} \times \sqrt{[(1 + 2\Phi_{\alpha}/c^2)/(1 + 2\Phi_{\beta}/c^2)]}$$

$$\text{Combining: } d\tau_{\alpha}/d\tau_{\beta} \approx (1 - v_{\alpha}^2/2c^2) \times (1 + \Delta\Phi_{\alpha\beta}/c^2) \approx 1 - v_{\alpha}^2/2c^2 + \Delta\Phi_{\alpha\beta}/c^2 + O(v^4/c^4, v^2\Phi/c^4)$$

Dropping higher-order terms:

$$d\tau_{\alpha} \approx d\tau_{\beta} [1 - v_{\alpha}^2/2c^2 + \Delta\Phi_{\alpha\beta}/c^2]$$

RECURSIVE INHERITANCE:

For nested succession child → parent → grandparent → ...:

$$d\tau_{\text{child}} = d\tau_{\text{parent}} [1 - v_{\text{child-parent}}^2/2c^2 + \Delta\Phi_{\text{child-parent}}/c^2]$$

$$d\tau_{\text{parent}} = d\tau_{\text{grandparent}} [1 - v_{\text{parent-grandparent}}^2/2c^2 + \Delta\Phi_{\text{parent-grandparent}}/c^2]$$

...

Composing:

$$d\tau_{\text{child}} = d\tau_{\text{root}} \times \prod_{i=1}^N [1 - v_i^2/2c^2 + \Delta\Phi_i/c^2]$$

For small corrections:

$$\ln(d\tau_{\text{child}}/d\tau_{\text{root}}) \approx \sum_{i=1}^N [-v_i^2/2c^2 + \Delta\Phi_i/c^2]$$

$$d\tau_{\text{child}}/d\tau_{\text{root}} \approx \exp[\sum_i (-v_i^2/2c^2 + \Delta\Phi_i/c^2)] \approx 1 + \sum_i (-v_i^2/2c^2 + \Delta\Phi_i/c^2)$$

TIED TO PREMISES: • P10: Time is hereditary (CORE PREMISE for this section) • P9: Motion and gravity affect time (mechanism) • P12: Individual refinements compound through succession

MATHEMATICAL STATUS: ✓ DERIVABLE in principle from SR + GR △

CRITICAL GAP: Functional form needs rigorous derivation △ Must specify:

Does “hereditary time” mean more than just cumulative SR+GR effects? △

If additional effect beyond standard relativity: MUST SPECIFY and JUSTIFY from premises

SUBSECTION V.5: LORENTZ TRANSFORMATIONS THROUGH NESTED SUCCESSION

From Premises P8, P9, P10, P12:

When comparing observations between pockets that are NOT directly nested (e.g., two galaxies in different galaxy clusters), Lorentz

transformations cannot be applied directly. Instead, transformations must be composed through the nested succession by first identifying a common ancestor pocket.

CRITICAL MODIFICATION TO STANDARD SR PROTOCOL:

In standard Special Relativity between two inertial frames A and B: • Direct Lorentz transformation $\Lambda(A \rightarrow B)$ connects observations • Assumes flat spacetime and direct kinematic relationship

In Successive Collision Theory with nested pockets: • Pockets α and ω may not share direct nesting relationship • Transformations must traverse through succession hierarchy • REQUIRES: Identify lowest common ancestor pocket β

PATH-FINDING ALGORITHM:

To compare observations between pockets α and ω :

STEP 1: Identify lowest common ancestor pocket β - Trace α 's parent chain: $\alpha \rightarrow \text{parent}_1(\alpha) \rightarrow \text{parent}_2(\alpha) \rightarrow \dots \rightarrow \beta$ - Trace ω 's parent chain: $\omega \rightarrow \text{parent}_1(\omega) \rightarrow \text{parent}_2(\omega) \rightarrow \dots \rightarrow \beta$ - β is first pocket appearing in both chains

STEP 2: Compose transformations UP from α to β $\Lambda(\alpha \rightarrow \beta) = \Lambda(\text{parent}_n \rightarrow \beta) \circ \dots \circ \Lambda(\text{parent}_1 \rightarrow \beta) \circ \Lambda(\alpha \rightarrow \text{parent}_1)$

STEP 3: Compose transformations DOWN from β to ω $\Lambda(\beta \rightarrow \omega) = \Lambda(\text{parent}_1 \rightarrow \omega)^{-1} \circ \Lambda(\text{parent}_2 \rightarrow \omega)^{-1} \circ \dots \circ \Lambda(\text{parent}_m \rightarrow \omega)^{-1}$

STEP 4: Complete transformation $\Lambda(\alpha \rightarrow \omega) = \Lambda(\beta \rightarrow \omega) \circ \Lambda(\alpha \rightarrow \beta)$

EXAMPLE:

Consider comparing time scales between: • α = Solar System (in Milky Way, in Local Group, in Virgo Supercluster) • ω = System in Andromeda (also in Local Group, in Virgo Supercluster)

Common ancestor β = Local Group (both are members)

Transformation chain: Solar System \rightarrow Milky Way \rightarrow Local Group \rightarrow
Andromeda \rightarrow Target System

$\Lambda(\text{Solar} \rightarrow \text{Target}) = \Lambda(\text{LG} \rightarrow \text{Andromeda})^{-1} \circ \Lambda(\text{Milky Way} \rightarrow \text{LG}) \circ$
 $\Lambda(\text{Solar} \rightarrow \text{Milky Way})$

Each component includes: • Lorentz boost from relative velocity •
Gravitational redshift factor from potential difference • Spatial rotation if
coordinate axes not aligned

MATHEMATICAL FORMULATION:

For pockets at depths n_α and n_ω in succession hierarchy, with common
ancestor at depth n_β (where $n_\beta < n_\alpha$ and $n_\beta < n_\omega$):

$\Lambda^{\mu\nu}(\alpha \rightarrow \omega) = [\prod_{i=n_\omega}^{n_\beta+1} \Lambda^{\mu\rho}(i \rightarrow i-1)] \circ [\prod_{j=n_\alpha}^{n_\beta+1}$
 $\Lambda^{\rho\nu}(j \rightarrow j-1)]^{-1}$

This is NOT simply $\Lambda(v_{\alpha\omega})$ where $v_{\alpha\omega}$ is relative velocity between α and
 ω .

WHY THIS MATTERS:

1. TIME SCALE COMPARISONS: When computing $d\tau_\alpha/d\tau_\omega$, cannot use
direct formula. Must compute: $d\tau_\alpha/d\tau_\omega = (d\tau_\alpha/d\tau_\beta) \times (d\tau_\beta/d\tau_\omega) =$
 $(d\tau_\alpha/d\tau_\beta) / (d\tau_\omega/d\tau_\beta)$

2. VELOCITY COMPOSITIONS: Velocities don't add linearly. For pockets α
 $\rightarrow \beta \rightarrow \omega$: $v_{\alpha\omega} \neq v_{\alpha\beta} + v_{\beta\omega}$

Instead, relativistic velocity addition: $v_{\alpha\omega} = (v_{\alpha\beta} + v_{\beta\omega}) / (1 + v_{\alpha\beta}$
 $v_{\beta\omega}/c^2)$

But in curved spacetime with nested succession, even this is modified by
gravitational contributions from intermediate pockets.

3. NON-COMMUTATIVITY OF BOOSTS: Successive Lorentz boosts in non-parallel directions produce Wigner rotation: $\Lambda(v_1) \circ \Lambda(v_2) \neq \Lambda(v_2) \circ \Lambda(v_1)$ when v_1 not parallel to v_2

Cumulative rotation angle through succession: $\theta_{\text{Wigner}} =$ accumulated rotation from non-collinear velocity changes

This could produce OBSERVABLE ANISOTROPIES in redshift measurements if parent pocket velocities vary by direction (see P52).

4. PATH DEPENDENCE: Question: Does transformation $\Lambda(\alpha \rightarrow \omega)$ depend on which common ancestor β is chosen?

Answer: Should NOT depend (due to associativity of transformations), BUT:

- Numerical precision issues in long chains - Possible systematic effects if pocket boundaries fuzzy (Section IV.6) - Requires verification: composition rule associative through all paths

TIED TO PREMISES: • P8: Nested succession of frames requires hierarchical transformations • P9: Time/space perception differs between pockets \rightarrow transformations needed • P10: Time is hereditary through succession \rightarrow must trace inheritance path • P12: Individual velocities and potentials \rightarrow each link in chain matters

OBSERVATIONAL IMPLICATIONS:

1. REDSHIFT CALCULATIONS (Section VIII): For photon from source in pocket α to observer in pocket ω :

- Must trace photon path through succession hierarchy
- Accumulate redshift contributions from each pocket boundary crossing
- Cannot use simple $z = v/c$ formula

2. VELOCITY MEASUREMENTS: Peculiar velocities of galaxies must account for:

- Parent frame motions (Local Group, Virgo, Laniakea, ...)
- Inherited velocities through succession
- Gravitational contributions from all parent pockets along path

3. ANISOTROPY PREDICTIONS: If parent frame has bulk velocity v_{parent} in grandparent frame (P63):

- Sources at different angles experience different accumulated boosts
- Predicted dipole pattern in redshift-distance relation
- Testable with precision surveys

MATHEMATICAL STATUS: ✓ ALGORITHM SPECIFIED: Path-finding through succession tree ✓ COMPOSITION RULE STATED: $\Lambda(\alpha \rightarrow \omega)$ via common ancestor Δ REQUIRES VERIFICATION: - Associativity: independent of path choice - Convergence: finite result for infinite succession ($n_{\beta} \rightarrow \infty$) - Wigner rotation accumulation: observably significant? Δ REQUIRES FROM SECTION IV: - Explicit $\Lambda^{\mu}_{\nu}(\alpha \rightarrow \beta)$ functional form - Pocket hierarchy structure (parent relationships) - Velocity and potential data for each link

SUBSECTION V.6: SPATIAL MEASUREMENT

INHERITANCE

From Premises P9 (length contraction), P12:

Just as time perception is inherited, spatial measurement standards are also modified through nested succession via SR length contraction and GR spatial curvature.

LORENTZ LENGTH CONTRACTION:

Object of rest length L_0 in pocket α , measured by observer in pocket β moving at velocity $v_{\alpha\beta}$:

$$L_{\text{measured}} = L_0 \sqrt{1 - v_{\alpha\beta}^2/c^2} = L_0/\gamma_{\alpha\beta}$$

Length contracts along direction of relative motion.

GRAVITATIONAL SPATIAL DISTORTION:

In curved spacetime with metric g_{ij} (spatial part):

$$dl^2 = g_{ij} dx^i dx^j$$

For Schwarzschild metric (example): $g_{rr} = (1 - 2GM/rc^2)^{-1}$

Radial distances stretched near massive object.

CUMULATIVE SPATIAL EFFECT:

Through nested succession, spatial scales transform via:

$$dx^i_{\text{(child)}} = [\Lambda^i_j(\text{parent} \rightarrow \text{child})] dx^j_{\text{(parent)}} \times [1 + \text{curvature corrections}]$$

Following path-finding algorithm from Subsection V.5, spatial measurements between non-nested pockets must also traverse common ancestor.

TIED TO PREMISES: • P9: Motion through space affects space perception (SR length contraction) • P12: Trajectories through gravitational fields refine space perception

MATHEMATICAL STATUS: ✓ DERIVABLE from SR + GR transformations Δ
Requires $\Lambda^\mu_\nu(\alpha \rightarrow \beta)$ from Section IV.4 Δ Less critical than time inheritance for redshift (Section VIII focuses on time)

SUBSECTION V.7: CONNECTION TO COSMOLOGICAL REDSHIFT

From Premises P9, P10, P13I:

Cumulative proper time differences across nested pocket succession are the foundation for cosmological redshift in SCT.

PHOTON FREQUENCY AND TIME:

Photon frequency ν relates to proper time via:

$$\nu = d\phi/d\tau$$

where ϕ is phase of electromagnetic wave.

For photon emitted in source pocket α with frequency ν_{source} (measured in α 's proper time τ_{α}) and received in observer pocket β with frequency ν_{obs} (measured in β 's proper time τ_{β}):

$$\nu_{\text{obs}}/\nu_{\text{source}} = d\tau_{\alpha}/d\tau_{\beta}$$

Redshift defined as:

$$z = (\nu_{\text{source}} - \nu_{\text{obs}})/\nu_{\text{obs}} = \nu_{\text{source}}/\nu_{\text{obs}} - 1$$

Therefore:

$$\mathbf{1 + z = \nu_{\text{source}}/\nu_{\text{obs}} = d\tau_{\beta}/d\tau_{\alpha}}$$

INTERPRETATION:

If observer's proper time runs faster than source's ($d\tau_{\beta} > d\tau_{\alpha}$), then photon is redshifted ($z > 0$).

From Subsection V.4: if source is in deeper nested pocket succession (more parent levels with cumulative time dilation), then $d\tau_{\text{source}} < d\tau_{\text{observer}} \rightarrow z > 0$.

CUMULATIVE REDSHIFT THROUGH SUCCESSION:

For photon traveling from source in pocket at depth n_{source} to observer at depth n_{obs} , passing through intermediate pockets:

$$1 + z_{\text{total}} = (d\tau_{\text{obs}}/d\tau_{\text{source}})_{\text{cumulative}} = \prod_{i=n_{\text{source}}}^{\{n_{\text{obs}}\}} (d\tau_{i+1}/d\tau_i)$$

From Subsection V.4:

$$1 + z_{\text{total}} \approx \exp[-\sum_i (v_i^2/2c^2 - \Delta\Phi_i/c^2)]$$

For $n_{\text{obs}} > n_{\text{source}}$ (observer in shallower pocket succession): $\sum_i \Delta\Phi_i < 0$ (cumulative deeper potential) $\sum_i v_i^2/2c^2 > 0$ (cumulative kinematic)

Net effect depends on whether gravitational or kinematic dominates.

CRITICAL NOTE ON PATH TRACING:

Following Subsection V.5, redshift calculation between source in pocket α and observer in pocket ω requires:

1. Identify photon path through succession hierarchy
2. Determine common ancestor pocket β (or multiple ancestors if path complex)
3. Accumulate time dilation factors along path:

$$1 + z = (d\tau_{\omega}/d\tau_{\beta}) / (d\tau_{\alpha}/d\tau_{\beta}) = \prod(\text{path } \beta \rightarrow \omega) (d\tau_{i+1}/d\tau_i) / \prod(\text{path } \beta \rightarrow \alpha) (d\tau_{j+1}/d\tau_j)$$

This is NOT simply a function of distance d , but depends on: • Specific pocket path through succession hierarchy • Velocities along each segment • Gravitational potentials at each pocket • Accumulated Wigner rotations (directional effects)

TIED TO PREMISES: • P10: Hereditary time differences → frequency shift • P13I: Acquired perception of space/time from parent succession (KEY for redshift) • P9: Mechanism via SR + GR

MATHEMATICAL STATUS: ✓ CONNECTION ESTABLISHED between hereditary time and redshift Δ CRITICAL TIER 1 DERIVATION REQUIRED (Section VIII): Must derive explicit $z = f(n_{\text{source}}, n_{\text{obs}}, \text{pocket parameters, path})$ Must show statistical emergence of Hubble law $z \propto d$ Must predict deviations from Λ CDM at high z

SUBSECTION V.8: OBSERVATIONAL CONSEQUENCES OF TIME INHERITANCE

From Premises P10, P13I:

If time perception is inherited through nested succession with cumulative modifications, several observational consequences follow:

CONSEQUENCE 1: REDSHIFT-DISTANCE RELATION

Deeper pockets (larger n) → slower proper time → larger redshift when observed from shallower pocket.

If distance d correlates with depth difference $\Delta n = n_{\text{source}} - n_{\text{obs}}$: $z \propto \Delta n \propto d$ (statistical Hubble law)

Derivation required (Section VIII).

CONSEQUENCE 2: CLOCK RATE VARIATIONS

Atomic clocks at different nesting levels would tick at different rates if compared directly.

GPS satellites already show this (gravitational + kinematic): $\Delta\tau/\tau \sim 10^{-9}$ per day (corrected in GPS algorithms)

Extended to cosmic scales: Clocks in different galaxy clusters may have $\Delta\tau/\tau \sim 10^{-5}$ to 10^{-4} differences from different parent frame embeddings.

Currently unobservable (can't synchronize clocks across Mpc scales), but principle established.

CONSEQUENCE 3: ANISOTROPIES FROM PARENT FRAME

If our observable universe has velocity v_{obs} relative to parent frame (P63), and different source directions correspond to different relative velocities in parent frame, then:

$z(\text{direction})$ varies with angle

Dipole anisotropy in redshift-distance relation: $\Delta z/z \sim (v_{\text{obs}}/c) \cos(\theta) \sim 10^{-3} \times \cos(\theta)$

for $v_{\text{obs}} \sim 600$ km/s (observed large-scale flow).

TESTABLE with future precision redshift surveys.

TIED TO PREMISES: • P10: Time is hereditary • P13I: Acquired time perception • P52: Our frame has velocity in parent frame

MATHEMATICAL STATUS: ✓ QUALITATIVE predictions identified Δ
QUANTITATIVE predictions require Tier 1 derivations (Sections VII, VIII)

SUBSECTION V.9: CRITICAL GAPS AND DERIVATION REQUIREMENTS

WHAT WE HAVE DERIVED:

✓ Proper time in single pocket: $d\tau = \sqrt{-g_{\mu\nu} dx^{\mu} dx^{\nu}}$, STANDARD GR ✓
Kinematic time dilation between pockets: factor $\sqrt{1 - v^2/c^2}$, STANDARD SR
✓ Gravitational time dilation between pockets: factor $\sqrt{g_{00}^{\alpha/g_{00}\beta}}$,
STANDARD GR ✓ Combined effect: $d\tau_{\alpha}/d\tau_{\beta} \approx 1 - v^2/2c^2 + \Delta\Phi/c^2$,

DERIVABLE from SR+GR ✓ Connection to redshift: $1 + z = d\tau_{\text{obs}}/d\tau_{\text{source}}$, STANDARD GR ✓ Path-finding algorithm through succession tree (Subsection V.5) ✓ Transformation composition via common ancestor

WHAT REQUIRES COMPLETION:

△ TIER 2 DERIVATION (Section XV):

HEREDITARY TIME FUNCTIONAL FORM

P10 states “time is hereditary” and “each pocket inherits base proper time from parent and passes refined version to child.”

Question: Is this MORE than cumulative SR+GR effects derived above, or IS IT EXACTLY the cumulative SR+GR?

Option A: Hereditary time = cumulative SR+GR (no new physics) Then:

$\tau_{\text{child}} = \tau_{\text{parent}} \times [1 - v^2/2c^2 + \Delta\Phi/c^2]$ as derived Status: ✓ COMPLETE (just standard relativity applied recursively)

Option B: Hereditary time includes additional effect beyond SR+GR Then: MUST SPECIFY what additional effect is MUST JUSTIFY from premises why SR+GR insufficient MUST DERIVE modified formula Status: △ NOT SPECIFIED, would require new physics (violates premise constraint “only GR+SR”)

RESOLUTION: Adopt Option A (hereditary time = cumulative SR+GR). P10’s “hereditary” language describes the RECURSIVE APPLICATION of standard relativistic effects through nested succession, not a new physical mechanism.

Action: Specify explicitly in Section XV that: $\tau_{\alpha}(\tau_{\beta}, v_{\alpha\beta}, \Delta\Phi_{\alpha\beta}) = \tau_{\beta} \times [1 - v_{\alpha\beta}^2/2c^2 + \Delta\Phi_{\alpha\beta}/c^2]$ is the COMPLETE functional form satisfying P10.

△ TIER 1 DERIVATION (Section VIII, Redshift):

Must derive $z = f(n_{\text{source}}, n_{\text{obs}}, \{v_i\}, \{\Delta\Phi_i\}, \text{pocket_path})$

Given cumulative time dilation formula from above, must: 1. Integrate along photon path through nested pockets 2. Sum contributions from each pocket boundary crossing 3. Account for path through common ancestor (Subsection V.5 algorithm) 4. Show statistical average yields Hubble-like relation $z \approx H_0 d/c$ 5. Identify deviations from Λ CDM (anisotropies, high- z departures)

This is CRITICAL for testability.

△ VERIFICATION REQUIREMENTS (Section XV):

TRANSFORMATION COMPOSITION ASSOCIATIVITY

Must prove that $\Lambda(\alpha \rightarrow \omega)$ is independent of path choice through succession tree: - For any two paths from α to ω through different common ancestors - Result should be identical (associativity of composition) - If path-dependent: physical meaning? (different light paths?)

WIGNER ROTATION ACCUMULATION

Must compute accumulated rotation from successive non-collinear boosts: - $\theta_{\text{total}} = \sum_i \theta_{\text{Wigner}}(v_i, v_{\{i+1\}}, \text{angle between})$ - Is this observably significant? ($\sim 10^{-6}$ rad cumulative?) - Produces quadrupole anisotropy pattern?

CONVERGENCE FOR INFINITE SUCCESSION

For $n_{\beta} \rightarrow \infty$ (common ancestor infinitely far up succession): - Does $\Lambda(\alpha \rightarrow \omega)$ converge to finite limit? - Does cumulative time dilation converge? - Physical interpretation if divergent

△ **REQUIRES FROM SECTION IV:**

- Pocket metrics $g_{\mu\nu}^{\alpha}$ • Transformation tensors $\Lambda^{\mu}_{\nu}(\alpha \rightarrow \beta)$ • Relative velocities $v_{\alpha\beta}$ from pocket dynamics • Potential differences $\Delta\Phi_{\alpha\beta}$ from pocket gravitational fields • Hierarchical structure (parent-child relationships)

Without these, hereditary time formula incomplete.

SUBSECTION V.10: SUMMARY, RELATIVISTIC INHERITANCE

ESTABLISHED IN THIS SECTION:

- ✓ Proper time in each pocket governed by metric $g_{\mu\nu}^{\alpha}$ (standard GR)
- ✓ Time dilation between pockets combines SR kinematic (γ factor) and GR gravitational ($\sqrt{g_{00}^{\alpha/g_{00}^{\beta}}}$) effects

- ✓ Hereditary time P10 realized as recursive application of SR+GR time dilation through nested succession:

$$d\tau_{\text{child}}/d\tau_{\text{parent}} = \sqrt{1 - v^2/c^2} \times \sqrt{g_{00}^{\text{child}/g_{00}^{\text{parent}}}}$$

- ✓ Cumulative time differences through N nesting levels:

$$d\tau_{\text{deepest}}/d\tau_{\text{shallowest}} = \exp[\sum_i (-v_i^2/2c^2 + \Delta\Phi_i/c^2)]$$

- ✓ Connection to redshift established: $1 + z = d\tau_{\text{obs}}/d\tau_{\text{source}}$

- ✓ CRITICAL ADDITION: Lorentz transformations between non-nested pockets require path-finding through succession hierarchy via common ancestor (Subsection V.5)

- Cannot apply direct transformation between arbitrary pockets • Must compose transformations: $\alpha \rightarrow \text{parent} \rightarrow \dots \rightarrow \text{ancestor} \rightarrow \dots \rightarrow \omega$ • Produces accumulated effects: non-commutativity, Wigner rotation • Enables anisotropic predictions from parent frame velocity

✓ Observational predictions identified: - Redshift-distance relation from depth correlation - Anisotropies from parent frame velocity (P63) - Path-dependent redshift effects - Possible clock rate variations (unobservable at current precision)

TIED TO PREMISES: • P9: Motion affects time (SR), gravity affects time (GR) • P10: Time is hereditary (recursive SR+GR) • P12: Individual velocities and potentials refine inheritance • P13I: Acquired time perception from parent succession • P8: Nested succession structure enables hierarchical transformations

CRITICAL DEPENDENCIES: • Section IV pocket geometry ($g_{\mu\nu}$, $\Lambda^{\mu\nu}$, $v_{\alpha\beta}$, $\Delta\Phi_{\alpha\beta}$) • Section VIII redshift derivation (z from cumulative $d\tau$ along paths) • Section XV verification of transformation associativity and convergence

NO NEW PHYSICS INTRODUCED: ✓ All formulas derive from standard SR and GR ✓ “Hereditary time” is conceptual framework, not new equation ✓ Path-finding algorithm is consequence of nested structure, not new physics ✓ Novelty is in APPLICATION (recursive through infinite nested succession, with transformation composition via common ancestors), not in mechanism (which is standard relativity)

REMAINING TASKS: \triangle Complete pocket geometry (Section IV Tier 2) \triangle Derive explicit redshift formula (Section VIII Tier 1) \triangle Verify transformation composition properties (Section XV) \triangle Specify pocket parameters from observations

SECTION IV: RELATIVISTIC INHERITANCE OF PROPER TIME AND SPACETIME PERCEPTION

The nested hierarchy of spacetime pockets established in Section III creates a fundamental consequence: each pocket inherits its baseline proper-time behavior from its parent pocket and passes a refined version to its child objects. This section formalizes the mechanism of hereditary time transmission (P10), derives the mathematical framework for proper-time inheritance through Special and General Relativistic effects (P9, P12), and identifies critical derivations required to connect inherited proper time to observable cosmological redshift.

IV.A, CONCEPTUAL FOUNDATION: TIME AS HEREDITARY (P9, P10, P12)

Premise Foundation

P9 establishes that each comoving frame possesses a shared perception of time arising from Special Relativistic time dilation: objects moving together at velocity v relative to a parent frame experience the same time dilation factor $\gamma = 1/\sqrt{1 - v^2/c^2}$, creating a common “clock rate” for that frame.

P10 introduces the hereditary principle: proper-time behavior propagates through the nested hierarchy like a recursive function, with each pocket inheriting a baseline from its parent and transmitting a modified version to its children.

P12 specifies the refinement mechanism: within a pocket, individual objects have local velocities and gravitational trajectories that further adjust the inherited proper-time baseline before passing it to child objects.

Operational Meaning

An observer in pocket α experiences time flow at rate $d\tau^{(\alpha)}/dt$ determined by: 1. Cumulative Lorentz factors from all parent pockets ($\alpha+1$, $\alpha+2$, $\alpha+3$, ...) 2. Cumulative gravitational redshift from all parent

gravitational potentials 3. Local refinements from motion and gravity within pocket α itself

This creates a proper-time “pedigree” where the observer’s clock rate encodes the entire history of nested frame motions and gravitational environments.

Connection to Observables

Hereditary proper-time differences between distant observers (in different pockets or at different depths in the hierarchy) manifest as cosmological redshift, replacing Λ CDM’s metric expansion interpretation with a proper-time differential interpretation (detailed in Section VII).

IV.B, SPECIAL RELATIVISTIC TIME DILATION ACROSS NESTED SUCCESSION

Single-Level Time Dilation (P9)

Consider observer in pocket α moving at velocity $v^{(\alpha)}$ relative to parent pocket $\alpha+1$. Standard SR time dilation:

$$d\tau^{(\alpha)} = d\tau^{(\alpha+1)} \sqrt{1 - v^{(\alpha)2}/c^2} = d\tau^{(\alpha+1)} / \gamma^{(\alpha)}$$

where: • $d\tau^{(\alpha)}$ is proper time interval measured by observer in pocket α •

$d\tau^{(\alpha+1)}$ is proper time interval measured by observer at rest in pocket

$\alpha+1$ • $\gamma^{(\alpha)} = 1/\sqrt{1 - \beta^{(\alpha)2}}$ with $\beta^{(\alpha)} = v^{(\alpha)}/c$ is Lorentz factor

DERIVATION FROM SR: Direct consequence of Lorentz transformation invariance of spacetime interval $ds^2 = -c^2dt^2 + dx^2 + dy^2 + dz^2$. For observer moving at velocity v in x-direction:

$$ds^2 = -c^2dt'^2 = -c^2dt^2 + v^2dt^2 = -c^2dt^2 (1 - v^2/c^2)$$

$$\rightarrow dt' = dt \sqrt{1 - v^2/c^2} = dt / \gamma$$

This is standard SR; no new physics invoked.

Multi-Level Cumulative Time Dilation (P10)

For observer in pocket α embedded within succession $\alpha+1, \alpha+2, \alpha+3, \dots$, proper time accumulates through recursive application:

$$d\tau^{\alpha} = d\tau^{\alpha+1} / \gamma^{\alpha} \quad d\tau^{\alpha+1} = d\tau^{\alpha+2} / \gamma^{\alpha+1} \quad d\tau^{\alpha+2} = d\tau^{\alpha+3} / \gamma^{\alpha+2} \dots$$

Composing these relations:

$$\begin{aligned} d\tau^{\alpha} &= d\tau^{\alpha+n} \times \prod_{k=0}^{n-1} [1 / \gamma^{\alpha+k}] \\ &= d\tau^{\alpha+n} \times \prod_{k=0}^{n-1} \sqrt{1 - \beta^{2\alpha+k}} \end{aligned}$$

This shows proper-time ratio between pocket α and distant ancestor pocket $\alpha+n$ depends on product of all intermediate Lorentz factors.

MATHEMATICAL COMMITMENT: The infinite product $\prod_{k=0}^{\infty} \sqrt{1 - \beta^{2\alpha+k}}$ must converge for hereditary time to be well-defined. This requires:

$$\sum_{k=0}^{\infty} \beta^{2\alpha+k} < \infty$$

Physically reasonable if relative velocities decrease at higher hierarchical levels ($\beta^{\alpha+k} \rightarrow 0$ as $k \rightarrow \infty$), consistent with large-scale homogeneity (P6).

DERIVABILITY REQUIREMENT: Must prove convergence from hierarchical dynamics and verify consistency with observed velocity dispersions at different scales.

Velocity Composition Across Levels

Relativistic velocity addition for successive frames:

$$v_{\text{total}} = (v_1 + v_2) / (1 + v_1 v_2 / c^2)$$

For small velocities ($v \ll c$), approximately additive: $v_{\text{total}} \approx v_1 + v_2$

For large velocities approaching c , saturates at c due to denominator.

In nested succession with N levels, total relative velocity between innermost and outermost frame:

$$v_{\text{rel}}(\alpha, \alpha+N) = v^{(\alpha)} \oplus v^{(\alpha+1)} \oplus \dots \oplus v^{(\alpha+N-1)}$$

where \oplus denotes relativistic velocity addition.

MATHEMATICAL REQUIREMENT: Must compute cumulative velocity composition and verify that it remains subluminal for all locally accelerated objects (P20), while permitting superluminal relative velocities between independently formed pockets (P21).

IV.C, GRAVITATIONAL TIME DILATION ACROSS NESTED SUCCESSION

Single-Potential Gravitational Redshift

Observer in gravitational potential $\Phi^{(\alpha)}$ experiences gravitational time dilation relative to observer at infinity (or in weaker potential):

$$d\tau_{\text{grav}} = dt \sqrt{g_{00}} \approx dt \sqrt{1 + 2\Phi/c^2} \approx dt (1 + \Phi/c^2)$$

where $g_{00} = -(1 + 2\Phi/c^2)$ is time-time component of metric in weak-field limit.

For observer deep in potential well $\Phi < 0$ (negative potential), proper time runs slower: $d\tau < dt$.

DERIVATION FROM GR: Schwarzschild metric for spherically symmetric mass M:

$$ds^2 = -(1 - 2GM/rc^2) c^2 dt^2 + (1 - 2GM/rc^2)^{-1} dr^2 + r^2 d\Omega^2$$

At fixed spatial position ($dr = d\Omega = 0$), proper time element:

$$d\tau = \sqrt{g_{00}} dt = \sqrt{1 - 2GM/rc^2} dt$$

For $r \gg r_s$ (Schwarzschild radius), weak-field approximation:

$$\Phi = -GM/r \rightarrow d\tau \approx (1 + \Phi/c^2) dt$$

Standard GR result; no new physics.

Multi-Potential Cumulative Gravitational Redshift (P10, P13)

Observer in pocket α experiences gravitational potentials from: 1. Local potential $\Phi_{\text{local}}^{\alpha}$ within pocket α 2. Parent potential $\Phi^{\alpha+1}$ from parent pocket 3. Grandparent potential $\Phi^{\alpha+2}$ from grandparent pocket 4. All higher-level potentials $\Phi^{\alpha+3}$, $\Phi^{\alpha+4}$, ...

Total gravitational time dilation (weak-field limit):

$$d\tau_{\text{grav}}^{\alpha} \approx dt [1 + \Phi_{\text{total}}^{\alpha}/c^2]$$

where:

$$\Phi_{\text{total}}^{\alpha} = \Phi_{\text{local}}^{\alpha} + \Phi^{\alpha+1} + \Phi^{\alpha+2} + \dots = \sum_{k=\alpha}^{\infty} \Phi^k$$

MATHEMATICAL COMMITMENT: The infinite sum $\sum_{k=\alpha}^{\infty} \Phi^k$ must converge. Physical expectation: gravitational potential from distant pockets decreases as $\Phi^k \propto -GM^{(k)}/r(k)$, and if pocket separations grow faster than masses, convergence is ensured.

DERIVABILITY REQUIREMENT: Must prove convergence from hierarchical structure (scaling relations from Section III) and verify that total potential remains finite.

Combination: SR + GR Time Dilation

Combining Special Relativistic velocity-induced time dilation and General Relativistic gravitational time dilation:

$$\begin{aligned} d\tau^{\alpha} &= dt \times [\text{kinematic factor}] \times [\text{gravitational factor}] \\ &= dt \times \sqrt{1 - v_{\text{total}}^2/c^2} \times (1 + \Phi_{\text{total}}/c^2) \end{aligned}$$

where: • v_{total} is cumulative velocity from all parent motions • Φ_{total} is cumulative gravitational potential from all parent structures

More precisely, for observer in pocket α :

$$d\tau^{\alpha} = d\tau^{\alpha+n} \times \prod_{k=0}^{n-1} \sqrt{1 - \beta^2(\alpha+k)} \times \exp[\sum_{k=\alpha}^{\alpha+n-1} \Phi^k/c^2]$$

Taking $n \rightarrow \infty$ (comparing to “absolute rest” at infinite hierarchical distance):

$$d\tau^{\alpha} = d\tau_{\infty} \times [\prod_{k=0}^{\infty} \sqrt{1 - \beta^2(\alpha+k)}] \times \exp[\sum_{k=\alpha}^{\infty} \Phi^k/c^2]$$

MATHEMATICAL REQUIREMENT: Both infinite product and infinite sum must converge for hereditary proper time to be well-defined.

IV.D, LOCAL REFINEMENTS WITHIN POCKET (P12)

Individual Object Trajectories

Premise P12 specifies that objects within pocket α possess individual velocities v_{local} and gravitational trajectories that refine the inherited baseline proper time before transmission to child objects.

Observer at position x within pocket α experiences: 1. Inherited baseline $\tau_{inherited}^{\alpha}$ from parent succession (Section IV.B, IV.C) 2. Local velocity correction from motion relative to pocket center-of-mass 3. Local gravitational correction from pocket’s own gravitational field

Total proper time:

$$d\tau_{observer} = d\tau_{inherited}^{\alpha} \times \sqrt{1 - v_{local}^2/c^2} \times (1 + \Phi_{local}(x)/c^2)$$

where: • $v_{local} = |v_{observer} - V_{CM}^{\alpha}|$ is velocity relative to pocket rest frame • $\Phi_{local}(x)$ is gravitational potential at observer position x within pocket α

Time Differential Across Pocket

Two observers at different positions x_1, x_2 within same pocket α experience proper-time differential:

$$\Delta\tau / \tau \approx [v_1^2 - v_2^2] / (2c^2) + [\Phi(x_1) - \Phi(x_2)] / c^2$$

For observers in different gravitational environments (e.g., surface of galaxy vs intergalactic void):

$$\Delta\Phi \sim GM_{\text{galaxy}} / R_{\text{galaxy}} \sim 10^{-6} c^2$$

$$\rightarrow \Delta\tau / \tau \sim 10^{-6}$$

This is small but measurable over cosmological timescales (billions of years).

OBSERVATIONAL COMMITMENT: Differential proper-time accumulation across large-scale structure could contribute to observed redshift scatter and peculiar velocities beyond simple Hubble flow.

Transmission to Child Objects

When observer forms child pocket (e.g., star system forms within galaxy), the child inherits observer's refined proper time as its new baseline:

$$\tau_{\text{baseline}}^{(\alpha-1)} = \tau_{\text{observer}}^{(\alpha)}$$

Child pocket then applies its own local refinements:

$$\tau^{(\alpha-1)} = \tau_{\text{baseline}}^{(\alpha-1)} \times [\text{local corrections for pocket } \alpha-1]$$

This creates multi-generational proper-time pedigree encoding entire ancestry of nested motions and gravitational environments.

IV.E, MATHEMATICAL FORMALISM: PROPER-TIME RECURSION RELATION

General Recursion Formula (P10)

Combining all effects (SR velocity, GR gravity, local refinements), proper-time relation between pocket α and parent $\alpha+1$:

$$d\tau^{\wedge}(\alpha) = d\tau^{\wedge}(\alpha+1) \times K^{\wedge}(\alpha)$$

where $K^{\wedge}(\alpha)$ is total modification factor:

$$K^{\wedge}(\alpha) = \sqrt{(1 - \beta^2(\alpha))} \times \exp[\Phi^{\wedge}(\alpha)/c^2] \times [1 + \text{local corrections}]$$

For pocket embedded n levels deep:

$$d\tau^{\wedge}(\alpha) = d\tau^{\wedge}(\alpha+n) \times \prod_{k=0}^{n-1} K^{\wedge}(\alpha+k)$$

Differential Form

Converting to differential equation for proper time as function of hierarchical depth:

$$d\tau^{\wedge}(\alpha) / d\tau^{\wedge}(\alpha+1) = K^{\wedge}(\alpha)$$

Taking logarithm:

$$d \ln \tau^{\wedge}(\alpha) / d\alpha = \ln K^{\wedge}(\alpha)$$

If $K^{\wedge}(\alpha)$ varies slowly with α , can approximate:

$$\tau^{\wedge}(\alpha) \approx \tau^{\wedge}(\alpha+n) \times \exp[\int_{\alpha}^{\alpha+n} \ln K(\alpha') d\alpha']$$

MATHEMATICAL REQUIREMENT: Must specify functional form $K(\alpha)$ from hierarchical dynamics. This requires: 1. Velocity distribution $\beta(\alpha)$ across hierarchy 2. Gravitational potential distribution $\Phi(\alpha)$ across hierarchy 3. Pocket spacing and mass scaling relations

DERIVABILITY GAP: Explicit functional forms $\beta(\alpha)$, $\Phi(\alpha)$ must be derived from N-body gravitational dynamics and large-scale structure formation. This is a major outstanding calculation.

IV.F, INHERITED QUANTITIES BEYOND PROPER TIME

While proper time is the primary inherited quantity (P10), P13 specifies that pockets inherit entire “perception of space and time” from parent succession. This includes:

1. Spatial Metric Inheritance

Spatial distances measured in pocket α depend on inherited metric from parent:

$$d \ell_{\text{measured}}^{\alpha} = \sqrt{g_{ij}^{\alpha} dx^i dx^j}$$

where spatial metric components g_{ij}^{α} are influenced by parent frame motion (length contraction) and parent gravitational field (spatial curvature).

Lorentz contraction from parent motion:

$$\ell_{\text{parallel}} = \ell_{\text{rest}} / \gamma^{\alpha} \text{ (along direction of motion)} \quad \ell_{\text{perpendicular}} = \ell_{\text{rest}} \text{ (perpendicular to motion)}$$

DERIVABILITY REQUIREMENT: Must derive full spatial metric g_{ij}^{α} from parent frame transformations and gravitational fields.

2. Light-Cone Structure Inheritance

The causal light-cone structure (which events can influence which others) is determined by metric $g_{\mu\nu}^{\alpha}$, which is inherited from parent frames.

Null geodesics (light paths) satisfy:

$$ds^2 = g_{\mu\nu}^{\alpha} dx^{\mu} dx^{\nu} = 0$$

Light-cone opening angle and propagation speed in local coordinates depend on inherited metric structure.

GR CONSISTENCY REQUIREMENT: Null geodesics must be continuous across pocket boundaries; no causality violations permitted.

3. Angular Momentum Orientation

Preferred axes and orientations (e.g., rotation axes, orbital planes) in pocket α may inherit alignment from parent pocket structure (P13A).

Observed large-scale alignment of galaxy spins and filament orientations could reflect this inheritance.

OBSERVATIONAL COMMITMENT: Statistical alignment of structures across large scales testable through galaxy shape correlations, cosmic web topology, and CMB anisotropy patterns.

4. Electromagnetic Environment

Magnetic and electric field configurations (P13F, P13G) may be partially inherited from parent pocket fields, then modified by local dynamics.

Primordial magnetic fields in our universe could have originated from parent pocket fields, amplified during collision (P25).

MATHEMATICAL GAP: Transformation of EM fields across nested frames requires full relativistic treatment; explicit derivation not yet completed.

IV.G, CONNECTION TO OBSERVATIONAL COSMOLOGY

Redshift from Hereditary Time (Preview of Section VII)

Two observers at different positions or depths in nested hierarchy measure different proper-time intervals for same physical process (e.g., atomic transition).

Redshift parameter z defined by:

$$1 + z = \lambda_{\text{observed}} / \lambda_{\text{emitted}} = \nu_{\text{emitted}} / \nu_{\text{observed}}$$

If photon emitted at time t_{emit} by source with proper-time rate $d\tau_{\text{source}}$ and observed at time t_{obs} by observer with proper-time rate $d\tau_{\text{obs}}$, then:

$$1 + z = (d\tau_{\text{obs}} / dt) / (d\tau_{\text{source}} / dt) = d\tau_{\text{obs}} / d\tau_{\text{source}}$$

From hereditary time framework:

$$1 + z = [\prod K_{\text{obs}}] / [\prod K_{\text{source}}]$$

where products extend over all nested frames influencing observer vs source.

DERIVABILITY REQUIREMENT: Must show that statistical distribution of $(\prod K_{\text{obs}}) / (\prod K_{\text{source}})$ over cosmological distances produces Hubble-like linear relation $z \propto d$ for nearby sources (Section VII).

CMB Temperature Variations

Hereditary proper-time differences across sky create temperature anisotropies in CMB:

$$\Delta T / T \approx \Delta\tau / \tau \sim [\Delta v^2 / c^2 + \Delta\Phi / c^2]$$

Dipole anisotropy: $\Delta T/T \sim 10^{-3}$ from our motion relative to parent frame (P54) Higher multipoles: $\Delta T/T \sim 10^{-5}$ from density perturbations and peculiar velocities

OBSERVATIONAL COMMITMENT: Predicted anisotropy pattern must match Planck CMB observations; deviations would falsify SCT's hereditary time framework.

IV.H, REQUIRED DERIVATIONS AND CONSISTENCY CHECKS

Required Mathematical Derivations

1. Convergence of infinite products and sums TASK: Prove that $\prod_{k=0}^{\infty} \sqrt{1 - \beta^{2^k}}$ and $\sum_{k=0}^{\infty} \Phi^{(k)}$ converge STATUS: Not yet proven; requires hierarchical velocity and potential scaling

2. Functional form $K(\alpha)$ TASK: Derive explicit $K(\alpha) = \sqrt{(1 - \beta^2(\alpha)) \exp[\Phi(\alpha)/c^2]}$ from dynamics STATUS: Requires solving for $\beta(\alpha)$, $\Phi(\alpha)$ from N-body simulations
3. Proper-time recursion solution TASK: Solve $\tau^\wedge(\alpha) = \tau^\wedge(\alpha+n) \prod K^\wedge(k)$ for arbitrary n , α STATUS: Formal structure established; explicit solution requires $K(\alpha)$
4. Spatial metric $g_{ij}^\wedge(\alpha)$ TASK: Derive full 3-metric from parent frame transformations STATUS: Lorentz transformation structure known; gravitational corrections require calculation
5. Redshift-distance relation $z(d)$ TASK: Derive Hubble law $z \propto d$ from statistical distribution of τ ratios STATUS: Deferred to Section VII; major outstanding derivation

Required GR/SR Consistency Checks

1. Lorentz invariance preservation CHECK: Verify that proper-time transformations preserve spacetime interval ds^2
2. Einstein equations satisfied CHECK: Metrics $g_{\mu\nu}^\wedge(\alpha)$ must satisfy $G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu} = (8\pi G/c^4) T_{\mu\nu}$ in each frame
3. Causality preservation CHECK: Light-cone structure must be consistent across all frames; no closed timelike curves
4. Energy-momentum conservation CHECK: $\nabla_\mu T^\mu{}_\nu = 0$ must hold in each pocket frame
5. Clock synchronization protocol CHECK: Operational definition of “simultaneous” events must be consistent with inherited proper time

Required Observational Consistency Checks

1. CMB dipole amplitude Our velocity relative to parent frame must match observed dipole $\Delta T/T \approx 3.36 \times 10^{-3}$
2. Local Hubble flow Predicted $\tau(z)$ relation must reproduce $H_0 \approx 67-73$ km/s/Mpc
3. Gravitational time dilation tests GPS satellites, pulsar timing, gravitational redshift experiments must be consistent with GR predictions
4. Large-scale velocity dispersions Predicted $\beta(\alpha)$ distribution must match observed bulk flows and peculiar velocities

SECTION V: SUCCESSIVE SUPERLUMINAL COLLISIONS: REPLACING THE HOT DENSE ORIGIN

The Lambda-CDM model attributes the hot dense early universe to backward extrapolation from an initial singularity at $t = 0$, requiring unverified quantum gravity and fine-tuned initial conditions. Successive Collision Theory replaces this singular origin with a cascade of superluminal collisions between pre-existing nested spacetime pockets (P20-P41), producing the hot dense plasma through shock thermalization of existing matter rather than creation ex nihilo. This section formalizes the collision mechanism, analyzes energy and entropy budgets, identifies curvature evolution requirements, and specifies mathematical tasks needed for quantitative predictions.

V.A, CONCEPTUAL FOUNDATION: SUPERLUMINAL COLLISIONS WITHOUT SR VIOLATION

Local vs Global Speed-of-Light Constraint (P20, P21, P22)

Premise P20 establishes that Special Relativity constrains local acceleration: objects accelerated from rest within our nested succession cannot exceed c .

Premise P21 specifies that SR does NOT constrain relative velocities between independently formed immense pockets whose scales exceed ours by factors $\sim 10^4$ - 10^5 . Two such pockets can have relative velocity $v_{rel} > c$ without violating SR locally within either pocket.

Premise P22 confirms that no physical laws are violated when two nested structures intersect with $v_{rel} > 2c$.

Physical Justification from GR

In General Relativity, the speed-of-light constraint applies to *local* measurement within a single coordinate patch. Two distant objects in different coordinate patches can have coordinate velocities exceeding c without causality violation.

Consider two pockets A and B, each with proper metric $g_{\mu\nu}^A$, $g_{\mu\nu}^B$. Within pocket A, all locally measured velocities satisfy $v < c$. Within pocket B, all locally measured velocities satisfy $v < c$. However, the relative velocity between pockets' centers of mass, $v_{rel} = |v_A - v_B|$, can exceed c because:

1. v_A and v_B are measured in different local frames
2. No signal propagates from A to B faster than c in any local frame
3. Causality is preserved: light cones remain timelike in each pocket's metric

This is analogous to cosmological horizons in Λ CDM: distant galaxies recede faster than c due to metric expansion, yet SR is not violated locally.

GR CONSISTENCY REQUIREMENT: Must verify that collision geometry preserves local light-cone structure in each pocket independently. No closed timelike curves or causality violations permitted.

Collision Kinematics (P23)

Premise P23 allows collision velocities far exceeding $2c$ (potentially $3c$, $7c$, $42c$, $67c$), creating extreme kinetic energy regimes:

$$E_{\text{kinetic}} = (1/2) M_{\text{pocket}} v_{\text{rel}}^2$$

For $M_{\text{pocket}} \sim 10^{\{53\}}$ kg (mass of observable universe) and $v_{\text{rel}} \sim 10c$:

$$E_{\text{kinetic}} \sim (1/2) \times 10^{\{53\}} \text{ kg} \times (10 \times 3 \times 10^8 \text{ m/s})^2 \sim 5 \times 10^{\{70\}} \text{ J} \sim 10^3 \times (M_{\text{pocket}} c^2)$$

Collision energy exceeds rest-mass energy by factors 10^3 or more, creating energy densities:

$$\varepsilon_{\text{collision}} \sim E_{\text{kinetic}} / V_{\text{collision}} \sim 10^{\{70\}} \text{ J} / (10^{68} \text{ m}^3) \sim 10^2 \text{ J/m}^3$$

This corresponds to temperatures (via $\varepsilon \sim a_{\text{radiation}} T^4$):

$$T_{\text{collision}} \sim (\varepsilon / a_{\text{radiation}})^{\{1/4\}} \sim 10^{\{30\}} \text{ K}$$

Such temperatures vastly exceed QCD phase transition ($\sim 10^{\{12\}}$ K) and electroweak symmetry breaking ($\sim 10^{\{15\}}$ K), creating quark-gluon plasma and exotic hadron states (P37).

MATHEMATICAL REQUIREMENT: Must compute collision energy deposition, shock heating, and thermalization timescales from hydrodynamic shock equations.

V.B, COLLISION SEQUENCE AND THERMALIZATION (P24-P29, P35-P40)

Multi-Stage Collision Cascade (P29, P35, P36)

Premise P29 specifies that multiple collisions with similar kinetic energies create our observable universe:

- First collision: pockets A and B intersect at $v_{\text{rel}}(1) > c$
- Second collision: debris from stage 1 collides with pocket C

at $v_{\text{rel}(2)} > c$ • Third collision: further intersections at $v_{\text{rel}(3)} > c$ • N-th collision: eventually $v_{\text{rel}(N)} < c$ as system dissipates energy

Premise P35 confirms early stages (1, 2, 3, possibly 4) are superluminal before slowing.

Premise P36 introduces observational degeneracy: we may be observing results of 4th, 5th, or 6th collision stage, not necessarily the first.

Energy Distribution Across Stages

For comparable kinetic energies $E_{\text{kinetic}(n)} \approx E_{\text{kinetic}(n-1)}$, total energy deposited:

$$E_{\text{total}} = \sum_{n=1}^N E_{\text{kinetic}(n)} \approx N \times E_{\text{kinetic}(1)}$$

Distributed over volume $V_{\text{thermalized}} \sim (\text{few} \times R_{\text{pocket}})^3$, this creates approximately homogeneous energy density:

$$\varepsilon_{\text{avg}} \sim N \times E_{\text{kinetic}} / V_{\text{thermalized}}$$

Small-scale inhomogeneities arise from random collision geometry (impact parameters, orientations), producing density fluctuations:

$$\delta\rho / \rho \sim \text{geometric_variations} \sim 10^{-5} \text{ to } 10^{-4}$$

This naturally produces CMB-like temperature fluctuations $\Delta T/T \sim \delta\rho/\rho$ without requiring inflationary quantum fluctuations.

OBSERVATIONAL COMMITMENT: Predicted $\Delta T/T$ spectrum must match Planck observations C_{ℓ} for $\ell = 2$ to 2500.

Thermalization Timescale (P37, P38)

Premise P37 identifies creation of exotic non-equilibrium states during superluminal collisions.

Premise P38 specifies these states persist through multiple collision generations before thermalizing.

Thermalization proceeds through: 1. Shock heating: kinetic \rightarrow thermal energy via shock fronts 2. Particle collisions: establishing kinetic equilibrium 3. Radiation processes: photon-matter coupling 4. Neutrino decoupling: weakly interacting particles escape

Characteristic thermalization time:

$$\tau_{\text{therm}} \sim 1 / (n \sigma v)$$

where n is particle number density, σ is interaction cross-section, v is typical particle velocity.

For quark-gluon plasma: $n \sim 10^{40} \text{ m}^{-3}$, $\sigma \sim 10^{-40} \text{ m}^2$, $v \sim c \rightarrow$

$$\tau_{\text{therm}} \sim 10^{-10} \text{ s}$$

For subsequent hadronization and cooling: $\tau_{\text{cool}} \sim$ few seconds to minutes (depending on expansion rate)

MATHEMATICAL GAP: Detailed numerical simulation of multi-stage thermalization with exotic intermediate states requires relativistic hydrodynamics + QCD equation of state, currently beyond analytical tractability.

Recombination Timescale Modification (P33)

Premise P33 offers two scenarios: (A) Extended thermalization region lowers initial temperature and density, allowing faster recombination ($\sim 380,000$ days or weeks vs years) (B) Extreme short-lived temperatures create exotic states that rapidly thermalize, enabling quick recombination despite high initial densities

Recombination rate determined by:

$$dX_e / dt = -\alpha_B(T) n_e n_p X_e + \beta(T) n_H (1 - X_e)$$

where X_e is ionization fraction, α_B is recombination coefficient, β is photoionization rate.

Effective recombination time:

$$t_{\text{recomb}} \sim 1 / (\alpha_B n_e)$$

For Λ CDM: $n_e \sim 10^6 \text{ m}^{-3}$, $\alpha_B \sim 10^{-19} \text{ m}^3/\text{s} \rightarrow t_{\text{recomb}} \sim 10^{13} \text{ s} \sim 380,000 \text{ yr}$

For SCT scenario (A): larger volume \rightarrow lower $n_e \rightarrow$ faster recombination per atom, but more atoms total; net effect depends on geometric factors.

For SCT scenario (B): rapid thermalization \rightarrow quick cooling \rightarrow faster recombination.

DERIVABILITY REQUIREMENT: Must compute recombination history from collision-based initial conditions and verify consistency with CMB decoupling redshift $z_{\text{dec}} \sim 1100$.

V.C, COLLISION GEOMETRY AND STRUCTURE FORMATION (P31-P34, P55-P56)

Grazing Collisions \rightarrow Rotating Structures (P30)

Impact parameter $b \sim R_{\text{pocket}}$ produces glancing collision. Angular momentum:

$$L = M_{\text{reduced}} \times v_{\text{rel}} \times b$$

where $M_{\text{reduced}} = M_A M_B / (M_A + M_B)$.

After thermalization, rotating plasma forms disk-like or spheroidal structures with rotation velocity:

$$v_{\text{rot}} \sim L / (M R) \sim (M_{\text{reduced}} / M) \times v_{\text{rel}} \times (b / R)$$

For $b \sim R$ and $M_{\text{reduced}} \sim M/2$:

$$v_{\text{rot}} \sim v_{\text{rel}} / 2$$

This creates flat rotation curves: objects at different radii have similar rotational speeds, mimicking dark matter phenomenology (P50-P53).

MATHEMATICAL REQUIREMENT: Must derive rotation curve $v_{\text{rot}}(r)$ from collision angular momentum distribution and gravitational settling dynamics.

Head-On Collisions \rightarrow Filamentary Structures (P31)

Small impact parameter $b \ll R_{\text{pocket}}$ produces direct collision.

Minimal angular momentum:

$$L \sim M v_{\text{rel}} b \rightarrow 0 \text{ as } b \rightarrow 0$$

Energy deposition creates elongated compression along collision axis.

Post-shock material forms filament aligned with collision direction:

$$\text{Length} \sim v_{\text{rel}} \times \tau_{\text{expansion}} \quad \text{Width} \sim \sqrt{(\text{pressure} / \text{density})} \times \tau_{\text{expansion}}$$

For multiple collisions at different orientations (P32), criss-crossing filaments create cosmic web topology: • Superfilaments: remnants of large-scale collisions • Supervoids: regions between collision zones • Nodes: filament intersections with high density

OBSERVATIONAL COMMITMENT: Predicted filament statistics (lengths, widths, orientations, connectivity) must match large-scale structure surveys (SDSS, 2dFGRS, BOSS).

Structure Formation Without Dark Matter (P46, P47)

Premise P46 asserts structure formation proceeds from collision-seeded over-densities without requiring dark matter particles.

Collision creates local density enhancements $\delta\rho/\rho \sim$ geometric variations. Gravitational instability amplifies perturbations:

$\delta\rho / \rho \propto t^\alpha$ (matter-dominated) or $\propto a$ (radiation-dominated)

Growth timescale:

$t_{\text{growth}} \sim 1 / \sqrt{G \rho_{\text{background}}}$

For $\rho_{\text{background}} \sim \text{critical_density} \sim 10^{-26}$ kg/m³:

$t_{\text{growth}} \sim 10^8$ yr

This is shorter than Λ CDM prediction (without dark matter), enabling earlier structure formation consistent with JWST observations of massive galaxies at $z > 10$ (addressed in Section IX).

Premise P47 predicts giant arcs, big rings, and large features as natural collision geometry remnants at scales \sim collision impact scale \sim Gpc.

OBSERVATIONAL COMMITMENT: Observed giant structures (Big Ring 1.3 Gly, Giant Arc 3.3 Gly) must match predicted collision scales.

V.D, ENERGY CONDITIONS AND CONSERVATION LAWS

Energy Budget Accounting

Total energy before collision: $E_{\text{initial}} = E_{\text{kinetic}}(A) + E_{\text{kinetic}}(B) + E_{\text{rest}}(A) + E_{\text{rest}}(B) = (1/2) M_A v_A^2 + (1/2) M_B v_B^2 + M_A c^2 + M_B c^2$

For $v_{\text{rel}} = |v_A - v_B| \gg c$, kinetic energy dominates rest mass: $E_{\text{kinetic}} \sim (1/2) M v_{\text{rel}}^2 \gg M c^2$

After collision: $E_{\text{final}} = E_{\text{thermal}} + E_{\text{radiation}} + E_{\text{kinetic,remnant}} + E_{\text{gravitational}}$

Energy conservation: $E_{\text{initial}} = E_{\text{final}}$

MATHEMATICAL REQUIREMENT: Must account for all energy channels:

1. Thermal energy: $E_{\text{thermal}} = \int \rho c_v T d^3x$ (internal heat)
2. Radiation: $E_{\text{radiation}} = \int u_{\text{radiation}} d^3x$ (photons, neutrinos)
3. Remnant kinetic: material still in bulk motion
4. Gravitational binding: negative contribution from bound structures
5. Gravitational waves: radiated during collision

Conservation check requires: $\Sigma(\text{all energy forms}) = \text{constant}$ (modulo radiation escaping to infinity)

Momentum Conservation

Total momentum before collision: $P_{\text{initial}} = M_A v_A + M_B v_B$

After collision (center-of-mass frame): $P_{\text{final}} = \int \rho v d^3x + P_{\text{radiation}}$

Momentum conservation: $P_{\text{initial}} = P_{\text{final}}$

For symmetric collision ($M_A \approx M_B$, $|v_A| \approx |v_B|$), center-of-mass approximately at rest. Asymmetric collision creates net momentum \rightarrow bulk flow in post-collision remnant.

OBSERVATIONAL COMMITMENT: Observed bulk flows (e.g., “dipole repeller”, Local Group motion) should correlate with collision asymmetry predictions.

Angular Momentum Conservation

Total angular momentum: $L_{\text{initial}} = L_A + L_B + M_A v_A \times r_A + M_B v_B \times r_B$

where first two terms are intrinsic spins, last two are orbital contributions.

After collision: $L_{\text{final}} = \int r \times (\rho v) d^3x + L_{\text{radiation}}$

Angular momentum conservation: $L_{\text{initial}} = L_{\text{final}}$

Collision impact parameter b determines final angular momentum distribution \rightarrow rotation curves, spin alignments, large-scale vorticity.

MATHEMATICAL REQUIREMENT: Must compute $L_{\text{final}}(b, v_{\text{rel}}, M_A, M_B)$ and verify consistency with observed rotation curves and large-scale flows.

Energy Conditions in GR

Einstein's equations permit arbitrary stress-energy tensors $T_{\mu\nu}$ provided they satisfy energy conditions:

1. Weak Energy Condition (WEC): $T_{\mu\nu} u^\mu u^\nu \geq 0$ for all timelike $u^\mu \rightarrow$ Energy density non-negative in all frames
2. Dominant Energy Condition (DEC): $T_{\mu\nu} u^\mu$ is non-spacelike for timelike $u^\mu \rightarrow$ Energy does not flow faster than light
3. Strong Energy Condition (SEC): $(T_{\mu\nu} - (1/2)T g_{\mu\nu}) u^\mu u^\nu \geq 0 \rightarrow$ Gravity is attractive for ordinary matter

During superluminal collision, exotic states (P37) may temporarily violate SEC due to extreme pressure gradients, but WEC and DEC should be preserved.

GR CONSISTENCY CHECK REQUIRED: Verify that collision stress-energy tensor satisfies WEC and DEC. SEC violation is permitted for transient exotic states but must be restored after thermalization.

V.E, ENTROPY PRODUCTION AND SECOND LAW

Entropy Increase During Collision (P25, P37, P38)

Collision converts ordered kinetic energy (low entropy) into disordered thermal energy (high entropy). Entropy increase:

$$\Delta S = S_{\text{final}} - S_{\text{initial}} > 0$$

Initial entropy (before collision): $S_{\text{initial}} \sim k_B \times (\text{number of microstates in kinetic energy}) \sim \text{small (coherent bulk motion)}$

Final entropy (after thermalization): $S_{\text{final}} \sim k_B \times (\text{number of microstates in thermal energy}) \sim k_B (E_{\text{thermal}} / T_{\text{avg}}) \sim \text{large}$

For $E_{\text{thermal}} \sim 10^{\{70\}} \text{ J}$, $T_{\text{avg}} \sim 10^{\{30\}} \text{ K}$: $S_{\text{final}} \sim 10^{\{40\}} k_B \sim 10^{\{17\}} \text{ J/K}$

Entropy per baryon: $s = S / N_{\text{baryon}} \sim 10^{\{17\}} / 10^{\{80\}} \sim 10^{\{-63\}} \text{ J/K per baryon}$

Observable universe today: $s_{\text{CMB}} \sim 10^{\{-23\}} \text{ J/K per baryon (photon entropy dominates)}$

CONSISTENCY CHECK REQUIRED: Must verify that collision-generated entropy matches or exceeds observed CMB entropy, ensuring second law is satisfied.

Entropy Evolution During Thermalization

Multi-stage thermalization (P38) increases entropy stepwise: $S(\text{stage 1}) < S(\text{stage 2}) < S(\text{stage 3}) < \dots < S(\text{final})$

Each collision stage: 1. Adds kinetic energy \rightarrow shock heating 2. Dissipates through viscosity, radiation 3. Increases phase-space volume explored by particles 4. Increases entropy

Total entropy production: $\Delta S_{\text{total}} = \sum_{n=1}^{\{N\}} \Delta S(\text{stage } n)$

As system cools and expands: • Entropy S increases (second law) • Entropy per comoving volume $s = S/V$ remains approximately constant (adiabatic expansion after thermalization) • Entropy per particle s/N remains constant (conserved particle number)

MATHEMATICAL REQUIREMENT: Compute entropy evolution $S(t)$ from thermodynamic relations and verify monotonic increase $dS/dt \geq 0$ throughout collision cascade.

Comparison to Λ CDM Entropy Budget

Λ CDM attributes high CMB entropy to initial conditions at Planck scale (unverified).

SCT produces high entropy through collision thermalization (verified physical process: shock heating).

Entropy per baryon in CMB: $s_{\text{CMB}} / N_{\text{baryon}} \sim (\text{photon number} / \text{baryon number}) \times k_B \sim \eta^{-1} \times k_B \sim (6 \times 10^{-10})^{-1} k_B \sim 10^9 k_B$

Collision must produce this or higher entropy. For $N_{\text{baryon}} \sim 10^{80}$, total entropy: $S_{\text{collision}} \geq 10^{80} \times 10^9 k_B \sim 10^{89} k_B \sim 10^{66} \text{ J/K}$

Energy-entropy relation during thermalization: $S \sim (E / T)$

For $E \sim 10^{70} \text{ J}$, $T \sim 10^{30} \text{ K}$: $S \sim 10^{70} / (10^{30} \text{ K} \times k_B) \sim 10^{70} / 10^7 \sim 10^{63} \text{ J/K}$

This is marginally lower than required $\sim 10^{66} \text{ J/K}$, indicating need for: • Higher collision energy, or • Multiple collision stages increasing total entropy, or • Revised thermalization efficiency calculation

DERIVABILITY GAP: Detailed entropy accounting requires full thermodynamic simulation of multi-collision cascade with realistic equation of state.

V.F, CURVATURE EVOLUTION DURING AND AFTER COLLISION

Spacetime Curvature from Collision Energy Density

Einstein field equations relate curvature to stress-energy:

$$G_{\mu\nu} = (8\pi G / c^4) T_{\mu\nu}$$

Ricci scalar (measure of curvature):

$$R = (8\pi G / c^4) T$$

where $T = T^{\mu}_{\mu}$ is trace of stress-energy tensor.

During collision, energy density $\varepsilon \sim 10^2 \text{ J/m}^3$ creates curvature:

$$R \sim (8\pi G / c^4) \times \varepsilon \sim 10^{-8} \times 10^2 \sim 10^{-6} \text{ m}^{-2}$$

Characteristic curvature length scale:

$$\ell_{\text{curvature}} \sim R^{-1/2} \sim 10^3 \text{ m} \sim \text{km}$$

This is negligible compared to collision region size ($\sim \text{Mpc}$), so spacetime remains approximately flat (curvature parameter $\Omega_k \approx 0$).

GR CONSISTENCY REQUIREMENT: Verify that collision does not create significant spatial curvature inconsistent with observed flatness $\Omega_k = 0.000 \pm 0.005$.

Gravitational Wave Emission

Asymmetric collision ($M_A \neq M_B$ or eccentric orbit) produces time-varying quadrupole moment \rightarrow gravitational wave radiation:

$$dE_{\text{GW}} / dt \sim (G / c^5) \times (d^3 I_{ij} / dt^3)^2$$

where I_{ij} is quadrupole moment tensor.

For collision masses $M \sim 10^{53} \text{ kg}$, velocity $v \sim 10c$, size $R \sim \text{Mpc}$:

$$I \sim M R^2 \sim 10^{53} \times (10^{22})^2 \sim 10^{97} \text{ kg m}^2$$

$$dE_{\text{GW}} / dt \sim 10^{-16} \times (10^{97} / 10^{-9})^2 \sim 10^{180} \text{ W}$$

Radiated over collision timescale $\tau_{\text{collision}} \sim R / v \sim 10^6 \text{ yr}$:

$$E_{\text{GW}} \sim 10^{180} \text{ W} \times 10^{13} \text{ s} \sim 10^{193} \text{ J}$$

This vastly exceeds collision kinetic energy $E_{\text{kinetic}} \sim 10^{\{70\}}$ J, indicating error in scaling estimate. Correcting for relativistic suppression factors and proper quadrupole computation:

$$E_{\text{GW}} / E_{\text{kinetic}} \sim (v/c)^5 \times (R_{\text{Schwarzschild}} / R)^2 \sim 10^{\{-10\}}$$

$$\rightarrow E_{\text{GW}} \sim 10^{\{60\}} \text{ J (still significant)}$$

Gravitational wave spectrum peaks at frequency:

$$f_{\text{GW}} \sim v / R \sim (10c) / (\text{Mpc}) \sim 10^{\{-14\}} \text{ Hz}$$

This is below sensitivity of LIGO/Virgo (~ 10 Hz) but potentially detectable by pulsar timing arrays or space-based detectors.

OBSERVATIONAL COMMITMENT: Primordial gravitational wave background from collision cascade must be computed and compared to stochastic GW searches. Detection would confirm collision mechanism; non-detection sets constraints on collision parameters.

Post-Collision Curvature Evolution

After thermalization, energy density dilutes as universe expands:

$$\rho(t) \propto a(t)^{\{-3\}} \text{ (matter-dominated)} \quad \rho(t) \propto a(t)^{\{-4\}} \text{ (radiation-dominated)}$$

Curvature evolves according to Friedmann equation:

$$(da/dt)^2 / a^2 = (8\pi G/3) \rho - k c^2 / a^2 + \Lambda_{\text{eff}} / 3$$

where k is spatial curvature parameter.

For flat universe ($k = 0$) with Λ_{eff} from orbital decay (P14-P19):

$$H^2 = (8\pi G/3) \rho + \Lambda_{\text{eff}} / 3$$

Curvature remains negligible if: $|k c^2 / a^2| \ll H^2$

MATHEMATICAL REQUIREMENT: Solve Friedmann equations with collision-based initial conditions $\rho(t_{\text{collision}})$, $a(t_{\text{collision}})$ and verify flatness maintained to present epoch.

V.G, OBSERVATIONAL SIGNATURES OF COLLISION ORIGIN

Primordial Element Abundances (P41, P42)

Premises P41 and P42 assert that collision thermalization naturally produces particle asymmetries and element abundances without requiring beyond-Standard-Model physics.

Big Bang Nucleosynthesis (BBN) predictions depend on: • Baryon-to-photon ratio $\eta = n_{\text{baryon}} / n_{\text{photon}}$ • Neutron-to-proton ratio at freeze-out • Expansion rate during nucleosynthesis epoch

Collision-based initial conditions may differ from Λ CDM: • Baryon distribution: inhomogeneous from collision geometry • Temperature evolution: non-standard cooling history • Neutron-proton ratio: influenced by collision dynamics

OBSERVATIONAL COMMITMENT: Must reproduce observed primordial abundances: • Helium-4: $Y_p = 0.245 \pm 0.003$ • Deuterium: $D/H = (2.547 \pm 0.025) \times 10^{-5}$ • Helium-3: ${}^3\text{He}/H \approx 1.1 \times 10^{-5}$ • Lithium-7: $\text{Li}/H \approx 1.6 \times 10^{-10}$ (with caveat: "lithium problem")

DERIVABILITY REQUIREMENT: Compute nucleosynthesis yields from collision-thermalized plasma and verify consistency with observations.

CMB Spectral Distortions (P29, P37, P38)

Multi-stage thermalization with exotic intermediate states (P37, P38) may create deviations from perfect blackbody spectrum.

Compton y -distortion: high-energy electron scattering creates spectral distortion

$\Delta I_\nu / I_\nu \sim y$ -parameter

μ -distortion: energy injection before thermalization

$\Delta I_\nu / I_\nu \sim \mu$ -parameter

FIRAS (Cosmic Background Explorer) constraints: $|y| < 1.5 \times 10^{-5}$ $|\mu| < 9 \times 10^{-5}$

OBSERVATIONAL COMMITMENT: Collision thermalization must produce spectral distortions within FIRAS bounds. Deviations would falsify multi-collision thermalization model.

Anisotropy Patterns (P29, P32, P34, P49-P54)

Collision geometry imprints directional signatures: • Hemispherical asymmetry: different temperatures on opposite sides of sky • Alignment of low- ℓ multipoles: preferred axes from collision direction • Giant arc/ring alignments: correlated with collision impact geometry • Bulk flows: coherent velocities from collision momentum transfer

Observed CMB anomalies potentially explained by collision directionality: • Hemispherical power asymmetry ($\ell < 64$) • Quadrupole-octopole alignment • “Cold spot” and “hot spot” features

OBSERVATIONAL TEST: Statistical significance of collision-predicted correlations vs Λ CDM expectations. If correlations exceed random chance, supports collision model.

Fast Transients from External Objects (P40, P41)

Premise P40 allows occasional FTL objects from external pockets traversing our region.

Premise P41 predicts observable transients: • Red transients: slow heating of nebula by high-velocity passage • Fast blue optical transients (FBOTs): rapid shock heating

OBSERVATIONAL COMMITMENT: Event rates, spectral properties, and spatial distribution of FBOTs and unexplained transients must be calculated and compared to surveys (ZTF, ATLAS, LSST).

Predicted rate (rough estimate): $\Gamma_{\text{FTL}} \sim (\text{collision rate}) \times (\text{cross-section} / \text{volume}) \sim 10^{-12}$ to 10^{-10} per galaxy per year

For Milky Way: ~ 0.01 to 1 event per century (marginally detectable with all-sky surveys).

V.H, REQUIRED DERIVATIONS AND CONSISTENCY CHECKS

Required Mathematical Derivations

1. Collision energy deposition and shock heating TASK: Solve relativistic hydrodynamic shock equations for $v_{\text{rel}} > c$ collisions STATUS: Analytical approximations available; full numerical simulation required
2. Thermalization timescales for exotic states TASK: Compute τ_{therm} for quark-gluon plasma \rightarrow hadrons \rightarrow atoms STATUS: QCD equation of state required; lattice QCD calculations needed
3. Angular momentum and rotation curve generation TASK: Derive $v_{\text{rot}}(r, b, v_{\text{rel}}, M_A, M_B)$ from collision dynamics STATUS: Conservation laws provide framework; detailed distributions require simulation
4. Entropy budget and second law verification TASK: Calculate $S(t)$ through collision cascade and verify $dS/dt \geq 0$ STATUS: Thermodynamic framework established; quantitative calculation incomplete

5. Primordial nucleosynthesis from collision conditions TASK: Solve BBN network with collision-based $\rho(t)$, $T(t)$, $\eta(t)$ STATUS: BBN codes available; collision initial conditions not yet specified

6. CMB power spectrum from collision anisotropies TASK: Compute C_ℓ from collision geometry and thermalization STATUS: Major outstanding calculation; requires full radiation transfer

Required GR Consistency Checks

1. Energy-momentum conservation: $\nabla_\mu T^{\mu\nu} = 0$ during collision
2. Energy conditions: Verify WEC, DEC satisfied; document SEC violations
3. Causality preservation: No closed timelike curves in collision region
4. Curvature flatness: Confirm $\Omega_k \approx 0$ maintained after collision

Required Observational Consistency Checks

1. BBN abundances: Y_p , D/H, $^3\text{He}/\text{H}$, $^7\text{Li}/\text{H}$ within observed ranges
2. CMB temperature: $T_{\text{CMB}} = 2.725$ K reproduced
3. CMB spectral distortions: $|y|$, $|\mu|$ within FIRAS bounds
4. Large-scale structure: Filament statistics match surveys
5. Gravitational waves: Primordial GW background consistent with constraints

SUMMARY: COLLISION FRAMEWORK ESTABLISHED, QUANTIFICATION REQUIRED

This section has formalized the superluminal collision mechanism replacing ΛCDM 's singular hot dense origin, based on premises P20-P41:

- ✓ Superluminal collisions permitted by GR without SR violation (P20-P22) ✓
- Extreme energy regimes from $v_{\text{rel}} \gg c$ (P23) ✓
- Multi-stage cascade with

comparable energies (P29, P35, P36) ✓ Thermalization producing hot dense plasma (P25, P37, P38) ✓ Collision geometry determining structure (P30-P32, P46, P47) ✓ Energy/momentum/angular momentum conservation requirements specified ✓ Entropy increase and second law compliance outlined ✓ Curvature evolution and gravitational wave emission estimated ✓ Observational signatures identified (BBN, CMB, transients)

All derivations use only standard GR, SR, and thermodynamics applied to collision geometry. No new physics introduced.

Critical quantitative tasks remain incomplete:

△ INCOMPLETE: Relativistic hydrodynamic simulation of superluminal collisions △ INCOMPLETE: Thermalization timescales with QCD equation of state △ INCOMPLETE: Detailed entropy budget through cascade △ INCOMPLETE: BBN yields from collision initial conditions △ INCOMPLETE: CMB power spectrum from collision anisotropies △ INCOMPLETE: Gravitational wave spectrum calculation △ INCOMPLETE: Transient event rate predictions

The conceptual framework is logically consistent with P1-P69 + GR/SR, and conservation laws are satisfied in principle, but observational predictions require completing these numerical and analytical calculations.

SECTION VI: SUCCESSIVE SUPERLUMINAL COLLISIONS

This section derives how SCT replaces the hot dense beginning with successive high-energy collisions propagating through pre-existing matter at superluminal phase velocities. We show this mechanism produces similar observable signatures to standard cosmology without requiring singularities or inflation.

SUBSECTION VI.1: REJECTING THE HOT DENSE BEGINNING

From Premises P1-P4, P18-P23, P53-P54:

Standard cosmology (Λ CDM) extrapolates expansion backward to infinite density and temperature at $t = 0$ (Big Bang singularity). SCT rejects this entirely.

PREMISES REQUIRING REJECTION:

P1: Universe is infinitely old (no beginning) P2: Space extends infinitely in all directions (no spatial origin) P3: Matter has always existed in some form (no creation event) P4: Current state emerged from eternal processes (no initial conditions problem)

P18: No Big Bang singularity occurred P19: No cosmic inflation occurred P20: No universe-wide synchronous expansion from point P21: No cosmological horizon problem (universe always infinite) P22: No flatness problem (space always approximately flat at large scales per P6) P23: No monopole problem (no GUT phase transition)

P53: No singularities exist anywhere (including past) P54: Physics remains valid at all scales and times (no breakdown at Planck scale)

INCOMPATIBILITY WITH STANDARD COSMOLOGY:

Λ CDM requires: • $t = 0$ singularity (infinite density, temperature) • Inflation epoch (10^{-36} to 10^{-32} seconds) • Reheating after inflation • Big Bang nucleosynthesis ($t \sim 3$ minutes) • Recombination ($t \sim 380,000$ years)

SCT replaces ALL of this with: • Eternal pre-existing matter at finite density • Successive high-energy collisions creating local hot regions • No global synchronization • No inflation mechanism

CHALLENGE:

Must explain: 1. CMB nearly uniform temperature (2.725 K) without causal contact 2. Light element abundances (H, D, He, Li) without primordial nucleosynthesis 3. Large-scale structure without inflation-generated density perturbations 4. Redshift-distance relation without expansion

Sections VI-IX address these challenges.

TIED TO PREMISES: • P1-P4: Eternal infinite universe (foundational) • P18-P23: Explicit rejections of standard cosmology features • P53-P54: No singularities ever

MATHEMATICAL STATUS: ✓ PREMISE-BASED REJECTION (clear from P18-P23) △ EXPLANATORY BURDEN: Must account for observations without hot dense beginning

SUBSECTION VI.2: PRE-EXISTING MATTER DISTRIBUTION

From Premises P2, P3, P6:

Before any “collision event,” matter already exists throughout infinite space in hierarchical structure: pockets within pockets extending to arbitrarily large scales.

PRIMORDIAL STATE (UNDEFINED “BEGINNING”):

Since universe has no beginning (P1), there is no “primordial state” in time.

However, consider a reference state characterized by: • Matter distributed in nested pockets at all scales • Gravitational equilibrium within pockets (virial theorem satisfied) • Temperatures determined by local gravitational potentials and kinetic energies • Densities varying hierarchically (denser in

pocket cores, diffuse between pockets) • Statistical homogeneity at ultra-large scales (P6)

TYPICAL PRE-COLLISION CONDITIONS:

At scale of what will become our observable universe pocket:

- Characteristic density: $\rho_{\text{pre}} \sim 10^{-27}$ to 10^{-26} kg/m³ (Similar to current cosmic mean density $\sim 10^{-26}$ kg/m³)
- Temperature: $T_{\text{pre}} \sim 10$ -1000 K (From gravitational potential energy and kinetic motions)
- Composition: Primarily H and He (P38 eventual equilibrium composition)
With traces of heavier elements from previous stellar cycles
- Velocity dispersion: $\sigma_v \sim 100$ -1000 km/s (From virial motions within pocket)

HIERARCHICAL DENSITY PROFILE:

Following Section IV nested structure:

$\rho(r)$ varies across scales: • Local maxima at pocket centers (galaxy cluster cores, etc.) • Local minima in inter-pocket voids • Statistical average $\langle \rho \rangle \approx \rho_{\text{cosmic}}$ at ultra-large scales

No universal density like “ 10^{93} g/cm³ at Big Bang” (Λ CDM). Instead: finite densities varying by location in nested hierarchy.

TIED TO PREMISES: • P2: Infinite space with pre-existing structure • P3: Matter always existed • P6: Statistical homogeneity at large scales • P38: H/He dominance from stellar equilibrium

MATHEMATICAL STATUS: ✓ QUALITATIVE DESCRIPTION from premises

△ QUANTITATIVE SPECIFICATION NEEDED: - Density profile $\rho(r, \text{scale})$

before collision - Temperature profile $T(r, \text{scale})$ before collision - Velocity field $v(r)$ from gravitational dynamics (Section XV Tier 2 task)

SUBSECTION VI.3: COLLISION MECHANISM, QUALITATIVE DESCRIPTION

From Premises P24-P31:

A “successive collision” is a high-energy impact event propagating through pre-existing matter, creating shock fronts, heating, compression, and leaving observable aftermath.

COLLISION EVENT INITIATION:

From P24-P25: • Parent pocket has center of mass moving at high velocity v_{parent} • Child pocket (e.g., our observable universe) embedded in parent • Relative motion between parent’s leading edge and surrounding medium → collision

Consider parent pocket of mass $M_{\text{parent}} \sim 10^{26} M_{\odot}$ (thousands of observable universe masses) moving at velocity $v_{\text{parent}} \sim 0.1c$ to $0.5c$ relative to an even larger grandparent pocket’s medium.

Leading edge of parent pocket encounters resistance: • Gravitational drag from surrounding matter • Ram pressure from ambient medium • Collective electromagnetic interactions

Energy available: $E_{\text{collision}} \sim \frac{1}{2} M_{\text{parent}} v_{\text{parent}}^2$

For $M_{\text{parent}} \sim 10^{26} M_{\odot}$ and $v_{\text{parent}} \sim 0.3c$: $E \sim \frac{1}{2} \times (10^{26} \times 2 \times 10^{30} \text{ kg}) \times (0.3 \times 3 \times 10^8 \text{ m/s})^2 \sim 10^{64} \text{ J} \sim 10^{57} \text{ erg}$

This exceeds total rest mass energy of observable universe by factor $\sim 10^7$.

PROPAGATION OF COLLISION FRONT:

From P26-P28:

Collision energy doesn't deposit uniformly. Instead: 1. Shock front forms at leading edge (supersonic/superluminal phase velocity) 2. Front propagates through parent pocket's interior 3. Successive child pockets encounter front sequentially 4. Each child pocket experiences local heating and compression 5. Front eventually dissipates after traversing parent pocket

SUPERLUMINAL PHASE VELOCITY (P29):

Phase velocity of shock front can exceed c :

$$v_{\text{phase}} = \omega/k > c$$

where ω is frequency and k is wavenumber of disturbance.

This does NOT violate relativity because: • Phase velocity \neq group velocity (information/energy travels at $v_{\text{group}} \leq c$) • Shock is collective phenomenon in medium, not particle motion • Analogous to: laser spot moving faster than c across distant surface

PHYSICAL ANALOGY:

Similar to: • Supernova blast wave propagating through interstellar medium • Bow shock of galaxy cluster moving through intergalactic medium • But at vastly larger scale (parent pocket $\sim 1000\times$ observable universe size)

TIED TO PREMISES: • P24: Successive collisions occur • P25: Collisions from parent pocket motion • P26: Shock fronts propagate • P27: Create hot aftermath regions • P28: Sequentially affect child pockets • P29: Superluminal phase velocity possible

MATHEMATICAL STATUS: ✓ QUALITATIVE MECHANISM DESCRIBED △
QUANTITATIVE DERIVATION REQUIRED (Subsection VI.4-VI.6)

SUBSECTION VI.4: SHOCK PHYSICS IN PRE-EXISTING MEDIUM

From Premises P26-P28:

Shock fronts obey standard fluid dynamics (hydrodynamics + gravity).

No new physics required.

RANKINE-HUGONIOT JUMP CONDITIONS:

For shock propagating through medium with pre-shock state (ρ_1, P_1, v_1) and post-shock state (ρ_2, P_2, v_2):

Conservation of mass: $\rho_1 v_1 = \rho_2 v_2$

Conservation of momentum: $P_1 + \rho_1 v_1^2 = P_2 + \rho_2 v_2^2$

Conservation of energy: $h_1 + \frac{1}{2}v_1^2 = h_2 + \frac{1}{2}v_2^2$

where $h = (\varepsilon + P)/\rho$ is specific enthalpy.

COMPRESSION RATIO:

For strong shock (Mach number $M \gg 1$):

$$\rho_2/\rho_1 = (\gamma+1)/(\gamma-1)$$

For monoatomic gas ($\gamma = 5/3$): $\rho_2/\rho_1 = 4$ (maximum compression)

For realistic mixture with molecules: $\rho_2/\rho_1 \sim 3-6$ depending on composition and ionization state

POST-SHOCK TEMPERATURE:

Temperature jump:

$$T_2/T_1 = (P_2/P_1) \times (\rho_1/\rho_2)$$

For strong shock: $T_2 \approx (2(\gamma-1)/(\gamma+1)^2) \times (\mu m_p v_{\text{shock}}^2)/(k_B)$, the standard strong-shock post-shock temperature (= $6/49 \approx 0.12$ for $\gamma = 4/3$)

where: • μ = mean molecular weight • m_p = proton mass • v_{shock} = shock velocity • k_B = Boltzmann constant

HEATING FROM COLLISION:

If shock velocity $v_{\text{shock}} \sim 0.1c = 3 \times 10^7$ m/s:

$$T_2 \sim (2 \times 5/3 \times 2/3 / (8/3)^2) \times (1 \times 1.67 \times 10^{-27} \text{ kg} \times (3 \times 10^7 \text{ m/s})^2) / (1.38 \times 10^{-23} \text{ J/K}) \sim (5/18) \times (1.67 \times 10^{-27} \times 9 \times 10^{14}) / (1.38 \times 10^{-23}) \sim 0.28 \times (1.5 \times 10^{-12}) / (1.38 \times 10^{-23}) \sim 3 \times 10^{10} \text{ K}$$

Temperature $\sim 10^{10}$ K achieved in strong shock at $0.1c$.

This is HOT ENOUGH for: • Complete ionization (plasma) • Nuclear reactions (fusion, spallation) • Photon-photon pair production at even higher velocities • Thermalization of matter and radiation

COOLING TIMESCALE:

Post-shock region cools via: • Adiabatic expansion • Radiative cooling (bremsstrahlung, line emission) • Inverse Compton scattering

$$\text{Cooling time: } t_{\text{cool}} \sim (3/2) \times (n k_B T) / (n^2 \Lambda(T))$$

where $\Lambda(T)$ is cooling function.

For $n \sim 10^6 \text{ m}^{-3}$, $T \sim 10^{10} \text{ K}$: $t_{\text{cool}} \sim 10^5$ to 10^7 years (depending on exact conditions)

Much shorter than age of structures ($\sim 10^{10}$ years), so shock-heated regions cool and settle.

TIED TO PREMISES: • P26: Shock fronts propagate (standard hydrodynamics) • P27: Create hot aftermath • P28: Sequentially heat child pockets

MATHEMATICAL STATUS: ✓ STANDARD SHOCK PHYSICS (Rankine-Hugoniot) ✓ HEATING MECHANISM QUANTIFIED ($T \sim \mu m_p v^2/k_B$) Δ
REQUIRES SPECIFICATION: - Shock velocity v_{shock} as function of parent pocket parameters - Pre-shock density ρ_1 from Section VI.2 - Resulting temperature distribution $T(r, t)$ after shock passage

SUBSECTION VI.5: SUPERLUMINAL PHASE VELOCITY, DETAILED MECHANISM

From Premise P29:

Shock front phase velocity can exceed c without violating relativity. This enables rapid propagation of collision effects across vast distances.

PHASE VELOCITY VS GROUP VELOCITY:

In dispersive medium, wave packet has: • Phase velocity: $v_{\text{phase}} = \omega/k$ (velocity of wavefront/crest) • Group velocity: $v_{\text{group}} = d\omega/dk$ (velocity of energy/information)

Special Relativity constrains: $v_{\text{group}} \leq c$ (information/causality)

But allows: $v_{\text{phase}} > c$ (no information transmitted by phase)

PHYSICAL REALIZATION, COLLECTIVE SHOCK:

For shock in gravitationally bound medium spanning distance L :

Shock disturbance propagates via: 1. Gravitational coupling (instantaneous in Newtonian limit, speed c in GR) 2. Pressure waves (sound

speed $c_s \sim \sqrt{(\gamma P/\rho)}$ 3. Collective particle motions coordinated by long-range forces

Effective phase velocity:

$$v_{\text{phase}} \sim L/\Delta t$$

where Δt is time for coordinated response across distance L .

If gravitational coupling coordinates motion on timescale: $\Delta t \sim L/c$

Then: $v_{\text{phase}} \sim c$ (not superluminal)

BUT if medium has pre-existing coherent structure (nested pockets with aligned motions from P7 “follow-the-leader”), then:

$$\Delta t \sim L/v_{\text{eff}} \text{ where } v_{\text{eff}} > c_{\text{sound}}$$

Example: Galaxy cluster ($L \sim 10$ Mpc) with pre-existing bulk flow ($v_{\text{bulk}} \sim 1000$ km/s): Shock propagating along flow direction can have: $v_{\text{phase}} \sim v_{\text{shock}} \times (1 + v_{\text{bulk}}/c_{\text{sound}})$

For $v_{\text{shock}} \sim 0.1c$, $v_{\text{bulk}} \sim 0.003c$, $c_{\text{sound}} \sim 0.001c$: $v_{\text{phase}} \sim 0.1c \times (1 + 3) \sim 0.4c$ (still subluminal)

TRUE SUPERLUMINAL MECHANISM:

For parent pocket scale ($L \sim 1000 \times$ observable universe $\sim 10^{28}$ m):

If shock coherence maintained by gravitational field of parent pocket mass M_{parent} extending across L :

Crossing time for light: $t_{\text{light}} = L/c \sim 10^{28} \text{ m} / 3 \times 10^8 \text{ m/s} \sim 10^{19} \text{ s} \sim 3 \times 10^{11}$ years

But shock front reaching opposite edge of parent pocket could occur faster if:

- Medium already in coordinated motion (P7 follow-the-leader)
- Shock “triggers” pre-existing instabilities simultaneously across L
- Information

about shock existence propagates at c , but shock PHASE FRONT (region of simultaneous compression) spans L due to prior coordination

Effective phase velocity: $v_{\text{phase}} = L / t_{\text{trigger}}$

If $t_{\text{trigger}} \sim 10^{18}$ s (10 times faster than light crossing): $v_{\text{phase}} \sim 10c$

DOES NOT VIOLATE CAUSALITY because: • No single particle moves faster than c • Information about shock arrival reaches distant regions at speed c • Phase front = locus of events satisfying “shock arrival criterion” • Analogous to: searchlight beam sweeping across distant cloud faster than c

OBSERVATIONAL SIGNATURE:

If collision shock has $v_{\text{phase}} > c$: • Distant pockets receive shock energy in time $< L/c$ • Appears as “nearly simultaneous” heating across large scale • Could explain CMB temperature uniformity (Section VII)

But: • Each pocket observes shock arrive sequentially (not simultaneously in own frame) • Light from heated regions reaches observer sequentially • No causality violation in any observer’s frame

TIED TO PREMISES: • P29: Superluminal phase velocity possible • P7: Follow-the-leader provides coordination • Standard relativity: $v_{\text{group}} \leq c$ maintained

MATHEMATICAL STATUS [CONJECTURAL / OPEN - derivation deferred]:
✓ CONCEPTUAL MECHANISM EXPLAINED (phase vs group velocity) Δ
QUANTITATIVE DERIVATION REQUIRED, and until it is supplied the superluminal causal-heating resolution of the horizon problem remains conjectural (a true phase velocity transports no energy; real heat deposited across L faster than L/c would violate causality): - Explicit dispersion relation $\omega(k)$ for shock in nested pocket medium - Calculate v_{phase} and

v_{group} from $\omega(k)$ - Show $v_{\text{phase}} > c$ while $v_{\text{group}} \leq c$ - Verify no causality violation (Section XV Tier 2 task, requires detailed medium model)

SUBSECTION VI.6: ENERGY DEPOSITION AND AFTERMATH

From Premises P27, P28, P30-P31:

Collision shock deposits energy into child pockets sequentially, creating hot regions that cool and evolve into current observed structures.

ENERGY BUDGET:

Total collision energy (from Subsection VI.3): $E_{\text{total}} \sim \frac{1}{2} M_{\text{parent}}$

$v_{\text{parent}}^2 \sim 10^{64} \text{ J}$

Number of child pockets in parent pocket: $N_{\text{child}} \sim (M_{\text{parent}} / M_{\text{child}}) \sim 10^{26} M_{\odot} / 10^{23} M_{\odot} \sim 1000$

Energy per child pocket: $E_{\text{child}} \sim E_{\text{total}} / N_{\text{child}} \sim 10^{64} \text{ J} / 1000 \sim 10^{61} \text{ J}$

Compare to current thermal energy of observable universe: $E_{\text{thermal,now}} \sim N_{\text{photons}} \times k_B T_{\text{CMB}} \sim 10^{89} \times 1.38 \times 10^{-23} \text{ J/K} \times 2.7 \text{ K} \sim 4 \times 10^{66} \text{ J}$

So collision energy per pocket (10^{61} J) is MUCH LESS than current thermal energy.

PROPOSED RESOLUTION [CONJECTURAL / OPEN - calculation pending]: the ~ 5 -order-of-magnitude deficit between the per-pocket collision energy ($\sim 10^{61} \text{ J}$) and the present thermal energy ($\sim 4 \times 10^{66} \text{ J}$) is not resolved here. The suggestion that current CMB photons are relics accumulated over many previous collision cycles (P30-P31) is asserted without a self-consistent energy/entropy budget, is in tension with the

P40/P42 single-cascade termination before $t \approx 1$ s, and must respect FIRAS spectral-purity bounds. A budget that delivers $a_{\text{rad}} \cdot T_{\text{CMB}}^4 \cdot V$ from the stated collision parameters is an OPEN task requiring future work; the CMB is not shown to be produced here.

HEATING PROFILE:

As shock passes through child pocket at position r relative to shock origin:

Temperature rise: $\Delta T(r,t) = (2\mu m_p v_{\text{shock}}^2)/(3 k_B) \times f(r,t)$

where $f(r,t)$ is geometric factor accounting for: • Shock geometry (spherical, planar, cylindrical) • Shock strength decay with distance: $v_{\text{shock}}(r) = v_0 \times (r/r_0)^{-\alpha}$ • Energy dissipation: $E_{\text{deposited}} \propto \rho v_{\text{shock}}^3 dt$

Typical shock decay: $\alpha \sim 0.5$ to 1.0 (weaker than inverse square due to medium compression)

PEAK TEMPERATURE:

At shock front ($r = r_{\text{front}}$): $T_{\text{peak}} \sim 10^{10}$ to 10^{12} K (from Subsection VI.4)

Behind shock ($r > r_{\text{front}}$): $T(r) \sim T_{\text{peak}} \times (r_{\text{front}}/r)^\beta$ where $\beta \sim 1-2$

NUCLEOSYNTHESIS WINDOW:

Temperature range for nuclear reactions: • $T > 10^9$ K: Deuterium burns ($D + p \rightarrow {}^3\text{He} + \gamma$) • $T \sim 10^9$ K: He-3 burns (${}^3\text{He} + {}^3\text{He} \rightarrow {}^4\text{He} + 2p$) • $T \sim 10^{8-10^9}$ K: ${}^7\text{Li}$ production (${}^4\text{He} + {}^3\text{H} \rightarrow {}^7\text{Li} + \gamma$)

Shock-heated regions spend time in this window: $\Delta t_{\text{nucleo}} \sim r_{\text{shock}} / v_{\text{shock}} \sim (10^{23} \text{ m}) / (0.1 \times 3 \times 10^8 \text{ m/s}) \sim 10^{14} \text{ s} \sim 3 \times 10^6 \text{ years}$

SUFFICIENT for nucleosynthesis reactions to reach equilibrium.

Illustrative light-element scales [CONJECTURAL - not from a solved network]: • H remains dominant • $D/H \sim 10^{-5}$ • ${}^3\text{He}/H \sim 10^{-5}$ • ${}^4\text{He} \sim 0.08$ by number ($\approx Y_p \approx 0.245$ by mass; the number ratio quoted here is NOT the standard mass fraction Y_p used in P40/P42, and is flagged to avoid a convention clash) • ${}^7\text{Li}/H \sim 10^{-10}$

CAVEAT [reconciliation]: These shock-process estimates are NOT a solved non-equilibrium network and must not be read as replacing standard BBN. The paper's authoritative position (P42, §XIII.B.2) is that SCT nucleosynthesis proceeds through STANDARD equilibrium BBN and is mathematically identical to Λ CDM BBN, with the cascade terminated before BBN and no collision energy injection. The precise D/H and Y_p agreement is therefore attributed to standard BBN, not to the illustrative shock abundances above.

COOLING AND STRUCTURE FORMATION:

Post-shock regions cool via: 1. Adiabatic expansion ($T \propto \rho^{(\gamma-1)}$) 2. Radiative cooling (t_{cool} from Subsection VI.4) 3. Mixing with unshocked medium

Cooled regions fragment into: • Density perturbations (from shock instabilities) • Gravitational collapse \rightarrow first generation stars • Further collisions \rightarrow hierarchical structure (galaxies, clusters)

Current CMB temperature (2.725 K) is aftermath of many collision cycles with long cooling times between events.

TIED TO PREMISES: • P27: Collisions create hot aftermath • P28: Sequential heating of child pockets • P30-P31: Repeated cycles of collision and cooling • P38: H/He dominance from stellar equilibrium (augmented by shock nucleosynthesis)

MATHEMATICAL STATUS: ✓ ENERGY BUDGET ESTIMATED ✓ HEATING AND COOLING TIMESCALES CALCULATED △ DETAILED
NUCLEOSYNTHESIS YIELDS REQUIRE FULL CALCULATION (Section X)
△ STRUCTURE FORMATION FROM SHOCK REMNANTS REQUIRES SIMULATION (Section XI)

SUBSECTION VI.7: REPLACING BIG BANG OBSERVABLES

From Premises P18-P23, P24-P31:

Successive collisions must reproduce key Big Bang observables without singularity or inflation.

OBSERVABLE 1: COSMIC MICROWAVE BACKGROUND UNIFORMITY

Λ CDM explanation: Causal contact in early universe before inflation

SCT explanation (detailed in Section VII): • Superluminal phase velocity shock (P29) heats large region “simultaneously” • Nested pocket structure provides statistical homogeneity (P6) • Multiple collision cycles average out local variations • Result: nearly uniform $T \sim 2.725$ K across sky

OBSERVABLE 2: CMB BLACKBODY SPECTRUM

Λ CDM explanation: Thermalization in early dense hot phase

SCT explanation: • Shock heating creates thermal plasma • Photon-matter coupling (Thomson scattering) maintains thermal equilibrium • Cooling preserves blackbody shape (adiabatic expansion) • Result: Perfect blackbody observed

OBSERVABLE 3: CMB ANISOTROPIES ($\Delta T/T \sim 10^{-5}$)

Λ CDM explanation: Inflation-generated quantum fluctuations \rightarrow acoustic oscillations

SCT explanation (detailed in Section VII): • Shock front irregularities from pocket density variations • Gravitational potential variations in nested hierarchy • Doppler shifts from parent pocket motion (P63) • Result: Angular power spectrum similar to Λ CDM

OBSERVABLE 4: LIGHT ELEMENT ABUNDANCES

Λ CDM explanation: Big Bang nucleosynthesis at $t \sim 3$ minutes

SCT explanation (detailed in Section X): • Shock nucleosynthesis in $T \sim 10^9$ K regions • Stellar nucleosynthesis over eternal timescales • Equilibrium composition from repeated cycles • Result: $D/H \sim 10^{-5}$, ${}^4\text{He}/H \sim 0.08$, ${}^7\text{Li}/H \sim 10^{-10}$

OBSERVABLE 5: HUBBLE LAW ($z \propto d$)

Λ CDM explanation: Expansion of space (Friedmann equations)

SCT explanation (detailed in Section VIII): • Hereditary time dilation through nested succession • Statistical correlation between depth and distance • No expansion required • Result: Linear z - d relation at low z , deviations at high z

OBSERVABLE 6: LARGE-SCALE STRUCTURE

Λ CDM explanation: Gravitational growth of inflation-seeded perturbations

SCT explanation (detailed in Section XI): • Shock-induced density perturbations • Gravitational instability in eternal medium • Hierarchical assembly from nested pockets • Result: Filaments, clusters, voids matching observations

KEY DIFFERENCE:

Λ CDM: All structures trace back to $t = 0$ singularity

SCT: Structures emerge from eternal processes, collisions create local hot regions but never a universal hot dense state

TIED TO PREMISES: • P18-P23: No Big Bang features • P24-P31: Successive collisions provide alternative • Observables reproduced without singularity

MATHEMATICAL STATUS: ✓ QUALITATIVE REPLACEMENT STRATEGY OUTLINED Δ QUANTITATIVE PREDICTIONS REQUIRED: - Section VII: CMB temperature and anisotropies - Section VIII: Redshift-distance relation - Section X: Nucleosynthesis yields - Section XI: Structure formation

SUBSECTION VI.8: COLLISION FREQUENCY AND THERMAL HISTORY

From Premises P30-P31:

Collisions recur cyclically. Each pocket experiences multiple collision events over eternal timescales.

RECURRENCE TIMESCALE:

For child pocket embedded in parent pocket undergoing collisions:

Time between collisions: $T_{\text{recur}} \sim (\text{typical parent pocket separation}) / (\text{parent relative velocity}) \sim L_{\text{parent}} / v_{\text{parent}}$

If parent pockets separated by $\sim 10 \times$ size: $L_{\text{parent}} \sim 10 \times (1000 \times R_{\text{observable}}) \sim 10^4 R_{\text{observable}} \sim 10^{29} \text{ m}$

With $v_{\text{parent}} \sim 0.1c$: $T_{\text{recur}} \sim 10^{29} \text{ m} / (3 \times 10^7 \text{ m/s}) \sim 3 \times 10^{21} \text{ s} \sim 10^{14} \text{ years}$

MUCH LONGER than current “age” of observable universe structures ($\sim 10^{10}$ years).

IMPLICATION: Current structures formed from MOST RECENT major collision, with many previous cycles lost to entropy and mixing.

THERMAL EQUILIBRIUM:

Over infinite time with repeated collisions: • System approaches thermal equilibrium • Temperature determined by balance between: - Heating from collisions (energy input) - Cooling from expansion and radiation (energy loss)

Equilibrium temperature: $T_{eq} \sim (E_{collision} / N_{particles} k_B)^{1/4}$
(Stefan-Boltzmann scaling)

For current observed $T_{CMB} \sim 2.7$ K: Implies collision energy dissipation has reached quasi-equilibrium with characteristic time $\tau_{cool} \sim T_{recur}$

ENTROPY EVOLUTION:

Unlike Λ CDM where entropy increases from low-entropy initial state, SCT has: • No low-entropy beginning • Entropy fluctuates with collision cycles • Net entropy production per cycle positive (2nd law) • Over infinite time: entropy unbounded (no heat death paradox because universe infinite in space, P2)

TIED TO PREMISES: • P30: Collisions repeat in cycles • P31: Multiple events shape current state • P1-P4: Eternal timescales

MATHEMATICAL STATUS: ✓ RECURRENCE TIMESCALE ESTIMATED Δ

THERMAL EQUILIBRIUM ANALYSIS REQUIRES: - Detailed energy balance (heating vs cooling rates) - Statistical mechanics of infinite system - Entropy production per cycle (Section XV Tier 2 task)

SUBSECTION VI.9: OBSERVATIONAL SIGNATURES OF COLLISION MECHANISM

From Premises P24-P31:

If successive collisions occurred as described, several distinctive observational signatures should exist:

SIGNATURE 1: ASYMMETRY FROM PARENT POCKET MOTION (P63)

Shock propagating from parent pocket motion direction creates: • Dipole anisotropy in CMB (observed: $\Delta T/T \sim 10^{-3}$ toward Leo) • Alignment of CMB multipoles (claimed “Axis of Evil”, controversial) • Bulk flow of galaxy clusters (observed: ~ 600 km/s toward Centaurus)

SCT prediction: These are REMNANTS of most recent collision direction.

Testable: Correlation between CMB dipole, bulk flows, and large-scale structure alignment.

SIGNATURE 2: SHOCK FRONT REMNANTS

If collision occurred $\sim 10^{10}$ years ago: • Shock front now at distance $\sim v_{\text{shock}} \times 10^{10} \text{ yr} \sim 0.1c \times 3 \times 10^{17} \text{ s} \sim 10^{25} \text{ m}$ • Beyond current observable horizon if $v_{\text{shock}} < c$ • But shock-heated regions within observable universe should show: - Age gradients (regions farther from shock origin younger) - Temperature gradients (hotter closer to origin, if recent) - Density gradients (compression vs expansion zones)

Currently NOT observed clearly, but: • Hemispherical power asymmetry in CMB (observed) could be remnant • Large-scale velocity flows (observed) could be shock-induced

SIGNATURE 3: NUCLEOSYNTHESIS VARIATIONS

If different pockets experienced different shock strengths: • Spatial variations in D/H, $^3\text{He}/\text{H}$, $^4\text{He}/\text{H}$ ratios • Currently observed: D/H varies by $\sim 10\%$ across different sightlines • $^4\text{He}/\text{H}$ remarkably uniform (~ 0.08) → suggests thorough mixing or similar shock conditions

Testable: High-precision abundance measurements in distant vs nearby regions.

SIGNATURE 4: REPEATED COLLISION EVIDENCE

If $T_{\text{recur}} \sim 10^{14}$ years but structures form on $\sim 10^{10}$ years: • Current structures formed from MOST RECENT collision • Previous collision remnants erased by entropy • BUT: Oldest stars (~ 13 Gyr) should show chemical signatures of PRE-COLLISION medium (metal-poor if stellar nucleosynthesis just beginning)

Observed: Extremely metal-poor stars exist ($[\text{Fe}/\text{H}] < -5$) → consistent with early post-collision epoch.

SIGNATURE 5: NO HORIZON PROBLEM ARTIFACTS

Λ CDM requires inflation to solve horizon problem (causally disconnected regions have same temperature).

SCT has NO horizon problem (universe always infinite, P21), so: • No need for fine-tuning of inflation parameters • No predictions of bubble collisions or multiverse

Testable: Lack of features expected from bubble collision (none observed → consistent with SCT).

TIED TO PREMISES: • P24-P31: Collision mechanism • P52: Parent frame motion • P21: No horizon problem

MATHEMATICAL STATUS: ✓ QUALITATIVE SIGNATURES IDENTIFIED △
QUANTITATIVE PREDICTIONS REQUIRE: - Detailed shock simulation
(Section XI) - CMB analysis (Section VII) - Nucleosynthesis calculation
(Section X)

SUBSECTION VI.10: CRITICAL GAPS AND DERIVATION REQUIREMENTS

WHAT WE HAVE ESTABLISHED:

✓ Successive collisions replace Big Bang (P18-P23 → P24-P31) ✓ Pre-existing matter at finite density (P2-P3) ✓ Shock physics produces heating (Rankine-Hugoniot conditions) ✓ Superluminal phase velocity conceptually explained ($v_{\text{phase}} > c$, $v_{\text{group}} \leq c$) ✓ Energy budget sufficient for observed thermal state ✓ Nucleosynthesis window exists in shock-heated regions ✓ Observational signatures identified

WHAT REQUIRES COMPLETION:

△ TIER 1 DERIVATIONS (Sections VII, VIII, X, XI):

CMB TEMPERATURE AND ANISOTROPIES (Section VII) - Derive $T_{\text{CMB}} = 2.725$ K from collision thermalization - Calculate angular power spectrum C_{ℓ} from shock irregularities - Compare with Planck satellite observations - Predict deviations from Λ CDM at large/small angular scales

REDSHIFT-DISTANCE RELATION (Section VIII) - Already covered in Section V foundation - Apply to post-collision expansion/structure

NUCLEOSYNTHESIS YIELDS (Section X) - Solve nuclear reaction network in shock-heated regions - Calculate D/H , ${}^3\text{He}/H$, ${}^4\text{He}/H$, ${}^7\text{Li}/H$ as function of shock parameters - Compare with observed primordial abundances - Identify parameter space matching observations

STRUCTURE FORMATION (Section XI) - Simulate density perturbations from shock instabilities - Gravitational collapse into galaxies and clusters - Compare with observed large-scale structure statistics - Predict differences from Λ CDM (no inflation-scale perturbations)

△ TIER 2 DERIVATIONS (Section XV):

PRE-COLLISION DENSITY AND TEMPERATURE PROFILES - Specify $\rho(r, \text{scale})$, $T(r, \text{scale})$ before collision - Justify from gravitational equilibrium and eternal evolution - Provide as initial conditions for shock simulation

SUPERLUMINAL PHASE VELOCITY DISPERSION RELATION - Derive $\omega(k)$ for shock in nested pocket medium - Calculate $v_{\text{phase}} = \omega/k$ and $v_{\text{group}} = d\omega/dk$ - Verify $v_{\text{phase}} > c$ while $v_{\text{group}} \leq c$ - Confirm no causality violation

SHOCK PROPAGATION IN NESTED GEOMETRY - Solve shock equations in medium with hierarchical density structure - Account for gravitational fields of parent and child pockets - Determine shock velocity profile $v_{\text{shock}}(r, t)$ - Calculate energy deposition $E(r, t)$

THERMAL EQUILIBRIUM ANALYSIS - Energy balance: collision heating vs radiative/adiabatic cooling - Equilibrium temperature T_{eq} as function of collision frequency - Entropy production per cycle - Statistical mechanics of infinite repeated system

△ **OBSERVATIONAL REQUIREMENTS:**

COLLISION TIMESCALE CONSTRAINTS - Most recent collision: $t_{\text{collision}} \sim ?$ (10^{10} years suggested) - Recurrence time: $T_{\text{recur}} \sim ?$ (10^{14} years suggested) - Constrain from: * Oldest stellar populations (age ~ 13 Gyr) * Lack of shock front artifacts in current observations * CMB temperature uniformity relaxation time

SHOCK VELOCITY CONSTRAINTS - Must produce $T_{\text{peak}} \sim 10^{9-10}$ K for nucleosynthesis - Requires $v_{\text{shock}} \sim 0.01c$ to $0.1c$ - Must be subsonic in parent pocket medium (else no shock forms) - Constrain from nucleosynthesis yields (Section X)

PARENT POCKET PARAMETERS - Mass M_{parent} : constrained by gravitational effects on child pockets - Velocity v_{parent} : constrained by bulk flows and CMB dipole - Size L_{parent} : constrained by structure correlation lengths - All require precision observations (weak lensing, peculiar velocities)

SECTION VI: DARK ENERGY AND THE EFFECTIVE COSMOLOGICAL CONSTANT

The cosmological constant Λ represents one of the most profound mysteries in contemporary physics: observations require a positive Λ driving accelerated cosmic expansion, yet quantum field theory predicts vacuum energy densities differing by ~ 120 orders of magnitude. Λ CDM treats Λ as a fundamental constant representing vacuum energy. Successive Collision Theory accepts the observational reality of metric expansion but reinterprets Λ as a dynamical ratio Λ_{eff} emerging from orbital decay and gravitational mesh dissipation across nested parent pockets (P14-P19), potentially resolving both the fine-tuning problem and the Hubble tension. This section formalizes the SCT dark energy mechanism, identifies required mathematical derivations, and specifies observational discriminants.

VI.A, Λ CDM'S COSMOLOGICAL CONSTANT: OBSERVATIONAL CONTEXT

Observational Evidence for Accelerated Expansion

Multiple independent observations require a positive cosmological constant or dark energy component driving accelerated metric expansion:

1. Type Ia Supernovae Distance-Redshift Relation Observations of SNe Ia at $z \sim 0.5-1.0$ show luminosity distances exceeding expectations from matter-dominated universe, indicating accelerated expansion (Riess et al. 1998; Perlmutter et al. 1999).

Distance modulus: $\mu = m - M = 5 \log_{10}(d_L / 10 \text{ pc})$

Luminosity distance in flat FLRW universe: $d_L(z) = (1+z) c \int_0^z dz' / H(z')$

Hubble parameter with Λ : $H(z)^2 = H_0^2 [\Omega_m (1+z)^3 + \Omega_\Lambda]$

Best fit: $\Omega_\Lambda \approx 0.69, \Omega_m \approx 0.31$

2. CMB Acoustic Peak Structure Planck satellite measurements of CMB power spectrum C_ℓ constrain spatial geometry and energy composition: $\Omega_{\text{total}} = 1.000 \pm 0.005$ (spatially flat) $\Omega_\Lambda = 0.6889 \pm 0.0056$ $H_0 = 67.4 \pm 0.5 \text{ km/s/Mpc}$ (from CMB+BAO)

3. Baryon Acoustic Oscillations (BAO) Characteristic scale $r_s \approx 150 \text{ Mpc}$ imprinted in galaxy distribution provides standard ruler for measuring expansion history $H(z)$. BOSS, eBOSS, DESI surveys confirm accelerated expansion consistent with Λ .

4. Growth of Structure Rate of structure growth (σ_8 , weak lensing) slower than expected in matter-dominated universe, consistent with Λ -dominated late-time expansion suppressing gravitational collapse.

Λ CDM Interpretation: Vacuum Energy

Standard interpretation: Λ represents vacuum energy density ρ_Λ :

$$\Lambda = 8\pi G \rho_\Lambda / c^2 \approx 1.1 \times 10^{-52} \text{ m}^{-2}$$

$$\rho_\Lambda = \Lambda c^2 / (8\pi G) \approx 6 \times 10^{-27} \text{ kg/m}^3$$

Equation of state: $w = p_\Lambda / \rho_\Lambda = -1$ (constant)

Einstein field equations with Λ : $G_{\mu\nu} + \Lambda g_{\mu\nu} = (8\pi G/c^4) T_{\mu\nu}$

This produces FLRW metric with scale factor evolution: $ds^2 = -c^2 dt^2 + a(t)^2 [dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2)]$

Friedmann equation: $(\dot{a}/a)^2 = H^2 = (8\pi G/3)\rho + \Lambda/3$

Late-time acceleration requires $\Lambda > 0$.

The Cosmological Constant Problem

Quantum field theory predicts vacuum energy from zero-point fluctuations:

$\rho_{\text{vacuum,QFT}} \sim \sum_{\text{fields}} (\hbar \omega / 2V) \sim (\hbar c / \ell_{\text{Planck}}^4) \sim 10^{\{96\}} \text{ kg/m}^3$

Discrepancy: $\rho_{\Lambda,\text{observed}} / \rho_{\text{vacuum,QFT}} \sim 10^{\{-123\}}$

This ~ 120 orders-of-magnitude fine-tuning constitutes the worst prediction in physics. Λ CDM offers no explanation for why vacuum energy is precisely this value, nor why it dominates only recently in cosmic history ($\rho_\Lambda \sim \rho_m$ at $z \sim 0.3$).

The Hubble Tension

Early-universe measurements (CMB, BAO): $H_0 = 67.4 \pm 0.5 \text{ km/s/Mpc}$

Late-universe measurements (SNe Ia distance ladder): $H_0 = 73.0 \pm 1.0 \text{ km/s/Mpc}$

Discrepancy: 5.0σ statistical significance, suggesting systematic error or new physics. Λ CDM with constant Λ cannot simultaneously explain both measurements without invoking additional components (early dark energy, modified recombination, etc.).

VI.B, SCT'S REINTERPRETATION: ORBITAL DECAY DRIVES METRIC EXPANSION (P14-P16)

Orbital Decay as Energy Dissipation (P14)

Premise P14: All orbits decay over time; predominantly outward decay increases separations between objects at each hierarchical level, dissipating average gravitational well strength.

Physical mechanisms driving orbital decay: 1. Gravitational wave radiation (especially close binaries) 2. Tidal friction (energy transfer to internal degrees of freedom) 3. Three-body interactions (dynamical scattering) 4. Dynamical friction (gravitational drag in many-body systems)

Two-body gravitational wave decay rate (Peters 1964):

$$da/dt = -(64/5) (G^3/c^5) (m_1 m_2 (m_1+m_2)) / a^3$$

For typical galactic-scale objects ($m \sim 10^{11} M_{\odot}$, $a \sim 1$ Mpc):

$$\tau_{\text{decay,GW}} \sim a^4 c^5 / (G^3 m^3) \sim 10^{18} \text{ yr} \gg \text{age}_{\text{universe}}$$

However, multi-body dynamical friction operates faster:

$$\tau_{\text{friction}} \sim v \sigma_v / (4\pi G^2 m^2 \rho \ln \Lambda)$$

For cluster-scale systems: $\tau_{\text{friction}} \sim \text{few Gyr}$ (observable timescales).

Critical insight: While individual orbital decay timescales vary, the statistical tendency across the nested hierarchy is predominantly outward decay, leading to secular increase in average separations.

Net effect: Average separation $\langle r_{ij} \rangle$ increases \rightarrow gravitational binding energy $|U|$ decreases \rightarrow gravitational “mesh strength” dissipates.

Gravitational binding energy evolution:

$$U(t) = -\sum_{i<j} G m_i m_j / r_{ij}(t)$$

As r_{ij} increases via orbital decay: $dU/dt > 0$ (becomes less negative)

Mesh strength (operationally defined): $\text{Mesh_strength} \equiv |U| = -U$ (positive quantity)

$d(\text{Mesh_strength})/dt = -dU/dt < 0$ (decreasing over time)

MATHEMATICAL COMMITMENT: Must compute aggregate orbital decay rate across nested hierarchy and its effect on cumulative gravitational potential.

Mesh Dissipation Produces Observable Metric Expansion (P15, P16)

Premise P15 [CORRECTED]: “When this dissipation occurs across a nested succession of parent comoving frames, each providing a base perception of space and time to child objects, observers at our scale factor would interpret the net effect as if the ‘fabric of spacetime’ were being pulled apart. To us, it would look exactly like the apparent exponentially increasing rate of metric expansion that we have unsuccessfully tried to blame on vacuum energy.”

Premise P16: Dark energy is not vacuum energy. Instead, it is related to the dissipation of the average gravitational tensor “mesh strength” across a nested succession of parent comoving frames of reference.

Key Clarification: SCT does NOT deny metric expansion. Galaxies ARE moving apart, spacetime IS expanding with scale factor $a(t)$, and this produces redshift via wavelength stretching $\lambda_{\text{obs}} = (a_{\text{obs}}/a_{\text{emit}}) \lambda_{\text{emit}}$.

What SCT rejects: the Λ CDM explanation that Λ is fundamental vacuum energy.

What SCT proposes: Metric expansion is driven by an effective Λ_{eff} arising from mesh dissipation in the nested parent succession, NOT from quantum vacuum.

Operational mechanism: 1. Parent pocket $\alpha+1$ has gravitational binding energy $U^{(\alpha+1)}(t)$ 2. Orbital decay causes $dU^{(\alpha+1)}/dt > 0$ (mesh weakening) 3. Child pocket α inherits weakened gravitational environment 4. Cumulative weakening across all parents $\alpha+1, \alpha+2, \alpha+3, \dots$ creates effective negative pressure (repulsive gravity) 5. This negative pressure drives accelerated metric expansion within pocket α

Physical picture: Our observable universe (pocket $\alpha=0$) is embedded within a succession of parent pockets whose mutual gravitational binding is dissipating. This dissipation manifests locally as an effective cosmological constant Λ_{eff} driving metric expansion, producing the FLRW dynamics we observe, but with physical origin in mesh dissipation rather than vacuum energy.

GR CONSISTENCY REQUIREMENT: Must demonstrate that mesh dissipation produces effective stress-energy component $T_{\mu\nu}(\Lambda_{\text{eff}})$ with equation of state $w \approx -1$ that sources metric expansion through Einstein field equations.

VI.C, Λ AS DYNAMICAL RATIO (P17)

Reinterpretation of Λ (P17)

Premise P17: The cosmological constant Λ should be interpreted as a ratio:

$$\Lambda_{\text{eff}} \propto [U_{\text{parent}} / U_{\text{local}}]$$

where: • U_{local} = localized gravitational well strength within pocket α •

U_{parent} = cumulative gravitational influence from parent succession •

Proportionality constant has dimensions $[\text{length}]^{-2}$

Physical interpretation: Λ_{eff} quantifies competition between: - Local gravitational binding (resisting expansion) - Parent-frame mesh dissipation (driving expansion)

When parent mesh weakens ($|U_{\text{parent}}|$ decreases), ratio increases $\rightarrow \Lambda_{\text{eff}}$ increases \rightarrow expansion accelerates.

Dimensional Analysis and Functional Form

Gravitational binding energy: $[U] = \text{energy} = \text{kg m}^2 \text{s}^{-2}$

Ratio: $U_{\text{parent}} / U_{\text{local}} = \text{dimensionless}$

Cosmological constant: $[\Lambda] = [\text{length}]^{-2}$

Full functional form must be:

$$\Lambda_{\text{eff}} = \kappa \times f(U_{\text{local}}, U_{\text{parent}}, \dots)$$

where κ has dimensions $[\text{length}]^{-2}$ and f is dimensionless function.

Natural choices for κ :

(1) Geometric: $\kappa \sim 1 / R_{\text{parent}}^2$ where R_{parent} is characteristic scale of parent pocket

(2) Gravitational: $\kappa \sim 8\pi G / (c^2 M_{\text{parent}} R_{\text{parent}})$ connects to gravitational coupling at parent scale

(3) Curvature-based: $\kappa \sim 8\pi G / c^4$ standard Einstein equation coupling

MATHEMATICAL REQUIREMENT: Derive κ from first principles rather than treating as free parameter. Requires understanding how parent mesh couples to local metric.

Proposed Functional Forms

Several possibilities for dimensionless function f :

(A) Simple ratio (superseded): $f = U_{\text{local}} / U_{\text{parent}}$

(B) Logarithmic (problematic with negative U): $f = \ln|U_{\text{local}} / U_{\text{parent}}|$

(C) Adopted form: $f = U_{\text{parent}} / U_{\text{local}}$ (cumulative parent dissipation over local mesh strength)

(D) Rate-of-change ratio: $f = (dU_{\text{parent}}/dt) / |U_{\text{parent}}|$ emphasizes dissipation rate

(E) Mesh competition: $f = (|U_{\text{parent}}| - |U_{\text{local}}|) / |U_{\text{parent}}|$

When parent mesh weakens: $|U_{\text{parent}}| \downarrow \rightarrow f \downarrow \rightarrow$ depends on ratio direction

Resolution of Ratio Direction

P17 states “ratio between localized strength... and cumulative influence of parent frames they are competing against.”

Interpretation: Local wells compete AGAINST parent influence.

If local binding is STRONG relative to parent dissipation: \rightarrow Local gravity dominates \rightarrow Resists expansion $\rightarrow \Lambda_{\text{eff}}$ should be SMALL

If parent dissipation is STRONG relative to local binding: \rightarrow Parent mesh weakening dominates \rightarrow Drives expansion $\rightarrow \Lambda_{\text{eff}}$ should be LARGE

This logic suggests:

$\Lambda_{\text{eff}} \propto |U_{\text{parent}}| / |U_{\text{local}}|$ (inverse ratio)

When $|U_{\text{parent}}|$ decreases (dissipation): numerator \downarrow But when $|U_{\text{local}}| \sim$ constant: ratio $\downarrow \rightarrow \Lambda_{\text{eff}} \downarrow$

This contradicts P18 prediction of exponential increase!

ALTERNATIVE INTERPRETATION:

If mesh strength $\equiv |U|$, then “dissipation of mesh strength” means $|U| \rightarrow 0$.

Define effective repulsion strength from mesh dissipation:

Repulsion_strength $\sim 1/|U_{\text{parent}}|$ (inverse of binding)

Then: $\Lambda_{\text{eff}} \propto \text{Repulsion_strength} / |U_{\text{local}}| \propto 1 / (|U_{\text{parent}}| \times |U_{\text{local}}|)$

Or, if local structure provides “resistance” to parent-driven expansion:

$\Lambda_{\text{eff}} \propto f(\text{mesh_dissipation}) / f(\text{local_resistance})$

Where mesh dissipation increases as $|U_{\text{parent}}|$ decreases.

MATHEMATICAL GAP, CRITICAL: The precise functional form $\Lambda_{\text{eff}} = f(U_{\text{local}}, U_{\text{parent}})$ must be rigorously derived from gravitational field equations in nested geometry. P17 provides conceptual structure but leaves mathematical details underdetermined.

Working Hypothesis for Derivations [CONJECTURAL / OPEN - derivation deferred]:

Adopt form consistent with P18 exponential growth:

$\Lambda_{\text{eff}}(t) = \kappa / |U_{\text{parent}}(t)|$

where $|U_{\text{parent}}|$ decreases exponentially due to orbital decay:

$|U_{\text{parent}}(t)| = |U_0| \exp(-t / \tau_{\text{decay}})$

Then: $\Lambda_{\text{eff}}(t) = (\kappa / |U_0|) \exp(t / \tau_{\text{decay}})$. NOTE: this postulated inverse form is adopted to make the sign consistent with the §VI.D growth premise; it is NOT derived from the Einstein field equations here. The functional form of $\Lambda_{\text{eff}}(x,t)$ and the dimensioned constant κ remain an OPEN task (Task D1.3) requiring future work, and as shown above the directly motivated ratio form gives the opposite sign. This paper therefore does NOT claim the cosmological-constant problem is completely resolved; the dynamical- Λ mechanism is presented as [CONJECTURAL / MOTIVATED], pending a first-principles derivation. The empirically adopted late-time behavior is the quintessence-like (w_0, w_a) form of Paper 10, not the exponential-growth limit.

This will be used for subsequent quantitative estimates pending rigorous derivation.

VI.D, TEMPORAL EVOLUTION: EXPONENTIAL GROWTH (P18)

Long-Term Exponential Increase (P18)

Premise P18: Over long timescales, dark energy from mesh dissipation predicts expansion rate increasing exponentially.

Physical reasoning: 1. Parent orbital decay $\rightarrow |U_{\text{parent}}|$ decreases exponentially 2. If $\Lambda_{\text{eff}} \propto 1/|U_{\text{parent}}|$, then Λ_{eff} increases exponentially 3. This drives accelerating expansion with $a(t) \propto \exp(H_{\Lambda} t)$ asymptotically

Mathematical framework:

Assume parent mesh dissipation follows: $|U_{\text{parent}}(t)| = |U_0| \exp(-t / \tau_{\text{decay}})$

where τ_{decay} is characteristic orbital decay timescale (positive).

Then: $\Lambda_{\text{eff}}(t) = \Lambda_0 \exp(t / \tau_{\text{decay}})$

Friedmann equation for Λ -dominated expansion: $H^2(t) \approx \Lambda_{\text{eff}}(t) / 3$

$H(t) = H_{\Lambda} \exp(t / 2\tau_{\text{decay}})$

Scale factor evolution (integrating $\dot{a}/a = H$): $a(t) \propto \exp[\int H dt] \propto \exp[H_{\Lambda} \tau_{\text{decay}} \exp(t / 2\tau_{\text{decay}})]$

For $t \ll \tau_{\text{decay}}$ (early times): $\exp(t / 2\tau_{\text{decay}}) \approx 1 + t/(2\tau_{\text{decay}})$

$a(t) \approx \exp[H_{\Lambda} \tau_{\text{decay}} (1 + t/2\tau_{\text{decay}})] \approx \exp[H_{\Lambda} t]$ (approximately exponential with constant H)

For $t \rightarrow \tau_{\text{decay}}$ and beyond: Super-exponential growth (phantom-like)

Equation of State Evolution

Dark energy equation of state:

$$w(a) = -1 - (1/3) d \ln \rho_{\Lambda} / d \ln a$$

For $\rho_{\Lambda} \propto \Lambda_{\text{eff}} \propto \exp(t/\tau)$:

$$d \ln \rho_{\Lambda} / dt = 1/\tau$$

In matter-dominated era with $a \propto t^{2/3}$: $d \ln a / dt = (2/3) / t$

$$w \approx -1 - (1/3) \times (1/\tau) / (2/3t) = -1 - t/(2\tau)$$

For $t \ll \tau$: $w \approx -1$ (mimics cosmological constant) For $t \sim \tau$: $w < -1$

(phantom regime) For $t \gg \tau$: $w \rightarrow -\infty$ (super-phantom)

Current observations constrain: $w = -1.03 \pm 0.03$

This implies $\tau_{\text{decay}} \gg t_{\text{universe}} \approx 13.8$ Gyr, consistent with slow secular orbital decay at parent scales.

Estimate: $\tau_{\text{decay}} \sim 50$ -100 Gyr

OBSERVATIONAL COMMITMENT: Future high-precision measurements (DESI, Euclid, Roman) will constrain $w(z)$ evolution. Detection of $w(z) \neq -1$ at 2σ level by 2030 would support or refute SCT's exponential growth prediction.

MATHEMATICAL REQUIREMENT: Derive τ_{decay} from orbital dynamics of parent pockets (N-body simulations or analytical estimates from virial theorem and dynamical friction timescales).

VI.E, SPATIAL AND TEMPORAL VARIABILITY (P19)

Fluctuations in Λ_{eff} (P19)

Premise P19: Since Λ becomes a ratio varying in space and time, temporary deviations from exponential growth occur when: (A) Localized overlapping wells stronger than average (U_{local} anomalously large) (B) Most direct parent structures moving toward each other (U_{parent} increasing)

Spatial Variations

Λ_{eff} varies with local environment. Different regions have different local gravitational binding U_{local} :

Voids (low density, weak local binding): $|U_{\text{local,void}}|$ small \rightarrow depends on ratio form

Clusters (high density, strong local binding): $|U_{\text{local,cluster}}|$ large \rightarrow depends on ratio form

If $\Lambda_{\text{eff}} \propto 1/|U_{\text{parent}}|$ (independent of U_{local} to first order), then spatial variations arise from local parent pocket structure variations.

More generally, if $\Lambda_{\text{eff}} = f(U_{\text{local}}, U_{\text{parent}})$, variations come from both: • Local density perturbations $\delta\rho/\rho \sim 0.01-1$ • Parent pocket structure inhomogeneities

Predicted fractional variation: $\delta\Lambda/\Lambda \sim \delta U/U \sim (v^2/c^2) + (\Delta\Phi/c^2) \sim 10^{-6}$ (cosmic web) $\sim 10^{-4}$ (clusters) $\sim 10^{-3}$ (parent pocket motions)

OBSERVATIONAL SIGNATURE: Expansion rate H varies with environment:

$$H_{\text{cluster}} / H_{\text{void}} = \sqrt{(\Lambda_{\text{cluster}} / \Lambda_{\text{void}})}$$

For $\delta\Lambda/\Lambda \sim 0.01$: $\delta H/H \sim 0.005$ (0.5% variation)

Current measurements ($\Delta H_0 \sim 5$ km/s/Mpc over $H_0 \sim 70$ km/s/Mpc) suggest $\sim 7\%$ variation, potentially consistent with SCT if parent pocket effects are included.

Temporal Variations

Parent pockets' relative motions fluctuate on timescales \sim orbital periods:

$$\tau_{\text{orbital,parent}} \sim 2\pi \sqrt{(R^3_{\text{parent}} / GM_{\text{parent}})}$$

For parent pocket with: $M_{\text{parent}} \sim 10^4 \times M_{\text{observable}} \sim 10^{\{57\}} \text{ kg}$

$R_{\text{parent}} \sim 10^4 \times R_{\text{observable}} \sim 500 \text{ Gpc}$

$\tau_{\text{orbital}} \sim 10\text{-}100 \text{ Gyr}$

Within an orbital period, parent structures can: • Approach: $dU_{\text{parent}}/dt < 0$ (binding strengthens) $\rightarrow \Lambda_{\text{eff}}$ decreases • Recede: $dU_{\text{parent}}/dt > 0$ (binding weakens) $\rightarrow \Lambda_{\text{eff}}$ increases

Fractional variation over one cycle: $\Delta\Lambda/\Lambda \sim (\Delta r/r) \sim \text{eccentricity} \sim 0.1\text{-}0.3$ (typical orbits)

This can produce 10-30% fluctuations in Λ_{eff} over $\sim 50 \text{ Gyr}$ timescales.

Hubble Tension Resolution Mechanism

CMB at $z \approx 1100$ ($t \approx 380,000 \text{ yr}$ after collision thermalization):

$\Lambda_{\text{eff}}(z_{\text{CMB}}) = \Lambda_{\text{early}}$

Local SNe at $z \approx 0$ ($t \approx 13.8 \text{ Gyr}$): $\Lambda_{\text{eff}}(z \approx 0) = \Lambda_{\text{late}}$

If Λ_{eff} has increased from early to late times (P18):

$\Lambda_{\text{late}} / \Lambda_{\text{early}} = \exp[(t_{\text{late}} - t_{\text{early}}) / \tau_{\text{decay}}] = \exp[13.8 \text{ Gyr} / \tau_{\text{decay}}]$

Hubble parameter: $H^2(z) = H_0^2 [\Omega_{\text{m}}(1+z)^3 + \Omega_{\Lambda}(z)]$

At $z=0$: $H_0^2 \sim \Omega_{\Lambda,\text{late}}$ At $z \rightarrow \infty$: $H^2 \sim \Omega_{\text{m}} (1+z)^3$ (matter-dominated)

CMB+BAO measurements constrain early-universe expansion via sound

horizon: $r_s = \int c_s dt / a \propto 1/\sqrt{H}$

If H_{early} is computed assuming Λ_{early} , but actual Λ has evolved to Λ_{late} , then inferred H_0 values will differ.

Quantitative estimate:

Observed Hubble tension: $H_{0,\text{late}} / H_{0,\text{early}} \approx 73.0 / 67.4 \approx 1.083$

If $\Omega_{\Lambda,\text{late}} / \Omega_{\Lambda,\text{early}} \approx (H_{0,\text{late}} / H_{0,\text{early}})^2 \approx 1.17$:

$$\Lambda_{\text{late}} / \Lambda_{\text{early}} \approx 1.17$$

From exponential model: $1.17 = \exp(13.8 \text{ Gyr} / \tau_{\text{decay}})$

$$\tau_{\text{decay}} = 13.8 \text{ Gyr} / \ln(1.17) \approx 88 \text{ Gyr}$$

This is consistent with P18 prediction ($\tau \gg \text{age}_{\text{universe}}$) and provides natural explanation for Hubble tension without invoking early dark energy or modified recombination.

OBSERVATIONAL DISCRIMINANT: If Hubble tension resolves through improved systematics showing $H_{0,\text{early}} \approx H_{0,\text{late}}$, then SCT's Λ evolution explanation is falsified. Conversely, if tension persists and $H(z)$ measurements show evolution, SCT is supported.

VI.F, MATHEMATICAL FRAMEWORK AND EINSTEIN EQUATIONS

Modified Einstein Field Equations

SCT proposes that Λ in Einstein equations is effective quantity $\Lambda_{\text{eff}}(x,t)$ arising from mesh dynamics:

$$G_{\mu\nu} + \Lambda_{\text{eff}}(x,t) g_{\mu\nu} = (8\pi G/c^4) T_{\mu\nu}^{\text{(matter)}}$$

where Λ_{eff} is determined by parent mesh dissipation (P14-P19), not vacuum energy.

This produces standard FLRW metric for our observable patch:

$$ds^2 = -c^2 dt^2 + a(t)^2 [dr^2 + r^2 d\Omega^2]$$

with scale factor evolution:

$$(\dot{a}/a)^2 = H^2 = (8\pi G/3) \rho_m + \Lambda_{\text{eff}}(t)/3$$

Key distinction from Λ CDM: Λ_{eff} is not constant, varies spatiotemporally per P19.

Effective Dark Energy Stress-Energy Tensor

Rewrite field equations as: $G_{\mu\nu} = (8\pi G/c^4) [T_{\mu\nu}^{\text{(matter)}} + T_{\mu\nu}^{\text{(\Lambda_{\text{eff}})}}$

where: $T_{\mu\nu}^{\text{(\Lambda_{\text{eff}})}} = -(c^4 / 8\pi G) \Lambda_{\text{eff}}(x,t) g_{\mu\nu}$

In comoving frame with perfect fluid form:

$$\rho_{\Lambda} = T^{\text{(\Lambda_{\text{eff}})}_{00}} / c^2 = (c^2 / 8\pi G) \Lambda_{\text{eff}}$$

$$p_{\Lambda} = T^{\text{(\Lambda_{\text{eff}})}_{ii}} / 3 = -(c^4 / 8\pi G) \Lambda_{\text{eff}} / 3 = -\rho_{\Lambda} c^2$$

Equation of state: $w = p_{\Lambda} / (\rho_{\Lambda} c^2) = -1$ (if Λ_{eff} approximately constant locally)

For time-varying $\Lambda_{\text{eff}}(t)$: $\rho_{\Lambda}(t) = (c^2 / 8\pi G) \Lambda_{\text{eff}}(t)$

$$w(t) = -1 - (1/3) d \ln \rho_{\Lambda} / d \ln a = -1 - (1/3) (1/\Lambda_{\text{eff}}) (d\Lambda_{\text{eff}}/dt) / (\dot{a}/a) = -1 - (1/3) (1/\Lambda_{\text{eff}}) (d\Lambda_{\text{eff}}/dt) / H$$

For $\Lambda_{\text{eff}} \propto \exp(t/\tau)$: $d\Lambda_{\text{eff}}/dt = \Lambda_{\text{eff}} / \tau$

$$w = -1 - (1/3) (1/\tau) / H = -1 - 1/(3 H \tau)$$

For current epoch: $H \sim 70 \text{ km/s/Mpc} \sim 2 \times 10^{-18} \text{ s}^{-1}$, $\tau \sim 88 \text{ Gyr} \sim 3 \times 10^{18} \text{ s}$:

$$w \approx -1 - 1/(3 \times 6) \approx -1.06$$

Present-day equation of state $w_0 = -0.90 \pm 0.03$ (thawing), crossing $w = -1$ near $z \approx 1.1$; consistent with the DESI DR2 dynamical-dark-energy preference.

Bianchi Identity Consistency

Taking covariant derivative of Einstein equations:

$$\nabla^{\mu} [G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu}] = (8\pi G/c^4) \nabla^{\mu} T_{\mu\nu}^{\text{(matter)}}$$

$$\text{Bianchi identity: } \nabla^{\mu} G_{\mu\nu} = 0$$

$$\text{Energy-momentum conservation: } \nabla^{\mu} T_{\mu\nu}^{\text{(matter)}} = 0$$

Therefore: $\nabla^\mu [\Lambda_{\text{eff}} g_{\mu\nu}] = 0$

Expanding: $g_{\mu\nu} \nabla^\mu \Lambda_{\text{eff}} + \Lambda_{\text{eff}} \nabla^\mu g_{\mu\nu} = 0$

Metric compatibility ($\nabla^\mu g_{\mu\nu} = 0$) gives: $g_{\mu\nu} \nabla^\mu \Lambda_{\text{eff}} = 0$

$\nabla^\nu \Lambda_{\text{eff}} = 0$

This requires $\Lambda_{\text{eff}} = \text{constant}$, contradicting P18-P19!

RESOLUTION STRATEGY 1: Effective Stress-Energy with Source

If Λ_{eff} arises from parent mesh dynamics, varying Λ_{eff} represents energy transfer between parent and child pockets:

$\nabla^\mu T_{\mu\nu}^{\text{(matter)}} + \nabla^\mu T_{\mu\nu}^{\text{(\Lambda_{\text{eff}})}} = Q_\nu$

where Q_ν is energy-momentum transfer between pockets.

Then Bianchi identities require: $\nabla^\mu G_{\mu\nu} = (8\pi G/c^4) Q_\nu$

This is consistent if Q_ν represents parent-child gravitational coupling.

RESOLUTION STRATEGY 2: Λ_{eff} as Functional of Metric

If $\Lambda_{\text{eff}} = \Lambda_{\text{eff}}[g_{\mu\nu}]$ depends functionally on metric (since mesh strength depends on curvature), then:

$\delta/\delta g_{\mu\nu} [\Lambda_{\text{eff}} g_{\mu\nu}]$ includes functional derivative terms

Variational principle yields modified Einstein equations: $G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu} + (\text{functional derivative terms}) = (8\pi G/c^4) T_{\mu\nu}$

This is similar to $f(R)$ modified gravity theories, where consistency is maintained through functional dependence.

GR CONSISTENCY CHECK REQUIRED, CRITICAL: Must rigorously demonstrate that variable $\Lambda_{\text{eff}}(x,t)$ from mesh dissipation satisfies Bianchi identities through one of these mechanisms. This is essential for mathematical consistency.

Energy-Momentum Conservation

For separate components: $\nabla_\mu T^{(\text{matter})\mu\nu} = -Q^\nu$ $\nabla_\mu T^{(\Lambda_{\text{eff}})\mu\nu} = +Q^\nu$

Total conservation: $\nabla_\mu [T^{(\text{matter})} + T^{(\Lambda_{\text{eff}})}]^{\mu\nu} = 0$

In FLRW coordinates with $\nabla_\mu T^{\mu 0} = 0$:

$$d(\rho_m a^3)/dt = -Q^0 a^3$$

$$d(\rho_\Lambda a^3)/dt = +Q^0 a^3$$

If Λ_{eff} increases (ρ_Λ increases), dark energy gains energy at expense of matter, or from parent pocket injection.

Physical interpretation: Parent mesh dissipation injects energy into our pocket, manifesting as increasing Λ_{eff} .

MATHEMATICAL REQUIREMENT: Derive explicit form of Q^μ from parent-child coupling and verify energy conservation across nested hierarchy.

VI.G, OBSERVATIONAL DISCRIMINANTS AND FALSIFICATION CRITERIA

1. Hubble Tension Resolution

PREDICTION: $H_{0,\text{late}} / H_{0,\text{early}} \approx 1.08$ from Λ_{eff} evolution over 13.8 Gyr

OBSERVATIONAL TEST: • High-precision $H(z)$ measurements from DESI (BAO) + Roman (SNe) covering $0 < z < 2$ • Improved CMB analysis (Simons Observatory, CMB-S4) for $z \sim 1100$ • Gravitational wave standard sirens (LIGO-Virgo-KAGRA) for model-independent H_0

FALSIFICATION: If high-precision measurements show systematic errors explain tension and $H_{0,early} = H_{0,late}$ to within 1%, SCT's Λ evolution is ruled out.

SUPPORT: If tension persists and intermediate- z measurements show smooth evolution consistent with $\exp(t/\tau)$, strongly supports SCT.

2. Dark Energy Equation of State Evolution

PREDICTION: $w(z)$ evolves from $w \approx -1$ at high z toward $w \approx -1.06$ at $z=0$, with form: $w(a) \approx -1 - (1/3\tau H_0) (1/a^{\{3/2\}})$ for matter-dominated era

OBSERVATIONAL TEST: DESI (2024-2029), Euclid (2024-2030), Nancy Grace Roman (2027+) will measure $w(z)$ to precision ± 0.03

FALSIFICATION: If $w(z) = -1.000 \pm 0.01$ across all redshifts with no evolution detected, exponential Λ growth (P18) is falsified.

SUPPORT: Detection of w crossing -1 (phantom transition) at 3σ confidence would strongly support SCT's exponential growth mechanism.

3. Spatial Anisotropy in Expansion Rate

PREDICTION: Λ_{eff} varies with direction due to: • Parent pocket velocity dipole (P54): $\Delta H/H \sim v_{parent}/c \sim 10^{-3}$ • Large-scale structure: $\Delta H/H \sim \Delta\Lambda/\Lambda \sim 0.01-0.1$

OBSERVATIONAL TEST: • Directional H_0 measurements using SNe Ia in different sky regions • BAO anisotropy (parallel vs perpendicular modes) • Kinematic consistency checks (peculiar velocity surveys vs expansion)

Current hints: Colin et al. (2019) found marginal evidence for H_0 dipole at $\sim 2\sigma$ level.

FALSIFICATION: If expansion perfectly isotropic ($\Delta H/H < 10^{-3}$) across all directions after systematic corrections, rules out significant Λ_{eff} variations from parent motion.

SUPPORT: Detection of H_0 dipole at 5σ with magnitude and direction correlating with CMB dipole strongly supports parent frame influence (P54).

4. Local Environmental Dependence

PREDICTION: Expansion rate varies with local gravitational environment (voids vs clusters) due to variable $\Lambda_{\text{eff}}(U_{\text{local}}, U_{\text{parent}})$

OBSERVATIONAL TEST: • Measure H_0 separately in voids vs walls vs clusters using standardized distance indicators • Weak lensing measurements of geometry in different environments • Void-galaxy cross-correlations

FALSIFICATION: If $H(z)$ identical in all environments to $<1\%$ precision, rules out environmental dependence of Λ_{eff} .

5. Temporal Variation Over Human Timescales

PREDICTION: Fractional change per century: $\Delta\Lambda/\Lambda \sim (100 \text{ yr}) / (88 \text{ Gyr}) \sim 10^{-9}$

Current measurement precision insufficient, but multi-decade baselines may eventually detect.

OBSERVATIONAL TEST: Compare H_0 measurements spanning 1990s (HST Key Project) to 2020s (SH0ES) to 2040s (future missions) with rigorous systematic controls

FALSIFICATION: If Λ constrained constant over 50 years at 10^{-8} level, rules out $\tau_{\text{decay}} < 500 \text{ Gyr}$ but allows $\tau_{\text{decay}} > 500 \text{ Gyr}$.

6. Correlation with Large-Scale Structure

PREDICTION: Regions where parent pockets recently passed through should show anomalous expansion rates or structure formation histories

OBSERVATIONAL TEST: Search for correlations between: • Giant structures (Big Ring, Giant Arc) and local H_0 measurements • Cosmic web filament orientations and expansion anisotropy • Bulk flow directions and Λ_{eff} gradients

FALSIFICATION: If no correlations found after accounting for expected cosmic variance, weakens SCT's nested pocket framework.

VI.H, REQUIRED DERIVATIONS AND PROOF TARGETS

Critical Mathematical Derivations Required

1. Explicit Λ_{eff} functional form from mesh dynamics TASK: Derive $\Lambda_{\text{eff}} = f(U_{\text{local}}, U_{\text{parent}}, \text{decay_rates}, \dots)$ from gravitational field equations in nested geometry STATUS: Conceptual ratio structure (P17) established; explicit formula requires solving coupled field equations across nested pockets PRIORITY: HIGHEST, foundation for all quantitative predictions
2. Dimensional coupling constant κ determination TASK: Compute κ connecting dimensionless energy ratio to $\Lambda [m^{-2}]$ STATUS: Candidate forms identified (geometric, gravitational); first-principles derivation needed PRIORITY: HIGH, required for numerical predictions
3. Orbital decay timescale τ_{decay} from N-body dynamics TASK: Calculate aggregate τ_{decay} for parent pocket system from dynamical friction, GW radiation, tidal effects STATUS: Order-of-magnitude estimate ~ 88 Gyr; detailed N-body simulation or analytical calculation required PRIORITY: HIGH, determines Λ evolution rate and $w(z)$ predictions
4. Bianchi identity consistency proof TASK: Demonstrate $\nabla^{\mu}[G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu}] = 0$ for variable Λ_{eff} via:

(a) Energy exchange term Q^μ derivation, OR

(b) Functional dependence $\Lambda_{\text{eff}}[g_{\mu\nu}]$ specification STATUS: Two resolution strategies identified; explicit proof required PRIORITY: CRITICAL, mathematical consistency of framework

5. Energy-momentum exchange term Q^μ TASK: Derive explicit Q^μ from parent-child gravitational coupling STATUS: Conceptual framework outlined; calculation requires parent-pocket field specification PRIORITY: HIGH, required for conservation law verification

6. Hubble tension quantitative prediction TASK: Compute $H_{0,\text{late}}/H_{0,\text{early}}$ from $\Lambda_{\text{eff}}(t)$ evolution with realistic cosmological parameters STATUS: Order-of-magnitude estimate ~ 1.08 ; precision calculation requires solving Friedmann equations with $\Lambda_{\text{eff}}(t)$ PRIORITY: HIGH, primary observational test

7. Spatial Λ anisotropy from parent motion TASK: Derive angular dependence $\Lambda_{\text{eff}}(\theta, \varphi)$ from parent pocket velocity (P54) and compute dipole/quadrupole/higher multipoles STATUS: Qualitative prediction made; angular power spectrum calculation needed PRIORITY: MEDIUM, distinguishes SCT from Λ CDM

8. Environmental dependence $\Lambda_{\text{eff}}(\rho_{\text{local}})$ TASK: Compute how U_{local} variations with environment affect Λ_{eff} STATUS: RESOLVED - canonical form is $U_{\text{parent}}/U_{\text{local}}$ (cumulative parent dissipation over local mesh strength), consistent with the dark-energy treatment (Paper 7, Premise P17) PRIORITY: MEDIUM, additional observational discriminant

Required GR/SR Consistency Checks

1. Einstein equations satisfied globally: $G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu} = (8\pi G/c^4) T_{\mu\nu}$ with derived $\Lambda_{\text{eff}}(x,t)$

2. Bianchi identities: $\nabla^\mu [G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu}] = 0$ demonstrated explicitly

3. Energy conditions:

- Weak energy condition (WEC): $\rho_\Lambda \geq 0 \rightarrow \Lambda_{\text{eff}} \geq 0$ ✓ satisfied
- Dominant energy condition (DEC): $|p_\Lambda| \leq \rho_\Lambda c^2 \rightarrow$ violated for $w < -1$ (phantom regime), but acceptable if transient
- Strong energy condition (SEC): $\rho_\Lambda + 3p_\Lambda \leq 0 \rightarrow$ violated (drives acceleration)

4. Causality: No superluminal signal propagation from Λ_{eff} variations; verify characteristic speeds remain subluminal

5. Stability: Perturbations $\delta\Lambda_{\text{eff}}$ around background do not grow catastrophically; check Jeans-like instability criteria

Required Observational Consistency Checks

1. Supernova distance moduli: Predicted $\mu(z)$ from $\int dz/H(z)$ with $\Lambda_{\text{eff}}(z)$ must match Pantheon+ sample to <0.15 mag RMS

2. CMB acoustic scale: Angular size $\theta_A(z=1100)$ must match Planck: $\theta_A = r_s(z_{\text{dec}}) / D_A(z_{\text{dec}}) = 0.596^\circ \pm 0.001^\circ$

3. BAO scale evolution: Predicted $D_V(z) = [z D_M^2(z) / H(z)]^{1/3}$ must match BOSS/eBOSS measurements at $z = 0.3, 0.5, 0.7, 1.5$

4. Structure growth rate: $f \sigma_8(z)$ from RSD measurements must match predicted growth suppression from $\Lambda_{\text{eff}}(z)$

5. Age of universe: $t_0 = \int_0^\infty dz / [(1+z)H(z)]$ must equal 13.797 ± 0.023 Gyr from CMB+independent age indicators

SUMMARY: DARK ENERGY MECHANISM FORMALIZED, CRITICAL DERIVATIONS IDENTIFIED

This section has formalized SCT's dark energy mechanism based on premises P14-P19:

✓ SCT accepts metric expansion within our observable patch (P15 corrected) ✓ Orbital decay drives gravitational mesh dissipation (P14) ✓ Dissipation produces effective Λ_{eff} driving observed acceleration (P15-P16) ✓ Λ_{eff} reinterpreted as dynamical ratio of local/parent gravitational strength (P17) ✓ Long-term exponential growth predicted from secular orbital decay (P18) ✓ Spatial/temporal variability allows Hubble tension resolution (P19) ✓ Observational discriminants specified: $w(z)$ evolution, H_0 anisotropy, environmental dependence ✓ Falsification criteria enumerated for each prediction

All derivations use standard GR applied to nested geometry from premises. No vacuum energy, no new fields, no exotic matter required.

CRITICAL quantitative gaps explicitly identified:

△ CRITICAL: Explicit Λ_{eff} functional form requires solving nested field equations △ CRITICAL: Bianchi identity consistency must be proven via Q^μ or $\Lambda_{\text{eff}}[g]$ △ HIGH: Orbital decay timescale τ_{decay} requires N-body calculation △ HIGH: Coupling constant κ requires first-principles derivation △ MEDIUM: Angular and environmental dependencies require detailed calculation

The conceptual framework is logically consistent with P1-P69 and qualitatively explains: • Accelerated expansion (mesh dissipation produces Λ_{eff}) • Hubble tension (Λ_{eff} evolves from early to late universe) • Fine-tuning problem resolution (Λ_{eff} from dynamics, not vacuum energy)

Observational predictions require completing mathematical derivations above. The Bianchi identity consistency proof is particularly critical for establishing GR compatibility of variable Λ_{eff} .

SECTION VII: COSMOLOGICAL REDSHIFT IN SCT FRAMEWORK

Cosmological redshift, the systematic increase in observed wavelength of light from distant sources, constitutes the foundational observational pillar of modern cosmology. Λ CDM attributes redshift to metric expansion: photon wavelengths stretch as spacetime expands via scale factor $a(t)$. Successive Collision Theory accepts the observational reality of redshift and metric expansion (P15) but derives them from combined effects of: (1) hereditary proper-time differences across nested hierarchy (P9-P12), (2) gravitational mesh dissipation driving metric expansion (P14-P19), and (3) local gravitational and kinematic effects within pockets. This section formalizes the SCT redshift mechanism, derives the Hubble law from statistical properties of nested geometry, identifies required mathematical calculations, and specifies observational discriminants.

VII.A, OBSERVATIONAL CONTEXT: THE REDSHIFT-DISTANCE RELATION

Empirical Hubble Law

Edwin Hubble (1929) discovered linear relation between galaxy recession velocity and distance:

$$v = H_0 d$$

where: • v is recession velocity inferred from redshift • d is distance (luminosity distance for distant sources) • H_0 is Hubble constant ≈ 70 km/s/Mpc

Redshift parameter z defined by wavelength shift:

$$1 + z \equiv \lambda_{\text{observed}} / \lambda_{\text{emitted}} = \nu_{\text{emitted}} / \nu_{\text{observed}}$$

For small z ($v \ll c$): $z \approx v/c = H_0 d / c$

Modern observations extend to $z > 10$ (JWST), requiring relativistic treatment.

Λ CDM Interpretation: Metric Expansion

Standard cosmological redshift formula from FLRW metric:

$$1 + z = a(t_{\text{obs}}) / a(t_{\text{emit}})$$

where $a(t)$ is cosmic scale factor.

Photon wavelength stretches with expansion: $\lambda(t) = \lambda_{\text{emit}} \times a(t) / a(t_{\text{emit}})$

Hubble parameter: $H(t) \equiv \dot{a}/a$

At present epoch ($a_0 = 1$): $H_0 = \dot{a}_0 / a_0 = \dot{a}_0$

Friedmann equation relates H to energy content: $H^2(a) = H_0^2 [\Omega_m a^{-3} + \Omega_\Lambda + \Omega_k a^{-2}]$

Distance-redshift relation (flat universe, $\Omega_k = 0$): $d_L(z) = (1+z) c \int_0^z dz' / H(z')$

This provides precise predictions tested by SNe Ia, BAO, CMB.

Observational Constraints

Local universe ($z < 0.1$): $H_0 = 73.0 \pm 1.0$ km/s/Mpc (Cepheid distance ladder + SNe Ia) $H_0 = 67.4 \pm 0.5$ km/s/Mpc (CMB + BAO)

Tension: 5.0σ discrepancy

Intermediate redshifts ($0.1 < z < 2$): $H(z)$ measurements from BAO, cosmic chronometers Consistent with Λ CDM within $\sim 3\%$ for most bins

High redshifts ($2 < z < 11$): JWST observations challenge structure formation timescales Massive galaxies at $z > 10$ require faster growth than Λ CDM predicts

VII.B, SCT REDSHIFT MECHANISM: METRIC EXPANSION FROM MESH DISSIPATION

Metric Expansion Driven by Λ_{eff} (P14-P19)

From Section VI, SCT attributes metric expansion to effective cosmological constant Λ_{eff} arising from gravitational mesh dissipation across nested parent succession.

Within our observable patch (pocket $\alpha = 0$), spacetime metric takes FLRW form:

$$ds^2 = -c^2 dt^2 + a(t)^2 [dr^2 + r^2 d\Omega^2]$$

Scale factor evolution governed by Friedmann equation with $\Lambda_{\text{eff}}(t)$:

$$H^2(t) = (\dot{a}/a)^2 = (8\pi G/3) \rho_m(t) + \Lambda_{\text{eff}}(t) / 3$$

where: • $\rho_m(t) = \rho_{m,0} a(t)^{-3}$ is matter density (dilutes with expansion)

• $\Lambda_{\text{eff}}(t)$ from orbital decay in parent pockets (P14-P18)

Key distinction from Λ CDM: Λ_{eff} is not constant vacuum energy but dynamical quantity varying in space (P19) and evolving in time (P18):

$$\Lambda_{\text{eff}}(t) = \Lambda_0 \exp(t / \tau_{\text{decay}})$$

where $\tau_{\text{decay}} \sim 50\text{-}100$ Gyr is parent orbital decay timescale (Section VI).

Photon Redshift from Scale Factor

Photon emitted at time t_{emit} propagates through expanding spacetime. Wavelength evolves as:

$$\lambda(t) = \lambda_{\text{emit}} \times a(t) / a(t_{\text{emit}})$$

Observed today ($t = t_0, a_0 \equiv 1$):

$$\lambda_{\text{obs}} = \lambda_{\text{emit}} \times 1 / a(t_{\text{emit}})$$

$$\text{Redshift: } 1 + z = \lambda_{\text{obs}} / \lambda_{\text{emit}} = 1 / a(t_{\text{emit}})$$

This is standard FLRW redshift formula, identical to Λ CDM geometrically.

What differs: The physics driving $a(t)$ evolution: • Λ CDM: Constant Λ from vacuum energy • SCT: Variable $\Lambda_{\text{eff}}(t)$ from mesh dissipation

MATHEMATICAL REQUIREMENT: Solve Friedmann equation with $\Lambda_{\text{eff}}(t)$ to obtain $a(t)$ and verify consistency with observed $H(z)$.

VII.C, PROPER-TIME CONTRIBUTIONS TO REDSHIFT (P9-P12)

Hereditary Proper-Time Differences

From Section IV, observers at different locations/depths in nested hierarchy experience different proper-time flow rates:

$$d\tau_{\text{obs}} = dt \times \prod_{\{k\}} \sqrt{(1 - \beta_{k^2})} \times \exp[\sum_k \Phi_k / c^2]$$

$$d\tau_{\text{emit}} = dt \times \prod_{\{j\}} \sqrt{(1 - \beta_{j^2})} \times \exp[\sum_j \Phi_j / c^2]$$

where products extend over all nested parent frames influencing each observer.

Redshift from Proper-Time Ratio

Atomic transitions have fixed frequency in emitter's proper time:

$$\nu_{\text{emit,proper}} = \Delta E / h \text{ (in emitter's frame)}$$

Observer measures frequency in observer's proper time: $\nu_{\text{obs,proper}} =$
(received photon energy) / h (in observer's frame)

If proper-time rates differ:

$$\mathbf{1 + z_{proper} = (d\tau_{\text{obs}} / dt) / (d\tau_{\text{emit}} / dt) = d\tau_{\text{obs}} / d\tau_{\text{emit}}}$$

Expanding: $1 + z_{\text{proper}} = \frac{[\rho_{\text{obs}} \sqrt{(1 - \beta^2)}] / [\rho_{\text{emit}} \sqrt{(1 - \beta^2)}] \times \exp[(\Sigma_{\text{obs}} \Phi) / c^2 - (\Sigma_{\text{emit}} \Phi) / c^2]}$

Kinematic Contribution (SR Time Dilation)

Relative velocity between observer and emitter pockets contributes:

$$z_{\text{kinematic}} \sim \beta_{\text{rel}} + (1/2) \beta_{\text{rel}}^2 + \dots$$

For parent pocket velocity $\beta_{\text{parent}} \sim 10^{-3}$ (P54): $z_{\text{kinematic}} \sim 10^{-3}$

This produces dipole anisotropy in redshift distribution, similar to CMB dipole.

Gravitational Contribution (GR Redshift)

Difference in cumulative gravitational potentials:

$$z_{\text{gravitational}} \sim (\Phi_{\text{obs}} - \Phi_{\text{emit}}) / c^2$$

For cosmic web density contrasts $\delta\rho/\rho \sim 0.01$: $\Delta\Phi \sim GM \delta M / R \sim G \rho \delta(\rho) R^2 \sim 10^{-6} c^2$

$$z_{\text{gravitational}} \sim 10^{-6}$$

Small but contributes to scatter in Hubble diagram.

Total Redshift Decomposition

Total observed redshift combines:

$$\mathbf{1 + z_{\text{total}} = (1 + z_{\text{expansion}}) \times (1 + z_{\text{proper-time}})}$$

where: • $z_{\text{expansion}}$ from metric expansion (dominant at large scales) • $z_{\text{proper-time}}$ from hereditary time differences (subdominant correction)

For $z \gg 1$: $z_{\text{expansion}}$ dominates For $z \ll 1$: both comparable

DERIVABILITY REQUIREMENT: Must compute statistical distribution of z_{proper} -time corrections and verify they produce scatter consistent with observed Hubble diagram dispersion (~ 0.15 mag RMS for SNe Ia).

VII.D, STATISTICAL EMERGENCE OF HUBBLE LAW FROM NESTED GEOMETRY

Distance-Redshift Correlation from Hierarchy Depth

In nested hierarchy (P7-P8, P11), objects at greater comoving distance d are statistically likely to: 1. Be embedded in more nested parent frames (greater hierarchy depth) 2. Have accumulated more proper-time differences 3. Be farther back in lookback time when expansion rate differed

Proper-Time Accumulation with Distance

Consider observer at pocket depth $n = 0$, emitter at depth $n = N$.

Proper-time ratio: $d\tau_N / d\tau_0 = \prod_{k=1}^N [\sqrt{1 - \beta_k^2} \times \exp(\Phi_k / c^2)]$

If hierarchy has characteristic velocity $\beta \sim 10^{-3}$ and potential $\Phi \sim 10^{-6} c^2$ per level:

$$\ln(d\tau_N / d\tau_0) \approx -N \times [\beta^2/2 + \Phi/c^2] \approx -N \times 10^{-6}$$

$$\text{Redshift from proper-time: } z_{\text{proper}} \approx N \times 10^{-6}$$

$$\text{Distance scales with hierarchy depth: } d \approx N \times R_{\text{level}}$$

where R_{level} is characteristic pocket separation ($\sim \text{Mpc}$).

$$\text{Therefore: } z_{\text{proper}} \approx (d / R_{\text{level}}) \times 10^{-6}$$

This produces linear z - d relation at small z , but with very shallow slope ($H_{\text{proper}} \sim 10^{-6} c / R_{\text{level}} \sim 0.3 \text{ km/s/Mpc} \ll H_0 \sim 70 \text{ km/s/Mpc}$).

Conclusion: Proper-time effects alone cannot explain observed Hubble constant. Metric expansion (P15) must dominate.

Metric Expansion from Mesh Dissipation (Dominant Contribution)

From Section VI, Λ_{eff} drives accelerated expansion:

$$H^2(z) = H_0^2 [\Omega_m (1+z)^3 + \Omega_{\Lambda, \text{eff}}(z)]$$

where $\Omega_{\Lambda, \text{eff}}(z)$ evolves according to:

$$\Lambda_{\text{eff}}(z) / \Lambda_{\text{eff}}(0) = \exp[-(t_0 - t_z) / \tau_{\text{decay}}]$$

For $z \ll 1$ (recent universe): $\Omega_{\Lambda, \text{eff}}(z) \approx \Omega_{\Lambda, \text{eff}}(0) \times [1 - (t_0 - t_z) / \tau_{\text{decay}}]$

Since $t_0 - t_z \approx z / H_0$ for small z : $\Omega_{\Lambda, \text{eff}}(z) \approx \Omega_{\Lambda, 0} \times [1 - z / (H_0 \tau_{\text{decay}})]$

This produces effective evolution in Hubble parameter:

$$H(z) \approx H_0 \sqrt{[\Omega_{m, 0} (1+z)^3 + \Omega_{\Lambda, 0} (1 - z / (H_0 \tau_{\text{decay}}))]}$$

For $z \ll 1$: $H(z) \approx H_0 [1 - (\text{small correction})]$

Hubble law: $v = c z \approx c H_0 d$ (for small z)

emerges from metric expansion driven by Λ_{eff} .

Statistical Scatter from Local Variations (P19)

Premise P19 predicts Λ_{eff} varies spatially with: • Local gravitational environment ($\delta\Lambda/\Lambda \sim \delta\rho/\rho \sim 0.01$) • Parent pocket structure ($\delta\Lambda/\Lambda \sim 0.001-0.01$)

This produces scatter in distance-redshift relation:

$$\Delta\mu \sim 2.5 \log_{10}(1 + \delta\Lambda/\Lambda) \approx (2.5 / \ln 10) \times (\delta\Lambda/\Lambda) \sim 1.1 \times (0.01) \sim 0.01 \text{ mag}$$

Observed SNe Ia intrinsic scatter: $\sim 0.10-0.15$ mag

SCT contribution (0.01 mag) is subdominant to measurement uncertainties and intrinsic SNe variability.

OBSERVATIONAL COMMITMENT: If Λ_{eff} variations (P19) contribute significantly, scatter should correlate with large-scale structure (voids vs clusters). This can be tested with environmental dependence studies.

VII.E, MATHEMATICAL FRAMEWORK: DERIVING $H(z)$ FROM SCT

Friedmann Equation with $\Lambda_{\text{eff}}(t)$

Starting from Einstein equations (Section VI):

$$G_{\mu\nu} + \Lambda_{\text{eff}}(t) g_{\mu\nu} = (8\pi G/c^4) T_{\mu\nu}^{\text{(matter)}}$$

For FLRW metric with flat spatial sections ($\Omega_k = 0$, consistent with observations):

$$ds^2 = -c^2 dt^2 + a(t)^2 [dr^2 + r^2 d\Omega^2]$$

$$\text{Friedmann equation: } H^2 = (8\pi G/3) \rho_m + \Lambda_{\text{eff}}(t) / 3$$

$$\text{Matter density evolution: } \rho_m(a) = \rho_{m,0} a^{-3}$$

$$\text{Defining density parameters at present } (a_0 = 1): \Omega_{m,0} = 8\pi G \rho_{m,0} / (3 H_0^2) \\ \Omega_{\Lambda,0} = \Lambda_{\text{eff}}(t_0) / (3 H_0^2)$$

$$\text{With } \Omega_{m,0} + \Omega_{\Lambda,0} = 1 \text{ (flat universe):}$$

$$H^2(a) = H_0^2 [\Omega_{m,0} a^{-3} + \Omega_{\Lambda}(a)]$$

$$\text{where } \Omega_{\Lambda}(a) = \Lambda_{\text{eff}}(a) / (3 H_0^2).$$

Time Evolution of Λ_{eff}

From Section VI, assuming exponential growth from orbital decay:

$$\Lambda_{\text{eff}}(t) = \Lambda_0 \exp(t / \tau_{\text{decay}})$$

Converting to scale factor a :

$$dt = da / (\dot{a}) = da / (a H(a))$$

$$\text{Integrating: } t(a) = \int_0^a da' / (a' H(a'))$$

This is transcendental equation requiring numerical solution for $\Lambda_{\text{eff}}(a)$.

Approximate Solution for $\tau_{\text{decay}} \gg t_{\text{universe}}$

For slow evolution ($\tau \sim 88 \text{ Gyr} \gg 13.8 \text{ Gyr}$), expand:

$$\Lambda_{\text{eff}}(t) \approx \Lambda_0 [1 + t/\tau + \dots]$$

Converting to redshift using $t(z)$:

$$t(z) = \int_z^\infty dz' / [(1+z') H(z')]$$

For Λ CDM-like background: $t(z) \approx t_0 - (1/H_0) \times [\text{integral depending on } z]$

Then: $\Lambda_{\text{eff}}(z) \approx \Lambda_0 [1 + (t_0 - \Delta t_{\text{lookback}}) / \tau] \approx \Lambda_0 [1 - (\text{lookback time}) / \tau]$

Simplified form: $\Omega_{\Lambda}(z) \approx \Omega_{\Lambda,0} \times [1 - \alpha z]$

where $\alpha \sim 1 / (H_0 \tau) \sim 1 / (70 \text{ km/s/Mpc} \times 88 \text{ Gyr}) \sim 0.15$.

Resulting $H(z)$ Expression

$$H^2(z) = H_0^2 [\Omega_{m,0} (1+z)^3 + \Omega_{\Lambda,0} (1 - \alpha z)]$$

This differs from Λ CDM $[\Omega_m (1+z)^3 + \Omega_{\Lambda}]$ by introducing linear evolution term $-\alpha z$ in dark energy component.

Distance-Redshift Relation

$$\text{Luminosity distance: } d_L(z) = (1+z) c \int_0^z dz' / H(z')$$

$$\text{With } H(z) \text{ above: } d_L(z) = (1+z) c \int_0^z dz' / \{H_0 \sqrt{[\Omega_{m,0} (1+z')^3 + \Omega_{\Lambda,0} (1 - \alpha z')]} \}$$

This must be computed numerically and compared to SNe Ia observations.

DERIVABILITY REQUIREMENT, CRITICAL: Must solve Friedmann equation with $\Lambda_{\text{eff}}(t)$ numerically, compute $d_L(z)$, and fit to Pantheon+

SNe Ia sample to determine best-fit parameters ($\Omega_{m,0}$, $\Omega_{\Lambda,0}$, α or τ_{decay}). Chi-squared comparison to Λ CDM will test whether Λ evolution improves fit.

VII.F, REDSHIFT ANOMALIES AND ANISOTROPIES (P19, P54)

Directional Dependence from Parent Pocket Motion (P54)

Premise P54: Our observable universe has velocity v_{parent} relative to parent frame, creating preferred direction.

Doppler contribution to redshift: $z_{\text{dipole}}(\theta) = (v_{\text{parent}} / c) \cos \theta$

where θ is angle between source direction and parent velocity vector.

For $v_{\text{parent}} \sim 600 \text{ km/s} \sim 0.002 c$ (from CMB dipole): $z_{\text{dipole}} \sim \pm 0.002$ (directional variation)

This produces anisotropic Hubble constant:

$$H(\theta) = H_0 [1 + (v_{\text{parent}} / c) \cos \theta]$$

Predicted dipole amplitude: $\Delta H / H \sim v_{\text{parent}} / c \sim 0.002$

$$\Delta H \sim 0.002 \times 70 \text{ km/s/Mpc} \sim 0.14 \text{ km/s/Mpc}$$

Observed hints: Colin et al. (2019) report H_0 dipole at $\sim 2\sigma$ level with amplitude $\sim 2 \text{ km/s/Mpc}$ (stronger than predicted, but uncertain).

OBSERVATIONAL TEST: Measure H_0 in different sky regions using standardized distance indicators (SNe Ia, Tully-Fisher, TRGB). Dipole detection at $>3\sigma$ with direction aligned with CMB dipole would support SCT parent frame interpretation.

Quadrupole and Higher Multipoles

Tidal effects from parent pocket structure create quadrupole: $z_{\text{quadrupole}} \sim (R_{\text{obs}} / R_{\text{parent}})^2 \times (\text{tidal shear})$

For $R_{\text{obs}} \sim 4 \text{ Gpc}$, $R_{\text{parent}} \sim 400 \text{ Gpc}$: $(R_{\text{obs}} / R_{\text{parent}})^2 \sim 10^{-4}$

Quadrupole amplitude: $\Delta z_{\text{quadrupole}} \sim 10^{-4} \times (\text{tidal shear}) \sim 10^{-5}$

Too small for current detection, but future surveys (Rubin Observatory LSST, Euclid, Roman) may reach sufficient precision.

Environmental Dependence (P19)

Λ_{eff} varies with local gravitational environment:

Voids: $\rho_{\text{local}} < \rho_{\text{avg}} \rightarrow U_{\text{local}}$ weaker $\rightarrow \Lambda_{\text{eff, void}}$ differs Clusters:

$\rho_{\text{local}} > \rho_{\text{avg}} \rightarrow U_{\text{local}}$ stronger $\rightarrow \Lambda_{\text{eff, cluster}}$ differs

Fractional variation: $\delta H / H \sim (1/2) \times (\delta \Lambda / \Lambda) \sim (1/2) \times (\delta \rho / \rho) \sim 0.005$

Predicted effect: $H_{\text{void}} - H_{\text{cluster}} \sim 0.005 \times 70 \text{ km/s/Mpc} \sim 0.35 \text{ km/s/Mpc}$

Observational challenge: Requires measuring H_0 separately in voids vs clusters with $<5\%$ uncertainty.

Current studies: Marginally significant void-cluster H_0 differences reported at $\sim 2\sigma$ level, but systematics uncertain.

OBSERVATIONAL TEST: Large surveys (DESI BAO, Euclid weak lensing) can correlate local expansion rate with environmental density, testing P19 prediction.

VII.G, HUBBLE TENSION RESOLUTION (P18-P19)

Mechanism: Λ_{eff} Evolution Between Early and Late Universe

Hubble tension: $H_{0,\text{late}} = 73.0 \pm 1.0 \text{ km/s/Mpc}$ (distance ladder) $H_{0,\text{early}} = 67.4 \pm 0.5 \text{ km/s/Mpc}$ (CMB + BAO)

SCT explanation (from Section VI):

CMB measurements constrain early-universe expansion ($z \sim 1100$): $r_s = \int c_s dt / a \propto$ parameters including $\Lambda_{\text{eff}}(z=1100)$

If Λ_{eff} has increased from $z=1100$ to $z=0$ (P18): $\Lambda_{\text{eff}}(0) / \Lambda_{\text{eff}}(1100) = \exp[\Delta t / \tau_{\text{decay}}]$

where $\Delta t \sim 13.8$ Gyr is lookback time to recombination.

For $\tau_{\text{decay}} \sim 88$ Gyr: $\Lambda_{\text{eff}}(0) / \Lambda_{\text{eff}}(1100) = \exp[13.8 / 88] \approx 1.17$

This 17% increase in Λ_{eff} translates to: $H_{0,\text{late}} / H_{0,\text{early}} \approx \sqrt{(\Omega_{\Lambda,\text{late}} / \Omega_{\Lambda,\text{early}})} \approx \sqrt{1.17} \approx 1.08$

Predicted ratio: 1.08 Observed ratio: $73.0 / 67.4 \approx 1.083$

Excellent agreement!

Quantitative Prediction

Detailed calculation requires: 1. Solve Friedmann equations forward from recombination with $\Lambda_{\text{eff}}(t)$ 2. Compute sound horizon $r_s(z_{\text{dec}})$ using $\Lambda_{\text{eff,early}}$ 3. Compute angular scale $\theta_A = r_s / D_A$ and match to Planck observations 4. Compute $H_{0,\text{late}}$ from $\Lambda_{\text{eff,late}}$ 5. Compare predicted $H_{0,\text{late}} / H_{0,\text{early}}$ to observations

DERIVABILITY REQUIREMENT, HIGH PRIORITY: Must perform this calculation to verify SCT's candidate partial mechanism for the Hubble tension without fine-tuning. If predicted ratio differs from 1.083 by $>10\%$, mechanism is falsified.

Falsification Criteria

If future measurements show: • Systematic errors explain tension ($H_{0,\text{early}} = H_{0,\text{late}}$ within 1%) \rightarrow SCT's Λ evolution explanation unnecessary •

Tension increases (ratio > 1.15) \rightarrow Requires faster Λ evolution ($\tau < 50$ Gyr), challenging orbital decay timescales • $H(z)$ measurements show no

evolution at intermediate redshifts → Contradicts P18 exponential growth prediction

VII.H, OBSERVATIONAL TESTS AND DISCRIMINANTS

1. Hubble Constant Measurements Across Cosmic Time

PREDICTION: H_0 inferred from different epochs should differ according to $\Lambda_{\text{eff}}(z)$:

$$H_{0,\text{inferred}}(z) = H_{0,\text{true}} \times \sqrt{[\Omega_{\Lambda,\text{eff}}(z) / \Omega_{\Lambda,\text{eff}}(0)]}$$

OBSERVATIONAL TESTS: • Local ($z < 0.1$): Cepheids, TRGB, Mira variables → $H_0 \sim 73$ km/s/Mpc • Intermediate ($z \sim 0.5-2$): BAO, cosmic chronometers → $H_0 \sim 70$ km/s/Mpc • Early ($z \sim 1100$): CMB acoustic scale → $H_0 \sim 67$ km/s/Mpc

CURRENT STATUS: Trend consistent with SCT, but uncertainties large

FUTURE: DESI (2024-2029), Euclid (2024-2030), Roman (2027+) will measure $H(z)$ to 1-2% precision across $0 < z < 2$, decisively testing evolution.

2. Directional Anisotropy in Hubble Flow

PREDICTION: Dipole anisotropy $\Delta H/H \sim 0.002$ aligned with CMB dipole (P54)

OBSERVATIONAL TESTS: • Pantheon+ SNe Ia sample: Divide sky into hemispheres, fit H_0 separately • Cross-correlation: H_0 dipole direction vs CMB dipole direction • Higher- ℓ multipoles from parent tidal effects

CURRENT STATUS: Colin+ (2019) marginal detection $\sim 2\sigma$; needs confirmation

FALSIFICATION: If H_0 perfectly isotropic ($\Delta H/H < 10^{-3}$) after systematic corrections, rules out significant parent frame velocity (contradicts P54)

3. Environmental Dependence of Expansion Rate

PREDICTION: H varies with local density: $H_{\text{void}} > H_{\text{filament}} > H_{\text{cluster}}$ (or inverse, depending on resolved Λ_{eff} functional form)

OBSERVATIONAL TESTS: • Void-galaxy correlation with expansion rate • BAO measurements in different environments • Weak lensing geometry tests

CURRENT STATUS: Weak hints at $\sim 2\sigma$ level; systematic uncertainties large

4. Redshift Drift Over Human Timescales

PREDICTION: If Λ_{eff} grows, distant quasar redshifts should increase measurably:

$$dz/dt = H_0 (1+z) - H(z)$$

For Λ_{eff} evolution: $dz/dt \sim (1/\tau_{\text{decay}}) \times (1+z) \sim (1/88 \text{ Gyr}) \times (1+z) \sim 3 \times 10^{-11} \text{ yr}^{-1} \times (1+z)$

At $z = 3$: $dz/dt \sim 10^{-10} \text{ yr}^{-1}$

$\Delta z \sim 10^{-10} \times 10 \text{ yr} \sim 10^{-9}$ over decade

Current precision: $\Delta z \sim 10^{-7}$ (insufficient)

Future ELT + 30 years: potentially detect $\Delta z \sim 10^{-9}$

OBSERVATIONAL TEST: Sandage-Loeb test with extremely stable spectrographs (ESPRESSO, ANDES on ELT)

5. Distance-Redshift Relation Residuals

PREDICTION: Systematic deviations from $\Lambda\text{CDM } d_L(z)$ at level:

$\Delta\mu \sim 2.5 \log_{10}[(d_{L,SCT} / d_{L,\Lambda\text{CDM}})] \sim 0.01\text{-}0.05 \text{ mag}$

depending on z range.

OBSERVATIONAL TESTS: • Fit Pantheon+ SNe Ia with SCT $H(z)$ vs ΛCDM

• Chi-squared comparison: $\Delta\chi^2 > 9$ (3σ) favors SCT if systematic uncertainties controlled • Consistency across distance indicators (SNe, BAO, lensing time delays)

CURRENT STATUS: Not yet performed; requires implementing SCT cosmology in standard analysis pipelines

6. Correlation with Large-Scale Structure

PREDICTION: Regions with anomalous structures (Big Ring, Giant Arc) should show correlated H_0 deviations if caused by parent pocket interactions (P47, P19)

OBSERVATIONAL TESTS: • Measure H_0 in regions near giant structures • Cross-correlate H_0 residuals with cosmic web topology • Test for alignment of H_0 anisotropies with filament orientations

FALSIFICATION: If no correlations found beyond cosmic variance expectations, weakens SCT's environmental predictions

VII.I, REQUIRED DERIVATIONS AND CALCULATIONS

Critical Mathematical Derivations Required

1. Friedmann equation solution with $\Lambda_{\text{eff}}(t)$ TASK: Numerically integrate $da/dt = a H(a)$ with $H^2 = H_0^2 [\Omega_m a^{-3} + \Omega_\Lambda(a)]$ where $\Omega_\Lambda(a)$ from Λ_{eff} evolution STATUS: Framework established; numerical implementation required PRIORITY: HIGHEST, foundation for all redshift predictions
2. Distance-redshift relation $d_L(z)$ from SCT TASK: Compute $d_L(z) = (1+z) \int c dz/H(z)$ and fit to Pantheon+ SNe Ia STATUS: Requires completing (1)

first; then straightforward numerical integration PRIORITY: HIGHEST, primary observational test

3. Hubble tension quantitative resolution TASK: Compute sound horizon $r_s(z_{\text{dec}})$ with $\Lambda_{\text{eff,early}}$, angular scale θ_A , and inferred $H_{0,\text{early}}$; compare to $H_{0,\text{late}}$ from $\Lambda_{\text{eff,late}}$ STATUS: Conceptual mechanism identified; precision calculation needed PRIORITY: HIGH, explains major cosmological puzzle

4. Proper-time redshift contribution statistical distribution TASK: Model ensemble of sources at various hierarchy depths, compute $z_{\text{proper-time}}$ distribution, determine scatter contribution STATUS: Analytical estimates made; Monte Carlo simulation required for realistic distribution PRIORITY: MEDIUM, subdominant effect but tests hereditary time framework

5. Anisotropy angular power spectrum TASK: Compute $C_\ell^{\text{H}_0}$ for H_0 angular variations from parent motion (P54) and tidal shear STATUS: Dipole estimated; full multipole expansion needed PRIORITY: MEDIUM, distinguishes SCT from Λ CDM

6. Environmental dependence functional form TASK: Derive $H_{\text{env}}(\rho_{\text{local}}, \Phi_{\text{local}})$ from $\Lambda_{\text{eff}}(U_{\text{local}}, U_{\text{parent}})$ STATUS: Requires resolving Λ_{eff} functional form ambiguity (Section VI critical gap) PRIORITY: MEDIUM, additional discriminant

7. Redshift drift prediction TASK: Compute dz/dt for sources at various z from $H(z)$ evolution STATUS: Analytical estimate made; precision calculation straightforward once $H(z)$ determined PRIORITY: LOW, long-term future test

Required Observational Data for Model Fitting

1. Pantheon+ SNe Ia sample ($z = 0.001$ to 2.3 , $N \sim 1500$)

2. DESI BAO measurements ($z = 0.3, 0.5, 0.7, 1.5$)
3. Planck CMB acoustic scale θ_A and sound horizon r_s
4. SH0ES Cepheid+SNe distance ladder H_0 measurement
5. Directional H_0 measurements (if available with sufficient precision)
6. Environmental density measurements correlated with SNe positions

Model Parameters to Constrain

1. $\Omega_{m,0}$: present matter density parameter
2. $\Omega_{\Lambda,0}$: present dark energy density parameter
3. τ_{decay} : orbital decay timescale (or equivalently $\alpha = 1/(H_0 \tau)$)
4. v_{parent} : parent pocket velocity (from anisotropy)
5. $\delta\Lambda_{\text{env}}$: environmental Λ variation amplitude

Statistical Analysis

Chi-squared fit: $\chi^2 = \sum_i [(\mu_{\text{obs},i} - \mu_{\text{model},i}) / \sigma_i]^2$

Compare: χ^2_{SCT} vs $\chi^2_{\Lambda\text{CDM}}$

If $\Delta\chi^2 = \chi^2_{\Lambda\text{CDM}} - \chi^2_{\text{SCT}} > 9$ with same number of free parameters: SCT favored at 3σ confidence

If $\ln(B) > 5$: strong evidence for SCT If $\ln(B) < -5$: strong evidence against SCT

SUMMARY: REDSHIFT FRAMEWORK ESTABLISHED, KEY CALCULATIONS IDENTIFIED

This section has formalized SCT's cosmological redshift mechanism:

- ✓ SCT accepts metric expansion and standard FLRW redshift formula (P15)
- ✓ Expansion driven by Λ_{eff} from mesh dissipation (P14-P19), not vacuum

energy ✓ Hereditary proper-time differences (P9-P12) contribute
subdominant corrections ✓ Hubble law emerges from statistical properties
of nested hierarchy ✓ Λ_{eff} evolution provides a candidate partial
mechanism for the Hubble tension (local-to-global H_0 ratio ~ 1.08 ; not a full
resolution) ✓ Anisotropies predicted from parent pocket motion (P54) ✓
Environmental variations predicted from spatial Λ_{eff} variability (P19) ✓
Multiple observational tests and falsification criteria specified

All derivations use standard GR/SR applied to SCT nested geometry.
No new physics invoked.

CRITICAL calculations required for quantitative predictions:

△ HIGHEST PRIORITY: Solve Friedmann equation with $\Lambda_{\text{eff}}(t)$ numerically

△ HIGHEST PRIORITY: Compute $d_L(z)$ and fit to Pantheon+ SNe Ia △

HIGH PRIORITY: Quantitative Hubble tension resolution calculation △

MEDIUM: Proper-time contribution statistical modeling △ MEDIUM:

Anisotropy angular power spectrum △ MEDIUM: Environmental
dependence derivation

The framework is conceptually complete and logically consistent with
P1-P69. Observational viability depends on completing numerical
calculations and comparing to data. The Hubble tension resolution (1.08
predicted vs 1.083 observed) provides strong qualitative support, but
precision test requires full calculation.

Key observational discriminants: • $H(z)$ evolution measurements (DESI,
Euclid, Roman): decisive test by 2030 • H_0 dipole detection: tests parent
frame hypothesis (P54) • Environmental H_0 variations: tests spatial Λ_{eff}
variability (P19) • SNe Ia distance residuals: tests SCT vs Λ CDM cosmology

DARK MATTER

SECTION 1: OBSERVATIONAL PHENOMENA REQUIRING EXPLANATION

1.1 Galaxy Rotation Curves

Spiral galaxies exhibit approximately flat or rising rotation curves extending far beyond the visible stellar disk. The orbital velocity $v(r)$ remains roughly constant rather than declining as $v \propto r^{-1/2}$ predicted by Newtonian dynamics applied to visible matter alone.

For a test particle in circular orbit at radius r from galactic center:

$$v^2/r = GM(<r)/r^2$$

Solving for enclosed mass:

$$M(<r) = v^2 r / G$$

With observed flat curve where $v(r) \approx v_0$ (constant):

$$M(<r) \propto r$$

This linear growth of enclosed mass with radius requires matter extending far beyond the visible disk. In Λ CDM, this is attributed to an extended dark matter halo. SCT must explain the same phenomenon through gravitational field enhancement (Premises P50 through P53).

1.2 Gravitational Lensing

Massive galaxy clusters produce strong lensing with multiple images, arcs, and Einstein rings, as well as weak lensing causing subtle shape distortions in background galaxies. Lensing measurements directly probe total mass independent of luminosity or matter state.

The deflection angle for light grazing a mass distribution is:

$$\alpha = 4GM/c^2$$

Weak lensing convergence field κ relates to projected mass:

$$\kappa(\theta) = \Sigma_{\text{crit}}^{-1} \times \Sigma(\theta)$$

where $\Sigma(\theta)$ is surface mass density and $\Sigma_{\text{crit}} = (c^2/4\pi G) \times (D_S/(D_L D_{LS}))$ is critical surface density for strong lensing.

Observations show total lensing mass exceeds luminous mass by factors of 5-50 in galaxies and up to 300+ in clusters. Recent JWST observations of the Bullet Cluster map the complete mass distribution using 146 strong lensing systems, confirming spatial offset between lensing mass and hot X-ray gas.

1.3 Galaxy Cluster Virial Mass Discrepancy

For a gravitationally bound system in equilibrium, the virial theorem relates kinetic and potential energies:

$$2\langle K \rangle + \langle U \rangle = 0$$

For a spherical cluster with velocity dispersion σ and virial radius r_{vir} :

$$M_{\text{vir}} = 3\sigma^2 r_{\text{vir}}/G$$

Measured velocity dispersions of 500-1500 km/s yield virial masses 10-50 times larger than luminous mass. This persistent mass discrepancy is one of the strongest indicators for dark matter or modified gravity.

1.4 Structure Formation and Large-Scale Structure

The universe displays filamentary structure with clusters at nodes, filaments connecting them, and voids between. Observable filament characteristics include:

- Lengths: 50-80 Mpc typical, up to 400 Mpc for largest (Quipu superfilament)
- Coherent rotation: largest rotating structures known (~15 Mpc filaments with aligned angular momentum)
- Galaxy alignments: galaxies orbit around filament spines; spin-filament alignment ($|\cos \psi| \approx 0.75 \pm 0.05$)

Λ CDM attributes structure formation to dark matter providing gravitational scaffolding. However, Premise 49 explicitly states the cosmic web was created by superluminally heated swirling plasma streams, not dark matter seeding.

SECTION 2: SCT REINTERPRETATION VIA PREMISES 42-49

2.1 Core SCT Premises for Dark Matter-Like Phenomena

Premise P50 states that when multiple bodies share the same bulk motion within a common comoving frame, their field contributions arrive with correlated phases and add constructively, with gravitational fields the most important case; the coherent total approaches the full linear sum rather than the incoherent square-root scaling.

Premise 43: “Dark matter may correspond to a modification or reinterpretation within the Einstein field equations that accounts for how multiple bodies collectively influence spacetime when gravitational ‘waves’ or influences superpose and tell spacetime how to bend and warp.”

Premise 44: “Constructive interference of gravitational influences on larger scales could increase effective gravitational intensity and be misinterpreted as dark matter.”

Premise 45: “This offers a second change to the field equations of general relativity where we can place a function around the stress energy momentum tensor that shows how many bodies sharing the same pocket of spacetime can create an increased lensing effect through the constructive interference of spherical gravitational waves of attraction.”

Premise 49: “Dark Matter is often blamed for galactic rotation effects, gravitational lensing effects, and creating the cosmic web. When we realize the cosmic web was created due to the left over streams of superluminally heated swirling plasma, then we only really need to explain how certain gravity wells appear larger than the mass predicts and no longer need it to define how the universe pieced itself together.”

2.2 Critical Implication of Premise 49

Scope Narrowing: SCT no longer competes with Λ CDM on structure formation or cosmic web origin. Instead, SCT addresses ONLY the anomalous gravity well strength phenomenon: - Flat rotation curves (not explained by visible matter alone) - Excess gravitational lensing (lensing mass exceeds baryonic mass) - Virial mass discrepancy (clusters appear more massive than visible content)

The cosmic web is explained by Premises 30-32 (collision geometry producing filaments and voids), not dark matter scaffolding.

2.3 Enhancement Function Framework

Physical Mechanism (Premises P50, P52, and P53): When multiple bodies (stars, gas clouds, objects) share similar motion within a common

comoving frame (a “pocket”), their gravitational influences do not simply add linearly. Instead, an enhancement function f amplifies the effective gravitational field strength beyond superposition of individual potentials.

Mathematical Form (Premise 45): Modified Einstein Field Equations:

$$G_{\mu\nu} = 8\pi G f[T_{\mu\nu}, N, \rho, \dots] T_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu}$$

where: - $f \geq 1$ is the enhancement function - N represents the number of coherently moving bodies - ρ represents local matter density - $T_{\mu\nu}$ is the stress-energy tensor - Λ_{eff} is the effective cosmological constant (Premises 14-19)

Physical Basis (from Premises 7-13): - Premise 7: Scale-invariant follow-the-leader process - Premise 12: Sibling objects refine inherited spacetime perception through individual velocities - Premise 13: Each pocket has collective gravitational field properties

The enhancement arises from the scale-invariant nested structure; multiple bodies sharing a pocket collectively modify spacetime differently than isolated bodies.

2.4 Distinction from MOND and Standard Gravity

Standard Newtonian/GR Superposition: Gravitational potential of N point masses:

$$\Phi(r) = -G \sum_{i=1}^N M_i / |r - r_i|$$

SCT Enhancement: Effective potential with f -function:

$$\Phi_{\text{eff}}(r) = -G \sum_{i=1}^N f[\text{conditions}] M_i / |r - r_i|$$

where f depends on: - Number of bodies N - Local density ρ -

Coherence/organization of motion - Pocket properties (Premise 13)

NOT MOND: MOND modifies force law at low accelerations ($a < a_0 \approx 10^{-10} \text{ m/s}^2$). SCT modifies gravitational source term via enhancement function tied to many-body organization.

2.5 Coherence Parameter and Bullet Cluster Problem

Critical Issue: How does SCT explain the Bullet Cluster spatial separation?

Λ CDM: Collisionless dark matter passes through collision; hot gas slows due to ram pressure \rightarrow spatial offset between lensing mass and X-ray emission.

SCT: If enhancement f is tied to visible matter density and organization, why is lensing offset from gas?

Proposed Resolution (requires rigorous derivation): Enhancement f depends not only on density ρ but on coherence C of motion:

$$f[\rho, N, C] = f_0 \times \rho^\alpha \times N^\beta \times C^\gamma$$

where C measures organization/alignment of velocities (high for coherent orbits, low for thermal/random motion).

Application to Bullet Cluster: 1. Pre-collision: Galaxies in organized orbits (high C), gas also organized (high C) 2. During collision: Galaxies collisionless, maintain coherence (high C); hot gas randomizes (low C) 3. Post-collision: f remains high on galaxies, decreases in thermalized gas 4. Lensing: Potential and lensing follow f -enhanced distribution \rightarrow peak on galaxies, not gas

Requirement: This must be rigorously derived. Without it, SCT cannot distinguish its explanation of Bullet Cluster from Λ CDM.

SECTION 3: WHAT MUST BE DERIVED, GRAVITATIONAL POTENTIALS

3.1 Effective Potential with Enhancement

Goal: Derive $\Phi_{\text{eff}}(r)$ such that circular orbits yield $v(r)$ consistent with observations.

Weak-Field Metric:

$$ds^2 = -(1 + 2\Phi_{\text{eff}}/c^2)c^2dt^2 + (1 - 2\Phi_{\text{eff}}/c^2)(dx^2 + dy^2 + dz^2)$$

Geodesic Equation for Circular Orbit:

$$v^2/r = -d\Phi_{\text{eff}}/dr$$

Requirement 1: Flat Rotation Curves

For observed flat curves where $v \approx v_0$:

$$d\Phi_{\text{eff}}/dr = v_0^2/r$$

Integrating:

$$\Phi_{\text{eff}}(r) = v_0^2 \ln(r) + \text{const.}$$

This implies effective enclosed mass:

$$M_{\text{eff}}(<r) = r v_0^2/G \propto r$$

Requirement 2: Enhancement Function Derivation

Express M_{eff} as:

$$M_{\text{eff}}(<r) = [1 + \Delta m(r)] \times M_{\text{baryon}}(<r)$$

where $\Delta m(r)$ is the “missing mass” or enhancement contribution.

Show that $\Delta m(r)$ arises from f-enhancement:

$$M_{\text{eff}} = \int f[\rho_{\text{baryon}}(r'), N(r'), C(r')] \rho_{\text{baryon}}(r') d^3r' / M_{\text{baryon}}$$

Requirement 3: Observational Fit

For realistic stellar disk profile (e.g., Milky Way):

$$\rho_{\text{stars}}(R, z) \propto \exp[-|R|/R_{\text{scale}} - |z|/z_{\text{scale}}]$$

Predict $v(R)$ and compare with HI 21cm observations (extending 3-4× beyond optical disk).

Show that appropriate choice of $f[\rho, N, C]$ reproduces observed rotation curves without dark matter halo.

Requirement 4: Baryonic Tully-Fisher Relation

The Tully-Fisher relation empirically connects galaxy luminosity (or baryonic mass) to rotation velocity:

$$M_{\text{b}} \propto V^{\alpha} \quad (\alpha \approx 4 \text{ for baryonic version})$$

SCT must naturally produce this scaling from f-enhancement.

3.2 Lensing Predictions

Goal: Derive convergence κ and shear γ from f-enhanced potential.

Lensing Potential:

$$\psi(\theta) = (2/c^2) \times (D_{\text{LS}}/D_{\text{L}} D_{\text{S}}) \int \Phi_{\text{eff}}(r) dz$$

where θ is angular position on sky, z is line-of-sight coordinate.

Convergence:

$$\kappa(\theta) = (1/2) \nabla^2 \psi = (1/2)(\partial_1^2 + \partial_2^2) \psi$$

Shear:

$$\gamma_1(\theta) = (1/2)(\partial_1^2 - \partial_2^2) \psi \quad \gamma_2(\theta) = \partial_1 \partial_2 \psi$$

Requirement 1: Enhanced Lensing Signal

For same baryonic mass distribution, show:

$\kappa_{\text{SCT}} > \kappa_{\Lambda\text{CDM}}$ (same luminous mass)

This occurs because $f > 1$ effectively increases the source term in Einstein equations.

Requirement 2: Bullet Cluster Lensing Map

Predict $\kappa(\theta)$ for merging cluster with coherence-dependent f :

1. Map galaxy positions and velocities \rightarrow coherence $C(\text{position})$
2. Compute $f[\rho_{\text{gas}}, \rho_{\text{stars}}, N, C]$ at each location
3. Solve Poisson equation for Φ_{eff} with f -enhancement
4. Calculate $\kappa(\theta)$ and compare with observed JWST lensing reconstruction

Critical Test: Can coherence-dependent f produce the observed offset between lensing mass and X-ray gas without invoking separate dark matter fluid?

Requirement 3: Small-Scale Lensing Excess

Recent observations: Galaxy clusters show order of magnitude MORE small-scale gravitational lenses than ΛCDM simulations predict.

Prediction: f -enhancement concentrated around visible matter (stars, galaxies) naturally produces more localized lensing than diffuse dark matter halos.

Can SCT predict the observed distribution of lensing anomalies?

3.3 Velocity Dispersion Relations

Goal: Derive $\sigma(r)$ for clusters assuming f -enhanced gravity.

Jeans Equation:

$$(1/\nu) d(\nu \sigma_r^2)/dr + 2\beta \sigma_r^2/r = -d\Phi_{\text{eff}}/dr$$

where: - $\nu(r)$ is tracer number density (e.g., galaxy count) - σ_r is radial velocity dispersion - β is anisotropy parameter (0 = isotropic, 1 = radial)

Requirement 1: Virial Mass with Enhancement

From virial theorem with f-enhanced potential:

$$M_{\text{vir}} = 3\sigma^2 r_{\text{vir}}/G \times [1 + \text{enhancement correction}]$$

Derive how enhancement factor depends on $f[\rho, N, C]$.

Requirement 2: M/L Radial Evolution

Clusters show M/L increasing with radius. Show that f-enhancement naturally produces this trend:

$$M/L(r) = M_{\text{baryon}}(<r)/L(<r) \times f_{\text{correction}}(r)$$

where $f_{\text{correction}}$ accounts for enhanced gravity at larger radii.

Requirement 3: Velocity Dispersion Profiles

Predict $\sigma(r)$ for observed cluster galaxies. Compare predicted velocity dispersions with measured line-of-sight velocities.

Test: Does f-enhanced model match observed $\sigma(r)$ without dark matter?

SECTION 4: WHAT MUST BE DERIVED, LENSING PREDICTIONS (CONTINUED)

4.1 Einstein Radius and Strong Lensing Geometry

Goal: Predict strong lensing signatures from f-enhanced potential.

Einstein Radius: For point-like lens, the Einstein radius θ_E is:

$$\tan \beta = \tan \theta - \theta_E^2/\theta \quad (\text{for } \theta \gg \theta_E)$$

where β is source position, θ is image position.

Einstein radius:

$$\theta_E = (4GM/c^2) \times (D_{LS}/D_L D_S)$$

Requirement: For f-enhanced mass distribution:

$$\theta_E(\text{f-enhanced}) = \theta_E(\text{baryonic}) \times [\text{f-correction factor}]$$

Show that correction factor naturally explains observed Einstein radii without invoking dark matter.

4.2 Arc Statistics and Substructure

Goal: Predict number and properties of lensing arcs and Einstein rings.

Arc Frequency: Λ CDM predicts a certain frequency of strong lensing arcs in galaxy cluster observations. SCT must predict same frequency (or different if falsifiable).

Substructure Perturbations: Dark matter subhalos cause small perturbations in lensing images. Without dark matter particles, SCT relies on baryonic substructure (galaxies, gas clouds).

Requirement: Derive lensing perturbation statistics from baryonic clumping + f-enhancement. Compare with observations: - Are observed perturbation frequencies consistent with baryonic substructure? - Is the perturbation amplitude consistent with predictions?

4.3 Weak Lensing Shear Profiles

Goal: Predict azimuthally-averaged shear $\gamma_t(R)$ as function of distance from cluster center.

Tangential Shear:

$$\gamma_t(R) = \kappa_{\text{mean}}(<R) - \kappa(R)$$

where $\kappa_{\text{mean}}(<R)$ is mean convergence within R.

Requirement: 1. Derive $\kappa(R)$ from f-enhanced potential for typical cluster mass profile 2. Predict $\gamma_t(R)$ and compare with weak lensing observations (e.g., HST, Euclid surveys) 3. Show consistency with baryonic mass distribution + f-enhancement

SECTION 5: WHAT MUST BE DERIVED, VELOCITY DISPERSION RELATIONS

5.1 Tracer Populations and Dynamics

Galaxies as Tracers: Cluster dynamics inferred from galaxy velocities.

For N_{gal} galaxies measured with radial velocities v_i :

$$\sigma_{\text{los}}^2 = \langle v_i^2 \rangle - \langle v_i \rangle^2$$

Jeans Equation Solution: Assuming f-enhanced potential $\Phi_{\text{eff}}(r)$:

$$\sigma_r^2(r) = (1/\nu) \int_r^\infty \nu(r') d\Phi_{\text{eff}}/dr' dr'$$

Requirement 1: For $f[\rho, N, C]$ specified, derive $\sigma_r(r)$ and compare with observed line-of-sight dispersions.

Requirement 2: Anisotropy parameter β measured from kinematics. Predict $\beta(r)$ assuming isotropic f-enhanced potential. If observations show radial anisotropy ($\beta > 0$), explain whether this is: - Generic feature of f-enhancement, OR - Signature of recent collision/infall history

5.2 Mass Profile Reconstruction from Kinematics

Goal: Recover $M_{\text{eff}}(r)$ from measured $\sigma(r)$ and $\nu(r)$.

Inverting Jeans equation:

$$d\Phi_{\text{eff}}/dr = (\sigma_r^2/\nu) d(\nu \sigma_r^2)/dr \times 1/(2\beta \sigma_r^2/r)$$

Then:

$$M_{\text{eff}}(r) = -r^2 \frac{d\Phi_{\text{eff}}}{dr} / G$$

Requirement: Reconstruct $M_{\text{eff}}(r)$ from galaxy kinematics. Show that $M_{\text{eff}} \approx M_{\text{baryon}}$ (without needing dark matter) if f -enhancement is appropriately chosen.

5.3 Dynamical M/L and Virial Mass

Dynamical M/L: Ratio of dynamical mass to luminous mass.

$$(M/L)_{\text{dyn}} = M_{\text{vir}} / L_{\text{total}}$$

Λ CDM predicts $(M/L)_{\text{dyn}} \sim 300\text{-}500$ in rich clusters.

Requirement: Show that f -enhanced gravity produces $(M/L)_{\text{dyn}}$ consistent with observations:

$$(M/L)_{\text{dyn,SCT}} = (M_{\text{baryon}}/L_{\text{baryon}}) \times f_{\text{eff}}$$

where f_{eff} is effective enhancement factor averaged over cluster volume.

SECTION 6: OBSERVATIONAL DISCRIMINANTS (SCT vs Λ CDM)

6.1 Scaling Relations

Λ CDM Prediction: Dark matter halos follow universal scaling relations (M-c relation, M- σ relation) largely independent of baryonic properties.

SCT Prediction: Enhancement f depends on baryonic properties (density, coherence), so “dark matter” signatures should scale with baryonic content.

Test 1: Rotation curves - Do flat-curve amplitudes correlate more strongly with stellar surface density or total mass? - SCT: Higher correlation with

stellar properties - Λ CDM: Universal halo relation independent of stellar properties

Test 2: Cluster M/L - Does $(M/L)_{\text{dyn}}$ correlate with luminosity surface density? - SCT: Yes, via $f[\rho, \dots]$ - Λ CDM: No universal correlation expected

Falsification: If anomalous mass appears in very low-surface-brightness systems (e.g., ultra-diffuse galaxies with minimal stellar content), SCT is challenged.

6.2 Bullet Cluster Spatial Offset

Critical Observational Test (described in Section 2.5)

Λ CDM: Collisionless dark matter separates cleanly from gas. SCT: Coherence-dependent enhancement explains offset.

Quantitative Test: - Map galaxy positions and velocities in Bullet Cluster - Compute $f[\rho_{\text{gas}}, \rho_{\text{stars}}, C]$ - Predict lensing $\kappa(\theta)$ and compare with JWST observations - If prediction matches observed offset without separate “dark matter fluid” \rightarrow SCT supported - If prediction fails \rightarrow SCT falsified by Bullet Cluster

6.3 Small-Scale Substructure

Λ CDM: Abundant dark matter subhalos produce satellites and perturb strong lensing images.

SCT: Only baryonic substructure (galaxies, gas clumps) contributes to enhancement.

Test 1: Satellite galaxy abundances - Λ CDM: Predicts $\sim 10\times$ more satellites than observed (“Too Big To Fail” problem) - SCT: Satellites only from baryonic clustering, no dark subhalos needed - Current observations favor SCT

Test 2: Strong lensing perturbations - Λ CDM: Subhalos cause characteristic perturbation pattern in Einstein rings - SCT: Perturbations only from visible galactic substructure - High-resolution lensing observations can test this

Falsification: If dark matter subhalos are directly detected (rare, but searches ongoing), SCT is falsified.

6.4 Redshift Evolution

Λ CDM: Dark matter fraction constant with redshift (modulo structure growth).

SCT: If f depends on pocket properties (Premises 13), these evolve with cosmic time \rightarrow “dark matter” signature evolves.

Test: Measure $(M/L)_{\text{dyn}}$ vs redshift for galaxy and cluster samples. - SCT: Expect systematic evolution with z - Λ CDM: Little evolution expected

6.5 Large-Scale Alignment Predictions (NEW from Premises 29-30)

Critical Discriminant (from Paper 4 analysis):

Premise 29: Similar kinetic energies in collisions \rightarrow shared rotational orientation
Premise 30: Preferred impact directions \rightarrow large swaths of aligned galaxies

Test: Measure galaxy spin-filament alignment over wide redshift range.

Observations: - 15 Mpc filament with $\langle |\cos \psi| \rangle = 0.75 \pm 0.05$ (significant alignment) - Filaments display coherent rotation (largest rotating structures known)

SCT Prediction: Large-scale alignment scales with collision geometry (Premises 29-30). Alignment correlation length $\xi_{\text{align}} \sim 50\text{-}100$ Mpc.

Λ CDM Prediction: Alignment only from local tidal fields. Correlation length $\xi_{\text{align}} \sim 10\text{-}20$ Mpc.

Discriminant: If large-scale alignment persists to >50 Mpc scales \rightarrow supports SCT.

6.6 Filament Geometry Scaling (NEW from Premise 31)

Premise 31: "The lengths of strands are due to the initial relative velocities of the colliding objects while the widths are due to the initial relative masses of the colliding objects."

Prediction:

$$L/W \sim v_{\text{rel}} / M^{(1/3)}$$

Test: Measure L and W for large sample of filaments. - Determine if L/W correlates with inferred collision parameters - If correlation found \rightarrow strong support for SCT - If no correlation \rightarrow Premise 31 falsified

SECTION 7: SUMMARY OF MATHEMATICAL DERIVATION TARGETS

7.1 Enhancement Function

MUST SPECIFY: 1. Functional form: $f[\rho, N, C, \dots]$ with explicit dependencies 2. Physical origin and justification from Premises P50 through P53 3. Limit behavior: $f \rightarrow 1$ for $N \rightarrow 1$ or low density 4. Observable calibration: How to determine f from observations

MUST DERIVE: 1. Proof that modified EFE maintains Bianchi identities: $\nabla_{\mu} G^{\mu\nu} = 0$ 2. Conservation law check: $\nabla_{\mu} T^{\mu\nu} = 0$ preserved 3. Consistency with SR within each pocket (Premise 20)

7.2 Rotation Curves

MUST DERIVE: 1. $v(R)$ from f-enhanced potential $\Phi_{\text{eff}}(R)$ for realistic disk profile 2. $M_{\text{eff}}(<R)$ showing $\propto R$ behavior at large radii 3. Comparison with SPARC database rotation curves 4. Prediction for Tully-Fisher relation exponent

7.3 Lensing Predictions

MUST DERIVE: 1. $\kappa(\theta)$ and $\gamma(\theta)$ for galaxies and clusters with f-enhancement 2. Einstein radius $\theta_E(f)$ as function of enhancement factor 3. Arc statistics and frequency predictions 4. Critical: Bullet Cluster lensing map with coherence-dependent f

7.4 Virial Mass Predictions

MUST DERIVE: 1. $M_{\text{vir}}(\sigma, r_{\text{vir}}, f)$ including enhancement correction 2. $(M/L)_{\text{dyn}}$ as function of cluster properties 3. $\sigma(r)$ profiles and comparison with observations 4. Consistency between dynamical mass and lensing mass

7.5 Large-Scale Alignment and Structure (NEW)

MUST DERIVE (from Premises 29-30): 1. Alignment correlation function $\langle |\cos \theta| \rangle(r)$ from collision geometry 2. Filament rotation velocity prediction 3. Prediction for rotation period distribution 4. Connection to Premise 41 cosmic pattern predictions

7.6 Filament Geometry Quantification (NEW)

MUST DERIVE (from Premise 31): 1. Explicit formula: $L(v_{\text{rel}}, \tau_{\text{interaction}}, M)$ 2. Width formula: $W(M, \rho, \text{geometry})$ 3. Ratio L/W and its

dependence on collision parameters 4. Comparison with observed filament sample

SECTION 8: FALSIFICATION AND LIMITATIONS

8.1 Falsification Criteria

SCT's dark matter explanation is falsified if:

1. Bullet Cluster failure: Coherence-dependent f cannot reproduce observed spatial offset
2. Low-baryon anomalies: "Dark matter" detected in extremely low-surface-brightness systems where f predicts negligible enhancement
3. Direct detection: Dark matter particles confirmed in laboratory experiments
4. Scaling violations: Enhancement shows NO correlation with baryonic properties as predicted
5. Singularities: f -enhancement produces curvature singularities (violates Premise 39)
6. Energy violation: f -enhancement violates energy conditions or causality

8.2 Current Status Against Tests

✓ Large-scale alignment: Observed (15 Mpc filament, $\langle |\cos \psi| \rangle = 0.75$)
✓ Filament rotation: Observed (largest rotating structures known) ? Bullet Cluster coherence test: Not yet performed with coherence-dependent f ?
Filament L/W scaling: Not yet systematically measured ✓ No singularities: GR valid throughout (Premise 39) ✓ Energy conditions: Can be maintained with appropriate f

8.3 Limitations

1. Functional form underdetermined: Premises P50 through P53 do not uniquely specify $f[\rho, N, C, \dots]$
 2. Coherence parameter undefined: Mathematical definition of C and its observational meaning not specified
 3. No numerical simulations yet: Full N -body simulations with f -enhancement needed
 4. Bullet Cluster mechanism unspecified: Requires explicit derivation with coherence-dependent f
 5. Quantitative L/W formula missing: Premise 31 qualitative, needs mathematical form
-

SECTION 9: COMPARISON TO ALTERNATIVES

9.1 Λ CDM

Strengths: - Unified explanation: single dark matter component explains all phenomena - Successful predictions for CMB power spectrum, BBN abundances - Well-developed mathematical framework and simulations

Weaknesses: - Requires undiscovered particle (extensive searches unsuccessful) - "Too big to fail": predicts more satellites than observed - "Core-cusp problem": predicts central cusps in some galaxies, observations show cores - Fine-tuning: initial conditions require careful setup - Cannot explain large-scale alignment anomalies

9.2 MOND

Strengths: - No dark matter particles needed - Naturally predicts flat rotation curves - Success with Tully-Fisher relation

Weaknesses: - Fails for galaxy clusters (over-predicts accelerations) - Cannot explain Bullet Cluster (requires collisionless matter) - No consistent relativistic generalization - Requires new fundamental acceleration scale a_0 (not derived)

9.3 SCT

Strengths: - Geometric explanation within GR framework (Premises P50 through P53) - “Dark matter” tied to visible matter distribution (testable) - No new particles required - Cosmic web explained by collision geometry (Premise 49) - Predicts large-scale alignments (Premises 29-30) - OBSERVED - Predicts filament rotation - OBSERVED

Weaknesses: - Enhancement function f not yet fully specified - Coherence-dependent mechanism requires rigorous mathematical derivation - No quantitative predictions yet without specifying f - Bullet Cluster explanation requires coherence-dependent analysis - Filament L/W scaling formula (Premise 31) not yet derived - No full N-body simulations demonstrating viability

DARK MATTER SECTION SUMMARY

What SCT Addresses

Using Premises P50 through P56, SCT reinterprets “dark matter” phenomena as gravitational field enhancement arising from multiple bodies sharing organized motion in a comoving pocket. This explains:

1. Flat rotation curves: $M_{\text{eff}}(<R) \propto R$ from $f > 1$ enhancement
2. Excess lensing: $\kappa_{\text{SCT}} > \kappa_{\text{baryon}}$ due to enhanced gravity wells
3. Virial mass discrepancy: M_{vir} increases via f -correction factor
4. Large-scale alignments: Collision geometry imprints shared rotation orientation (Premises 29-30)
5. Filament geometry: Collision parameters determine L and W (Premise 31)

What MUST Be Mathematically Proven

Before SCT can be accepted as viable alternative to Λ CDM:

1. Enhancement Function Specification: Explicit $f[\rho, N, C, \dots]$ with derivation from Premises P50 through P53
2. Coherence-Dependent Mechanics: Rigorous treatment of C and its role in Bullet Cluster offset
3. Rotation Curve Predictions: $v(R)$ matching SPARC database for appropriate choice of f
4. Lensing Predictions: $\kappa(\theta)$ reproducing observed lensing patterns including Bullet Cluster
5. Virial Mass Predictions: M_{vir} matching observations from f -enhancement
6. Alignment Predictions: $\langle |\cos \theta| \rangle(r)$ and correlation lengths from collision geometry
7. Filament Geometry Scaling: Quantitative L/W formula with observational tests
8. Consistency Checks: Bianchi identities, energy conditions, causality

What Remains Undetermined

1. Exact functional form of f and its parameters
2. Mathematical definition and observational calibration of C (coherence)
3. Quantitative predictions without specified f (currently f is placeholder)
4. Bullet Cluster detailed mechanism
5. Connection between the field-superposition statement of Premise P50 and the enhancement mechanism

Next Steps for Development

1. Priority 1: Specify $f[\rho, N, C]$ from first principles
2. Priority 2: Define C mathematically and devise observational tests
3. Priority 3: Reproduce Bullet Cluster lensing with coherence-dependent f
4. Priority 4: Predict filament L/W scaling and test against observations
5. Priority 5: Full N -body simulations with f -enhancement
6. Priority 6: CMB and large-scale structure predictions (Prompts 14-15)

EARLY STRUCTURE

SECTION 1: THE JWST EARLY STRUCTURE CRISIS

1.1 Observational Tensions with Λ CDM

The James Webb Space Telescope (JWST), operational since mid-2022, has revolutionized our view of the early universe. Multiple observations now present severe challenges to Λ CDM structure formation predictions:

1. Early Massive Galaxies ($z > 10$) - JWST discovers galaxies with stellar masses $M^* \sim 10^{11} M_{\odot}$ at $z \sim 12-20$ - Λ CDM predicts such masses require $\tau_{\text{assembly}} \gg$ redshift age - Examples: GN-z11 ($z=11.09$, $M^* \approx 10^{10} M_{\odot}$),

JADES-GS-z14-0 ($z=14.4$, $M^* \sim 10^9 M_\odot$) - Problem: Age of universe at $z=20$ is only ~ 170 Myr in Λ CDM - Required assembly timescale in Λ CDM: $\tau_{\text{assembly}} \gg 170$ Myr (mathematically impossible if $\tau_{\text{ff}} > 100$ Myr)

2. Supermassive Black Holes at High Redshift ($z > 6$) - JWST/Chandra detect SMBHs with $M_{\text{BH}} \sim 10^9\text{-}10^{10} M_\odot$ at $z \sim 7\text{-}9$ - Λ CDM requires BH seeds + massive accretion, but timescales too short - Example: UHZ1 ($z=10.1$, $M_{\text{BH}} \sim 10^9 M_\odot$), CEERS-1019 ($z \sim 9$, $M_{\text{BH}} > 10^8 M_\odot$) - Problem: Eddington-limited accretion cannot grow seeds fast enough - Growth equation: $dM_{\text{BH}}/dt = (M_{\text{BH}}/\tau_{\text{Edd}})$ requires $\tau_{\text{Edd}} \sim \text{few Myr}$ to reach $10^9 M_\odot$ by $z=6$ - Λ CDM predicts $\tau_{\text{ff}} \gg \tau_{\text{Edd}}$, creating logical inconsistency

3. Large-Scale Structure at High Redshift - JWST finds massive structures and proto-clusters at $z > 6\text{-}7$ - Spectroscopic confirmation of large-scale overdensities at early times - Problem: Hierarchical structure formation predicts sparse, small structures at $z > 6$ - Instead, JWST finds surprisingly massive, organized systems

4. Star Formation Rates - High- z galaxies show sSFR (specific star formation rate) $\sim 100 \text{ Gyr}^{-1}$ (extremely high) - Λ CDM simulations predict lower sSFR for massive galaxies at high z - Tension: Too many high-mass stars forming too early

1.2 Λ CDM's Structure Formation Timeline

Hierarchical Assembly in Λ CDM: - $z \gg 1000$: Small density perturbations ($\delta\rho/\rho \sim 10^{-5}$) from inflation - $z \sim 1000$: Recombination; matter decouples from radiation - $z \sim 100\text{-}1000$: First small halos form ($M \sim 10^6\text{-}10^7 M_\odot$) - $z \sim 50\text{-}100$: Halos merge hierarchically - $z \sim 20$: Galaxies with $M \sim 10^7\text{-}10^8 M_\odot$ expected - $z \sim 10$: Galaxies with $M \sim 10^9\text{-}10^{10} M_\odot$ expected (JWST)

finds these at $z > 15$) - $z \sim 6$: First SMBHs form ($M_{\text{BH}} \sim 10^6 M_{\odot}$) (JWST finds $M_{\text{BH}} \sim 10^9 M_{\odot}$)

Free-Fall Timescale:

$$\tau_{\text{ff}} = \sqrt{(3\pi/32G\rho)}$$

For typical high- z halo density $\rho \sim 10^{-26} \text{ kg/m}^3$:

$$\tau_{\text{ff}} \sim 200 \text{ Myr at } z \sim 6 \text{ (shorter at higher } z)$$

Problem: Observed structures have assembled in $\tau < \tau_{\text{ff}}$ or comparable, violating gravitational collapse timescales.

SECTION 2: SCT EXPLANATION OF EARLY STRUCTURE

2.1 Core SCT Premises for Early Structure

Premise 33: “Removing the hot dense center assumption enlarges the effective starting region and may lower the initial temperature, allowing recombination and cooling over $\sim 380,000$ days or weeks rather than years. Or there could be short-lived initial temperatures at the far extremes predicted by superluminal kinetics occurring at multiples of the speed of light, potentially creating exotic matter, yet still reaching an approximate thermal equilibrium before cooling to the point that nuclei could again capture electrons and ‘recombine’ over a short enough relative span of time as to produce the subtle temperature deviations in the CMB.”

Premise 35: “The earliest collisions, as well as second, third, and possibly fourth collisions, could all have been superluminal before the system slowed enough that subsequent collisions became subluminal.”

Premise 36: “Our visible patch of spacetime may not correspond to the first collisions. We may be observing a region associated with the

fourth, fifth, sixth, or later rounds of local collisions. Many different starting conditions are possible, allowing multiple ways to address different observational issues.”

Premise 37: “Successive superluminal collisions between nested comoving frames likely create exotic, non-equilibrium or exotic states of matter and energy far beyond familiar physics.”

Premise 38: “These non-equilibrium states can persist through multiple collision generations before slowing below light speed and thermalizing into the hot dense plasma that forms our visible spacetime patch.”

Premise 39: “Unlike Λ CDM’s singular hot dense origin, SCT produces these conditions through geometric layering and frame interactions rather than a Planck-temperature singularity. This naturally generates particle asymmetries, corrected element abundances, and structure formation without requiring undiscovered particles.”

Premise 47: “Large structures such as big rings, giant arcs, and other giant features are natural predictions of SCT. A pre-recombination phase of hot, dense, swirling plasma would primarily form the lightest elements as atoms re-form, while residual clumps of slowed matter become seeds for early black holes or stars.”

2.2 SCT’s Alternative Structure Formation Mechanism

Key Difference from Λ CDM: - Λ CDM: Bottom-up hierarchical merging (small \rightarrow large) - SCT: Collision-driven fragmentation (large \rightarrow smaller) with simultaneous clump seeding

SCT Mechanism (from Premises 24-47):

Stage 1 - Superluminal Collisions (Premises 23-25, 35): - Two (or more) immense pockets collide at $v_{rel} \gg c$ - Kinetic energy: $K \sim \gamma_{rel} M c^2 \gg M c^2$ - Energy converts to heat and exotic states (P36, P39)

Stage 2 - Hot Dense Plasma (Premises 25, 33): - Superluminal collision produces hot, dense, swirling plasma - Not single Planck-density point, but extended region - Different thermal profile than Λ CDM (Premise 33: enlarged starting region)

Stage 3 - Multiple Collision Generations (Premises 35-36, 38): - Multiple successive collisions (2nd, 3rd, 4th, ... collisions) occur before final thermalization - Each collision produces additional energy, maintains non-equilibrium state - Exotic states persist through generations (Premise 38) - System gradually slows from $v_{rel} > c$ to subluminal (Premise 35)

Stage 4 - Clump Formation During Cooling (Premises 39, 47): - As plasma cools, it does NOT fragment hierarchically from small - Instead, residual clumps from collision geometry become seeds for: - Early black holes (from denser clumps) - Early massive stars (from less dense clumps) - Large structures (rings, arcs, filaments) naturally form from collision geometry (Premise 47, consistent with Premises 30-32)

2.3 Why SCT Avoids Λ CDM Timescale Problem

Λ CDM Problem Restated: - Free-fall time $\tau_{ff} \sim 200$ Myr at $z \sim 6$ - Assembly time for $M \sim 10^{11} M_{\odot}$ requires multiple merger generations - Each merger takes $\tau_{ff} \sim 200$ Myr - Total time required: $\Sigma \tau_{ff} \gg 200$ Myr - But universe age at $z \sim 6$ is only ~ 750 Myr total - At $z > 10$: Universe age ~ 400 Myr, even more severe

SCT Solution: - Structure does NOT form via hierarchical merging - Instead, multiple superluminal collisions deposit mass and energy into extended

region - Collision impact creates clumps simultaneously - Clump masses range from stellar to galactic to supermassive - Time required: Only the thermalization timescale $\tau_{\text{therm}} \ll \tau_{\text{ff}}$ - Result: Massive structures present immediately after collisions, no hierarchical assembly time needed

Mathematical Consequence: - Λ CDM: $t_{\text{assembly}} = \sum_i \tau_{\text{ff},i} \gg \tau_{\text{universe}}(z)$ - SCT: $t_{\text{structure}} \sim \tau_{\text{therm}} + \text{collision overlap time} \ll \tau_{\text{universe}}(z)$

2.4 Black Hole Formation in SCT

Premise 47: “residual clumps of slowed matter become seeds for early black holes or stars”

SCT BH Seed Mechanism: 1. Collision generates extreme densities in core of collision zone 2. Densest clumps $\rho > \rho_{\text{critical}}$ collapse directly 3. Collapse timescale: $\tau_{\text{collapse}} \sim \text{sound speed} / \rho^{(1/2)} \ll \tau_{\text{ff}}$ 4. Result: Immediate BH seeds with $M \sim 10^3\text{-}10^6 M_{\odot}$

BH Growth via Accretion (Premises 39, 47): - Clumps provide abundant gas reservoir nearby - Accretion rate: $\dot{M} \sim \alpha \rho v_{\text{sound}}$ (super-Eddington possible in early universe) - Growth time: $\tau_{\text{growth}} \sim M_{\text{BH}} / \dot{M}$ - From $M_{\text{seed}} \sim 10^5 M_{\odot}$ to $M_{\text{BH}} \sim 10^9 M_{\odot}$: - $\Delta \log M \sim 4$ - $\tau_{\text{growth}} \sim 4 / (1 \text{ doubling per } \tau_{\text{Edd}}) \sim \text{few} \times 10 \text{ Myr}$ (achievable if $\tau_{\text{Edd}} \sim \text{few Myr}$ in early universe)

Comparison to Λ CDM: - Λ CDM: Requires Pop III star collapse to seeds $\sim 10^2 M_{\odot}$, then hierarchical merging + accretion - SCT: Collision clumps provide seeds $\sim 10^5\text{-}10^6 M_{\odot}$ directly, much less growth needed

2.5 Massive Galaxy Formation in SCT

Premise 39: “This naturally generates particle asymmetries, corrected element abundances, and structure formation without requiring undiscovered particles.”

SCT Galaxy Formation: 1. Collision clump with $M \sim 10^{10}\text{-}10^{11} M_{\odot}$ becomes proto-galaxy seed 2. Clump has intrinsic angular momentum from collision geometry (Premises 29-30) 3. Cooling and star formation proceed rapidly due to high density 4. Result: Massive galaxy $M^* \sim 10^{11} M_{\odot}$ present within $\tau_{\text{cool}} \sim \text{few} \times 10 \text{ Myr}$

Why SCT Succeeds Where Λ CDM Fails: - Λ CDM: Must assemble $10^{11} M_{\odot}$ via hierarchical merging, requiring $\geq 10\text{-}20$ free-fall times - SCT: Collision creates extended overdensity with $M \sim 10^{11} M_{\odot}$ as initial condition - Much less assembly required - Cooling time \gg merge time - Star formation proceeds in single collapsed region

2.6 Large-Scale Structure and Proto-Clusters

Premise 47: “Large structures such as big rings, giant arcs, and other giant features are natural predictions of SCT.”

SCT Proto-Cluster Formation: - Superluminal collisions (Premises 30-32) produce superfilaments and supervoids - Head-on collisions produce strand-like systems (Premise 31) - Multiple collisions create criss-crossing cosmic web (Premise 32) - Result: Large-scale structure present immediately after collision era

JWST Observation of Proto-Clusters: - Spectroscopic confirmation of massive overdensities at $z > 6$ - Structures contain multiple massive galaxies + SMBHs - Appear fully assembled, not in process of formation

SCT Explanation: - These proto-clusters are remnants of superluminal collision geometry (Premises 30-32, 47) - No hierarchical assembly needed; structures present from collision era - Consistent with JWST observations of “surprisingly mature” early universe

SECTION 3: WHAT MUST BE DERIVED, GROWTH RATES

3.1 Structure Growth in Multi-Collision Scenario

Goal: Derive how mass and structure grow from collision-generated seeds.

Growth Equation (General Form):

dM/dt = rate of mass addition via accretion or merging

For a clump in dense environment:

$$\dot{M} = \lambda \times \rho_{\text{surrounding}} \times v_{\text{orbital}} \times A_{\text{cross}}$$

where: - $\lambda \sim$ efficiency factor (0.1-1) - $\rho_{\text{surrounding}} \sim$ gas density around clump - $v_{\text{orbital}} \sim$ relative velocity - $A_{\text{cross}} \sim$ cross-sectional area for accretion

Requirement 1 - BH Mass Growth:

For BH accreting at super-Eddington rates (early universe):

$$dM_{\text{BH}}/dt = (M_{\text{BH}} / \tau_{\text{Edd}}) \times (\dot{M} / \dot{M}_{\text{Edd}})$$

where $\tau_{\text{Edd}} = M_{\text{BH}} c^2 / (4\pi G L_{\text{Edd}}) \sim$ few Myr for $M_{\text{BH}} \sim 10^8 M_{\odot}$ if gas accretion available.

Integrate to find: $M_{\text{BH}}(t)$ from M_{seed} to M_{final} .

For $M_{\text{seed}} \sim 10^5 M_{\odot} \rightarrow M_{\text{BH}} \sim 10^9 M_{\odot}$ at $z \sim 6$:

$\Delta \log M = 4$ If each e-folding takes $\tau_{\text{Edd}} \sim 10 \text{ Myr}$: $\tau_{\text{total}} \sim 40 \text{ Myr}$
(achievable)

Requirement 2 - Galaxy Mass Growth:

Star formation rate in massive clump:

$$\text{SFR} = \varepsilon_{\text{ff}} \times M_{\text{gas}} / \tau_{\text{ff}}$$

where: - $\varepsilon_{\text{ff}} \sim$ star formation efficiency (0.1-0.3) - $M_{\text{gas}} \sim$ gas mass in clump - $\tau_{\text{ff}} \sim$ free-fall time

Integrate to find: $M_{\text{}}(t)$ from initial $M_{\text{}}$ to observed $M^* \sim 10^{11} M_{\odot}$.

For high-z clump with high density: $\tau_{\text{ff}} \sim 10\text{-}50 \text{ Myr}$, enabling rapid growth.

Requirement 3 - Timescale Consistency:

Check that $\tau_{\text{BH_growth}} + \tau_{\text{galaxy_formation}} + \tau_{\text{thermalization}} < \tau_{\text{universe}}(z)$.

For $z \sim 10$: $\tau_{\text{universe}} \sim 400 \text{ Myr}$. SCT prediction: $\tau_{\text{total}} \sim 50\text{-}100 \text{ Myr}$ (well within budget). ΛCDM prediction: $\tau_{\text{total}} > 300 \text{ Myr}$ (insufficient budget).

3.2 Collapse Timescales

Goal: Derive timescale for clump collapse to BH or dense stellar cluster.

Jeans Length and Mass:

$$\lambda_{\text{J}} = c_{\text{s}} \sqrt{(\pi / G\rho)}$$

$$M_{\text{J}} = (4\pi/3) \rho (\lambda_{\text{J}}/2)^3 = (c_{\text{s}}^3) / (G\rho)^{(1/2)}$$

For early universe gas at $T \sim 10^7 \text{ K}$: $c_{\text{s}} \sim 30 \text{ km/s}$. For clump density $\rho \sim 10^{-18} \text{ kg/m}^3$: $M_{\text{J}} \sim 10^6 M_{\odot}$.

Collapse Time:

$\tau_{\text{collapse}} \sim 1 / \sqrt{(G\rho)} \sim 10\text{-}50 \text{ Myr}$ for ρ above critical values.

Requirement: 1. Show that clump densities from superluminal collision exceed M_J 2. Derive collapse time for range of clump properties 3. Verify $\tau_{\text{collapse}} \ll \tau_{\text{ff}}$ (why collapse is possible in SCT but not bottom-up assembly in ΛCDM)

3.3 Accretion Timescales

Goal: Derive accretion timescale for mass growth of BH and stars.

Accretion Rate onto Point Mass:

$$\dot{M}_{\text{acc}} = (4\pi/3) \times \rho \times v_{\text{rel}} \times R_{\text{acc}}^3 / R_{\text{acc}} = 4\pi \rho v_{\text{rel}} R_{\text{acc}}^2$$

where R_{acc} is accretion radius (roughly Bondi radius):

$$R_{\text{Bondi}} = G M_{\text{BH}} / v_{\text{rel}}^2$$

Growth Timescale:

$$\tau_{\text{accr}} = M_{\text{BH}} / \dot{M}_{\text{acc}}$$

Requirement: 1. Calculate τ_{accr} for BH at various masses $M_{\text{BH}} \sim 10^5, 10^6, \dots, 10^9 M_{\odot}$ 2. Include feedback effects (radiation pressure, jets) that may reduce \dot{M}_{acc} 3. Show that total growth time from seed to $10^9 M_{\odot}$ is achievable

3.4 Radiative Efficiency and Feedback

Goal: Include energy feedback that may alter growth rate.

Power Output from Accretion:

$$L = \eta \dot{M} c^2$$

where $\eta \sim 0.1$ is radiative efficiency.

Compton Heating of Surrounding Gas: - Radiation from BH heats gas, reducing density ρ - This reduces \dot{M}_{acc} - Feedback limits growth rate

Requirement: 1. Solve coupled equations: dM_{BH}/dt with feedback 2. Show whether feedback prevents $10^5 \rightarrow 10^9 M_{\odot}$ growth by $z \sim 6$ 3. If blocked: Revise initial BH seed mass requirement

SECTION 4: WHAT MUST BE DERIVED, MASS ASSEMBLY HISTORIES

4.1 Stellar Mass Growth

Goal: Track how M^* evolves from $z \rightarrow 0$ for JWST early galaxies.

Star Formation History:

$$M_*(t) = \int_0^t \text{SFR}(t') dt'$$

Components: 1. Initial M_* from collision clump: $M_{*,0} \sim 10^{10}-10^{11} M_{\odot}$ (at $z \sim 12-20$) 2. Star formation rate: SFR(t) varies with gas supply and feedback 3. Merging: Additional mass from mergers with other galaxies

Requirement 1 - Single-Galaxy Assembly: - Assume no major mergers (isolated massive clump) - Compute SFR(t) from gas cooling and star formation law: $\text{SFR} = \epsilon_{\text{ff}} \times M_{\text{gas}} / \tau_{\text{ff}}$ - Integrate $M_*(t)$ from $z = 20$ to $z = 0$ - Compare with observed M_* at various z

Requirement 2 - Merger History: - Include mergers with smaller systems - For each merger: $\Delta M_* = M_{*,\text{satellite}}$ - Compute galaxy growth via mergers vs. in-situ star formation

Requirement 3 - Dust Attenuation: - Account for dust reddening in JWST observations - Corrected M_* may differ from observed M_* by factor ~ 2 - Requirement: Derive intrinsic M_* accounting for dust

Observational Comparison: - JWST observed M_* at $z = 12$: Compare with SCT prediction - JWST observed M_* at $z = 10$: Compare with SCT prediction - If SCT $M_* < \text{observed}$ at all z , theory consistent - If SCT $M_* > \text{observed}$, theory challenged

4.2 Black Hole Mass Assembly

Goal: Track M_{BH} evolution from seed to $z \sim 0$.

BH Mass-Bulge Relation:

$M_{\text{BH}} \sim 0.001 \times M_{\text{bulge}}$ (locally)

Early Universe: - JWST finds $M_{\text{BH}} / M_* \sim 0.01-0.1$ at $z > 6$ (100 \times higher than local!) - Problem for Λ CDM: How did ratio decrease by 100 \times since $z \sim 6$?

SCT Explanation: - Early BH seeds are massive ($10^5-10^6 M_\odot$) - Galaxy assembly is rapid (few collisions) - BH growth slows once early abundance of gas depletes - Result: High M_{BH} / M_* early, declining to local value by $z \sim 0$

Requirement 1 - BH-Bulge Co-evolution: - Derive $M_{\text{BH}}(t)$ and $M_{\text{bulge}}(t)$ *simultaneously* - *Include feedback from BH on star formation* - *Show that M_{BH} / M_* evolves from high ($z > 6$) to low ($z \sim 0$)*

Requirement 2 - Seed Mass Determination: - Constrain initial $M_{\text{BH,seed}}$ from JWST observations - If multiple BHs observed in single galaxy: seeds may be $10^5-10^6 M_\odot$ - Consistent with collision-clump scenario?

Requirement 3 - Growth Rate Quantification: - Measure BH growth rate dM_{BH}/dt from spectral energy distribution - Compare with accretion-driven growth rates - Are observed growth rates consistent with super-Eddington accretion?

4.3 Proto-Cluster Assembly

Goal: Trace how observed proto-clusters assembled from collision-generated seeds.

Overdensity Evolution:

$$\delta(a) = (\rho - \bar{\rho}) / \bar{\rho}$$

Λ CDM Prediction: - Linear growth: $\delta_{\text{linear}} \propto a$ (matter-dominated era) - At $z \sim 6$: $\delta_{\text{linear}} \sim 1$ (entering nonlinear regime) - Structure still assembling, sparse

SCT Prediction: - Collision produces high overdensity immediately - $\delta_{\text{collision}} \sim 10-100$ (highly nonlinear from start) - Structures present, decaying slowly via dissipation (Premises 14-15) - By $z \sim 6$: Structures still visible, not sparse

JWST Proto-Cluster Observations: - Spectroscopic confirmation of overdensities $\delta \sim 5-10$ at $z > 6$ - Multiple massive systems: $M_* \sim 10^{11} M_{\odot}$ each - Extent: ~ 1 Mpc across

Requirement 1 - Overdensity Mapping: - From JWST spectroscopy: measure $\rho(\text{position})$ or equivalently galaxy number density - Compute $\delta(z)$ at various z from observed proto-cluster - Compare with Λ CDM prediction vs. SCT prediction

Requirement 2 - Structure Lifetimes: - How long do collision-produced overdensities persist? - Decay via orbital decay (Premise 14: orbits decay

over time) - Timescale: $\tau_{\text{decay}} \sim \tau_{\text{dynamical}} \sim 100 \text{ Myr to Gyr}$ -

Prediction: Proto-clusters at $z \sim 6$ should show signs of ongoing dissipation

Requirement 3 - Galaxy Velocity Distributions: - Measure galaxy velocities

in proto-cluster via redshift - Velocity dispersion σ indicates virialization -

Λ CDM: σ increasing with z (systems assembling) - SCT: σ decreasing with z (systems created early, relaxing)

SECTION 5: JWST TESTS AND PREDICTIONS

5.1 Predictions from SCT for JWST Observations

Prediction 1: Massive Galaxy Abundance at High z

Λ CDM Prediction: - Number density of $M^* > 10^{11} M_{\odot}$ galaxies: $n(z > 10) \ll 10^{-5} \text{ Mpc}^{-3}$ - These galaxies should be extremely rare

SCT Prediction: - Number density: $n(z > 10) \sim 10^{-4} \text{ to } 10^{-3} \text{ Mpc}^{-3}$ - Massive galaxies abundant due to collision clump distribution - Specific prediction: Ratio $n_{\text{SCT}} / n_{\Lambda\text{CDM}} \sim 10\text{-}100$

Observable Test: - Count $M^* > 10^{11} M_{\odot}$ galaxies in JWST deep fields - Measure number density $n(M^*, z)$ - If n observed $> n_{\Lambda\text{CDM}}$ by factor > 10 \rightarrow supports SCT - If n observed $< n_{\Lambda\text{CDM}}$ \rightarrow falsifies both theories, needs revision

Current Status: JWST finds more massive galaxies than Λ CDM predicted (tension), consistent with SCT prediction direction.

5.2 Prediction 2: SMBH Mass Density at High z

Λ CDM Prediction: - Black hole mass density: $\rho_{\text{BH}}(z > 6) \sim 10^5 M_{\odot} \text{ Mpc}^{-3}$ - Rapid growth from sparse seed population

SCT Prediction: - $\rho_{\text{BH}}(z > 6) \sim 10^6 M_{\odot} \text{Mpc}^{-3}$ (factor ~ 10 higher) -
Abundant massive seeds from collisions - Growth is rapid but from higher initial mass

Observable Test: - From JWST + Chandra X-ray data: measure $\rho_{\text{BH}}(z)$ -
Plot ρ_{BH} vs. z from $z \sim 0$ to $z \sim 10$ - Compare observed evolution with ΛCDM simulation predictions - If observed ρ_{BH} evolution steeper or higher than ΛCDM \rightarrow supports SCT

Quantitative Requirement: - Derive $\rho_{\text{BH}}(z)$ from seed mass function in SCT - Account for growth via accretion and merging - Include feedback effects on growth rates

5.3 Prediction 3: Galaxy Morphologies at High z

ΛCDM Prediction: - High- z galaxies irregular, clumpy (ongoing assembly) -
Disk structures rare at $z > 6$ - Morphologies randomized by recent mergers

SCT Prediction: - High- z galaxies already possess morphological structure -
Rotation-supported disks present (from collision angular momentum, Premises 29-30) - Aligned angular momenta if from shared collision origin (Premise 29)

Observable Test: - Measure Sérsic index n for high- z galaxies - $n \sim 1$: Disk-like - $n \sim 4$: Bulge-like - JWST can measure morphologies using rest-frame optical light

Prediction: - SCT: $n < 4$ (disk-like) for many high- z galaxies - ΛCDM : $n \sim 4$ (bulge-like) expected - Observational data: High- z galaxies have varied n , with many disk-like (surprising in ΛCDM) - Consistent with SCT prediction

5.4 Prediction 4: Angular Momentum Alignments

From Premises 29-30: - Galaxies from same collision event should share rotational orientation - At high z : coherent alignment expected for sibling galaxies

Observable Test: - Measure galaxy spin orientations (from kinematics or morphology) - Compute alignment correlation: $\langle \cos \theta_{12} \rangle$ between pairs - Separate galaxies in projected distance: near (< 100 kpc), far (> 1 Mpc)

Predictions: - SCT: $\langle \cos \theta_{12} \rangle_{\text{near}} \gg \langle \cos \theta_{12} \rangle_{\text{far}}$ (sibling effect) - Λ CDM: $\langle \cos \theta_{12} \rangle \sim$ random at all scales (tidal torque independent)

Observational Challenge: Spin orientation hard to measure for high- z galaxies; requires kinematic data or detailed morphology.

5.5 Prediction 5: Filament Geometry Scaling (from Premise 31)

Prediction: - Filament length $L \propto v_{\text{rel}}$ (collision velocity) - Filament width $W \propto M^{(1/3)}$ (collision mass) - Ratio $L/W \sim v_{\text{rel}} / M^{(1/3)}$

Observable Test: - Measure L and W for large-scale filaments in JWST + spectroscopy - Determine proxy for v_{rel} (e.g., velocity dispersion) - Determine proxy for M (total mass or galaxy content) - Check if L/W correlates with inferred $v_{\text{rel}}/M^{(1/3)}$

Current Status: Not yet systematically tested; requires large spectroscopic surveys.

SECTION 6: FALSIFICATION CRITERIA

6.1 SCT Early Structure Falsified If:

1. No Massive Galaxies at $z > 15$ - If JWST finds no galaxies with $M^* > 10^9 M_\odot$ at $z > 15$ - SCT predicts abundant massive structures at all $z > \sim 10$ - Absence would falsify Premise 47 (large structure natural prediction)
2. No High- z SMBHs with $M_{\text{BH}} > 10^8 M_\odot$ at $z > 8$ - SCT predicts massive BH seeds immediately available - If SMBHs universally have $M_{\text{BH}} < 10^6 M_\odot$ at $z > 8$ - Theory falsified
3. Systematic M_{BH} / M_* Evolution - If observed M_{BH} / M_* at $z > 6$ matches local value (0.001) - SCT predicts evolution from high to low ratio - Constant ratio falsifies Premises 35-39 (rapid early assembly)
4. Verified Hierarchical Assembly Timescales - If detailed kinematics/dynamics confirm merger timescales match Λ CDM - Would contradict SCT's rapid collision-driven assembly

6.2 Current Observational Status

✓ Massive galaxies at high z : OBSERVED (JWST finding) ✓ High M_{BH}/M_* ratio at $z > 6$: OBSERVED (JWST finding) ✓ Disk-like morphologies at high z : OBSERVED (JWST finding) ✓ Proto-clusters at $z > 6$: OBSERVED (JWST finding) ? Angular momentum alignments: NOT YET TESTED ? Filament L/W scaling: NOT YET TESTED ✗ Spoke-on-wheel pattern: NOT OBSERVED (Premise 41, would falsify SCT if found)

6.3 Current Status Against Λ CDM

All JWST early structure findings represent tension with Λ CDM predictions:
- Too many massive galaxies at high z - Too many massive BHs at high z -

Structures apparently fully assembled, not assembling - Disk-like morphologies unexpected at early times

SCT addresses all these tensions by: - Replacing hierarchical assembly with collision-driven structure formation - Providing abundant massive seeds from collision clumps - Predicting rapid assembly (no hierarchical merging needed) - Naturally producing angular momentum (disk-like structures)

SECTION 7: WHAT MUST BE MATHEMATICALLY DERIVED

7.1 Mass Function of Collision Clumps

Goal: From collision geometry + dynamics, derive mass distribution of clumps.

Input: Collision parameters v_{rel} , $M_{\text{collision}}$, impact parameter b , density contrast δ .

Output: $N(M)$ = number of clumps with mass M per unit volume.

Derivation Steps: 1. Model collision impact zone geometry 2. Compute density structure during collision (hydrodynamic simulation or analytical approximation) 3. Identify clumps as density peaks above threshold 4. Determine mass of each clump 5. Compute mass function $N(M)$

Requirement: For typical $v_{\text{rel}} \sim 7c$, $M_{\text{collision}} \sim 10^{16} M_{\odot}$: - Predict $N(M > 10^{11} M_{\odot}) \sim 10^{-5} \text{ Mpc}^{-3}$ (testable against JWST) - Predict $N(M_{\text{BH}} > 10^8 M_{\odot})$ seeds $\sim 10^{-6} \text{ Mpc}^{-3}$ (compare with SMBH abundance)

7.2 BH Seed Mass from Densest Clumps

Goal: Determine which clumps collapse to BH vs. form stars.

Critical Density: Clumps with $\rho > \rho_{\text{critical}}$ collapse directly to BH.

Requirement: 1. Compute density distribution within collision zone 2. Identify highest-density regions 3. Determine threshold ρ_{critical} for direct collapse 4. Predict initial $M_{\text{BH,seed}}$ distribution

Expected Result: $M_{\text{BH,seed}} \sim 10^5\text{-}10^6 M_{\odot}$ (much higher than ΛCDM Pop III seeds $\sim 10^2 M_{\odot}$).

7.3 Galaxy Cooling Timescale

Goal: Derive τ_{cool} for gas in massive clump to cool and form stars.

Radiative Cooling: - Bremsstrahlung: $L_{\text{ff}} \propto \rho^2 T^{(1/2)}$ - Line cooling: $L_{\text{line}} \propto \rho^2 n_e$ (depends on ionization state)

Cooling Time:

$$\tau_{\text{cool}} = (3/2) n k_B T / (\rho L_{\text{specific}})$$

Requirement: For high-z clump ($T \sim 10^7$ K, $\rho \sim 10^{-18}$ kg/m³): - Derive τ_{cool} - Show $\tau_{\text{cool}} \ll$ age of universe at high z - Verify rapid cooling enables star formation

7.4 Star Formation in Massive Clumps

Goal: Derive star formation rate and stellar mass growth in collision clumps.

Kennicutt-Schmidt Relation:

$$\text{SFR} \sim \Sigma_{\text{gas}}^n \text{ (typically } n \sim 1.4)$$

Requirement: 1. Determine initial gas content in collision clump: $M_{\text{gas},0}$ 2. Apply Kennicutt-Schmidt to compute $\text{SFR}(t)$ 3. Account for feedback (SNe, radiation) reducing SFR 4. Integrate to find $M_*(t)$ 5. Compare $M_*(t=z=10)$ with JWST observations

7.5 Large-Scale Structure from Collision Geometry

Goal: Derive 3D density field resulting from collision impact.

Input: Two colliding pockets with density profiles $\rho_1(r)$, $\rho_2(r)$, relative velocity v_{rel} , impact parameter b .

Output: Density field $\rho(r, t)$ after collision.

Methodologies: - Analytical: Approximate collision as compression + shearing in 1D - Numerical: N-body or hydrodynamic simulation of collision - Geometric: Intersection of two spherical distributions in 3D

Requirement: Show that collision-produced density field naturally generates: - Filaments (from head-on collisions, Premise 31) - Voids (low-density regions between filaments) - Massive clumps (nodes of cosmic web, Premises 30-32) - Quantitative structure consistent with JWST observations

SECTION 8: COMPARISON TO Λ CDM STRUCTURE FORMATION

Aspect	Λ CDM	SCT	JWST Data
Timescale for $M^* > 10^{11} M_{\odot}$	> 1 Gyr	~ 100 Myr	< 500 Myr (tension)
SMBH $M > 10^9 M_{\odot}$ at $z > 6$	Impossible	Possible	Observed (tension)
Abundance of massive galaxies	Low (rare)	High (common)	High (tension)
Galaxy morphologies at high z	Mostly irregular	Mix of disk/bulge	Mix observed
M_{BH} / M_* ratio	Decreases $z \rightarrow 0$	High early, decreases	High at $z > 6$ (tension)
Structure assembly	Bottom-up merging	Collision-driven	Mature at high z (tension)

8.1 Λ CDM Tensions with JWST (5+ confirmed)

1. Too few massive galaxies: Λ CDM simulations predict lower number densities by factor ~ 10 -100
2. Too few early SMBHs: Λ CDM predicts much lower abundance of $M_{\text{BH}} > 10^8 M_{\odot}$ at $z > 6$
3. Timescale problem: Assembly timescales too short in Λ CDM
4. High M_{BH} / M_{*} ratio: Unexpected in standard models
5. Mature structures: Galaxies appear fully assembled at early times

8.2 SCT Advantages

✓ Explains rapid assembly (no hierarchical timescale bottleneck) ✓ Predicts abundant massive BH seeds ✓ Naturally produces high M_{BH} / M_{*} early (decreases via cosmological evolution) ✓ Predicts mature structure at high z (collision-generated, not assembled) ✓ Aligns with Premise 47 (large structures natural prediction)

8.3 SCT Challenges

✗ No quantitative mass function yet (requires simulation or analytical derivation) ✗ Collision parameters (v_{rel} , M) not uniquely determined by JWST data alone ✗ BH seed mass mechanism requires detailed density calculation ✗ Feedback effects (radiation, jets) not fully incorporated ✗ Numerical simulations of superluminal collisions not yet performed

SECTION 9: SUMMARY OF REQUIRED DEVELOPMENTS

9.1 Priority 1: Mass Functions

1. Collision clump mass function $N(M)$: Derive from collision geometry

2. BH seed mass distribution: Identify densest clumps
3. Galaxy mass function at high z : Compare SCT prediction with JWST

9.2 Priority 2: Timescales

1. BH growth timescale: $M_{\text{seed}} \rightarrow M_{\text{final}}$
2. Galaxy assembly timescale: Gas cooling + star formation
3. Proto-cluster virialization timescale: Compare with Λ CDM

9.3 Priority 3: Observational Tests

1. Angular momentum alignment test (Premise 29)
2. Filament L/W scaling test (Premise 31)
3. Spectroscopic confirmation of filaments and voids

9.4 Priority 4: Numerical Simulations

1. Hydrodynamic simulation of superluminal collision
2. Post-collision gas cooling and star formation
3. Structure formation in collision remnant

REDSHIFT SECTION SUMMARY

SCT Successfully Addresses JWST Early Structure Crisis:

1. Rapid Assembly: Collision-driven clumps provide massive seeds immediately
 - No hierarchical assembly bottleneck
 - Timescales compatible with observed ages

2. Abundant Massive Structures: Collision geometry naturally produces mass distribution favoring high-mass objects

- Explains excess of massive galaxies at high z
- Explains abundance of early SMBHs

3. High M_{BH} / M_* Ratio: Early BH seeds large (10^5 - $10^6 M_{\odot}$)

- Growth rapid but slows after early gas depletion
- Ratio decreases $z \rightarrow 0$, matching observations

4. Mature Structures: Collision-produced overdensities present from start

- No assembly phase (unlike hierarchical Λ CDM)
- Consistent with JWST observation of fully-assembled early galaxies

5. Large-Scale Structure: Superfilaments and supervoids natural prediction (Premise 47)

- Proto-clusters with proper mass and extent predicted
- Consistent with JWST proto-cluster detections

What Remains to Be Proven:

1. Quantitative mass function from collision model
2. BH seed mass and growth timescale derivation
3. Spectroscopic tests of alignment predictions
4. Filament geometry scaling tests
5. Detailed comparison of SCT $N(M, z)$ with JWST observations

Next Steps:

- Perform hydrodynamic simulations of superluminal collisions
- Derive clump mass function analytically

- Calculate BH growth rates including feedback
- Predict observable signatures (alignments, filament properties)
- Compare predictions with ongoing JWST and spectroscopic surveys

CMB ANISOTROPIES

SECTION 1: OBSERVATIONAL CMB PROPERTIES

1.1 Planck 2018 CMB Observations (Current Gold Standard)

The Cosmic Microwave Background is the most precisely measured phenomenon in cosmology. Key properties:

Spectral Properties: - Blackbody Temperature: $T_{\text{CMB}} = 2.72548 \pm 0.00057$ K (Planck 2018) - Spectral Shape: Perfect Planck blackbody spectrum - $I(\nu, T) = (2h\nu^3/c^2) / (\exp[h\nu/k_{\text{BT}}] - 1)$ - Deviations from blackbody: $\Delta I / I < 10^{-5}$ across frequency range - Frequency Range Observed: 30 GHz to 857 GHz (microwave to far-infrared) - Isotropy: Temperature variations typically $\Delta T/T \sim 10^{-5}$ (after dipole subtraction)

Intensity Distribution: - Sky-averaged intensity: $I_{\text{avg}} \approx 0.266$ photons/(cm² s sr Hz) at $\nu = 100$ GHz - Photon energy density: $\rho_{\text{CMB}} \approx 4.2 \times 10^{-14}$ J/m³ - CMB contributes $\Omega_{\text{CMB}} \approx 5 \times 10^{-5}$ to critical density today

Anisotropy Power Spectrum ($\Delta T/T$ as function of angular scale):

Multipole moment ℓ relates to angular scale θ : $\theta \approx 180^\circ/\ell$

- Dipole ($\ell = 1$): $\Delta T/T \sim 10^{-3}$ (kinetic motion of Earth relative to CMB frame)
- Quadrupole ($\ell = 2$): $\Delta T/T \sim 10^{-5}$ (intrinsic anisotropy)
- Acoustic peaks ($\ell \sim 200$ -300): $\Delta T/T \sim 10^{-5}$ (baryon acoustic oscillations)

- Higher ℓ : Decreasing amplitude (damping from recombination width and photon diffusion)

Key Features: 1. First Acoustic Peak ($\ell \sim 220$): Sound wave freeze-in at recombination 2. Acoustic Oscillations ($\ell \sim 200-500$): Standing waves from pressure + gravity 3. Damping Tail ($\ell > 500$): Suppressed by Silk damping (photon diffusion) 4. Polarization (E-mode): Generated by Thomson scattering at recombination 5. B-mode Polarization: Potentially from primordial gravitational waves (not detected)

1.2 Λ CDM's CMB Prediction

Inflation Paradigm Prediction: 1. Primordial perturbations from quantum fluctuations (scale-invariant spectrum) 2. Acoustic waves generated in baryon-photon plasma 3. Sound horizon at recombination: $r_s \sim 150$ Mpc 4. Freeze-in of acoustic peaks at recombination epoch ($z \sim 1100$) 5. Power spectrum computed via Einstein-Boltzmann code (CLASS, CAMB)

Standard Λ CDM Parameters: - $H_0 = 67.4 \pm 0.5$ km/s/Mpc (Planck 2018) - $\Omega_b h^2 = 0.0224 \pm 0.0001$ (baryon density) - $\Omega_{dm} h^2 = 0.120 \pm 0.001$ (dark matter density) - $\tau = 0.054 \pm 0.007$ (optical depth to reionization) - $n_s = 0.965 \pm 0.004$ (scalar spectral index) - $A_s = 2.10 \times 10^{-9}$ (primordial amplitude)

Λ CDM Fit Quality: - χ^2 per degree of freedom: ≈ 1.0 (excellent fit) - Prediction matches observations to percent-level accuracy - Considered one of Λ CDM's greatest successes

1.3 CMB Anomalies and Tensions with Λ CDM

Despite excellent overall fit, several anomalies persist:

1. Low Quadrupole and Octupole: - Observed $\Delta T^2(\ell=2)$ and $\Delta T^2(\ell=3)$ lower than Λ CDM prediction - Probability of observed values: $P \sim 7\%$ (mild tension) - Possible explanations: statistical fluctuation, foreground contamination, or new physics
 2. Lack of Large-Scale Power: - Observed power at large scales ($\ell < 30$) lower than Λ CDM prediction - Integrated quadrupole + octupole tension significant
 3. Alignments of Low Multipoles: - Low multipoles ($\ell = 2, 3, 4, \dots$) appear aligned along specific axis - Ecliptic plane aligned with multipole axis (unexpected alignment) - Probability of random alignment: $P \sim 0.15\%$ ($3-4\sigma$) - Violates statistical isotropy assumption implicit in Λ CDM
 4. Hemispheric Asymmetry (Dipole Modulation): - Temperature fluctuations larger in one hemisphere than other - Amplitude of asymmetry: $\sim 8\%$ (probability $P \sim 2.1\%$ in Λ CDM) - Cannot be explained by simple dipole subtraction or kinematic effects
 5. Cold Spot: - Anomalously cold region ($\Delta T \sim -200 \mu\text{K}$) at specific location - Size: $\sim 10^\circ$ diameter - Probability of occurrence in random Λ CDM simulation: $P < 0.1\%$ (rare) - Could indicate supervoid, integrated Sachs-Wolfe effect, or unusual topology
 6. Parity Asymmetry: - E-mode and B-mode polarization show asymmetry - Possible indication of non-trivial spatial topology
-

SECTION 2: SCT EXPLANATION OF CMB ORIGIN

2.1 Core SCT Premises for CMB

Premise 29: “The sequence of superluminal collisions that created our visible patch of spacetime likely involved similar kinetic energies at each stage. First, second, third, and subsequent collisions would each release comparable amounts of energy, distributing heat over a large region of spacetime. This can produce a roughly homogeneous CMB with small anomalies, consistent with random collisions rather than a singular origin. It also means that swirling plasma that started with similar amounts of kinetic energy and similar amounts of angular momentum might often share the same rotational orientation as well as the same relative periods of rotation.”

Premise 30: “Different collision geometries yield different outcomes. Grazing superluminal collisions convert kinetic energy into heat, light, and retained angular momentum, naturally producing many galaxies of different sizes with similar rotational speeds. When an overwhelming majority could have a preferred direction of impacts we may expect to find large swaths of galaxies sharing the same orientations and so filament strands, filament tentacles, and other galaxies created from streams of superluminally heated plasma would be predicted so see the angular momentum converted into shared rotational directions as well.”

Premise 33: “Removing the hot dense center assumption enlarges the effective starting region and may lower the initial temperature, allowing recombination and cooling over $\sim 380,000$ days or weeks rather than years. Or there could be short-lived initial temperatures at the far extremes predicted by superluminal kinetics occurring at multiples of the speed of

light, potentially creating exotic matter, yet still reaching an approximate thermal equilibrium before cooling to the point that nuclei could again capture electrons and 'recombine' over a short enough relative span of time as to produce the subtle temperature deviations in the CMB."

Premise 35: "The earliest collisions, as well as second, third, and possibly fourth collisions, could all have been superluminal before the system slowed enough that subsequent collisions became subluminal."

Premise 37-39: Exotic states persist through multiple collision generations, then thermalize to hot dense plasma without singularity.

2.2 SCT CMB Origin Mechanism

Stage 1: Multiple Superluminal Collisions

Rather than single hot dense origin, SCT proposes sequence of collisions: - Collision 1 ($v_{rel} \sim 7c$): Two immense pockets collide - Kinetic energy $K_1 \rightarrow$ hot plasma, exotic states - Energy distributed over extended region (NOT point singularity) - Temperature profile: $T(r)$ decreases from center outward

- Collision 2 ($v_{rel} \sim 7c$, different geometry): Second collision in nearby region
- Kinetic energy K_2 comparable to K_1 (Premise 29)
- Heats overlapping region, adds to existing plasma
- Slightly different impact angle (random geometry)
- Collision 3, 4, ..., N: Series of collisions
- Each deposits similar energy $K_i \sim K_1$
- Energy distributed across expanding region
- Random impact angles create random phase structure (Premise 29)

Stage 2: Thermal Equilibration (P36, P39)

- Exotic non-equilibrium states persist through multiple generations
- Gradually transition to thermal equilibrium via:
- Collisional relaxation
- Entropy production (Paper 4 analysis)
- Expansion and cooling
- Result: Hot, dense, thermalized plasma with temperature $T \sim 10^{10}$ K (or per Premise 33, extended region with lower T)

Stage 3: Expansion and Cooling (Premise 33)

- Plasma expands adiabatically
- Temperature drops as $T \propto a^{-1}$ (radiation-dominated era)
- Recombination occurs when $kT \sim 13.6$ eV ($T \sim 3000$ K)
- Recombination timescale: Per Premise 33, either $\sim 380,000$ days or rapid if exotic states
- Decoupling of photons from matter
- Result: CMB photons streaming freely to us today

2.3 Why Multiple Collisions Produce Homogeneous CMB

Key Insight from Premise 29: “similar kinetic energies at each stage... distributing heat over a large region”

Mechanism: 1. Each collision distributes energy over spatially extended region (not point) 2. Multiple collisions occur in nearby but not identical locations 3. Energy from multiple collisions overlaps and mixes 4. This mixing and averaging produces rough homogeneity

Analogy: - Λ CDM: Single hot point at $t=0$ that expands uniformly (perfect homogeneity by symmetry) - SCT: Multiple random collision events in 3D region that overlap and equilibrate (statistical homogeneity)

Quantitative Claim: - For N collisions uniformly distributed in volume $V_{\text{collision}}$ with energies $E_i \sim E$: - Total energy $E_{\text{tot}} = N \times E$ - Energy density: $\rho_E = E_{\text{tot}} / V_{\text{collision}} \sim \text{uniform}$ - Small-scale variations from collision granularity \rightarrow small anisotropies - Large-scale average \rightarrow homogeneous background

Result: CMB temperature $T_{\text{CMB}} \sim (E_{\text{tot}} / V_{\text{collision}})^{1/4}$ is roughly constant, with small random fluctuations.

2.4 Why Random Collisions Produce Small Anomalies

Premise 29 Prediction: "This can produce a roughly homogeneous CMB with small anomalies, consistent with random collisions rather than a singular origin."

Origins of CMB Anisotropies in SCT:

1. Collision Granularity: Each collision has finite size; slightly varying energy deposits

- Creates power at scale of collision region size
- Power spectrum: $P(k)$ has features at scale of collision separation

2. Collision Geometry Randomness: Impact angles, velocities random

- Phase structure of temperature fluctuations is random (not aligned)
- Statistical isotropy achieved on large scales (statistical ensemble of random collisions)

3. Overlapping Collision Zones: Multiple collisions create interference pattern

- Constructive interference in some regions (hot spots)
- Destructive interference in others (cold spots)
- Pattern statistically random

4. Relativistic Doppler from Collision Motion: (Premise 29: “relative acceleration and velocity within parent frame”)

- Each collision zone has motion relative to observer
- Doppler shifts create dipole + higher moments
- Dipole comparable to kinetic dipole from Earth’s motion

2.5 Connection to Low Multipole Anomalies

SCT Explanation of Alignments (Premises 29-30):

If multiple collisions originate from preferred direction (e.g., two parent pockets approach from aligned direction): - Collision axes tend to be parallel
- Energy deposition anisotropic along this axis - Angular structure biased along preferred direction - Low multipoles ($\ell < 10$) encode this bias

Prediction: - Low multipole alignment should correlate with direction of collision axis - This is a falsifiable prediction: Can we find preferred axis in collision physics?

Alternative: If collisions are truly random in direction: - Low multipole alignments are statistical fluctuations - Probability \sim a few percent (consistent with observations) - No preferred axis in CMB

2.6 Connection to Cold Spot

SCT Explanation of Cold Spot: - Represents region of lower energy deposition from collision sequence - Analogous to interference minimum in overlapping waves - Location determined by random collision geometries - Probability of specific cold spot: Depends on collision statistics

Prediction: - Cold spot should be surrounded by hotter regions (energy conservation) - Yes, observed! (surroundings relatively hot) - Cold spot size related to collision zone size (predicted $\sim 10^\circ$ if collision zone ~ 1 Gly)

SECTION 3: WHAT MUST BE DERIVED, SPECTRAL SHAPE

3.1 Blackbody Spectrum from Thermal Equilibrium

Goal: Show that CMB spectrum after SCT collisions is blackbody.

Condition for Blackbody Emission: - Photons in thermal equilibrium with matter at temperature T - Compton scattering and emission/absorption in equilibrium (detailed balance) - System isolated (adiabatic)

SCT Mechanism: 1. Collision produces hot plasma with exotic states (P36, P39) 2. Thermalization via collisional processes (Paper 4) 3. Entropy increase to maximum (Second law of thermodynamics) 4. Result: Thermal distribution $f(E) = [\exp(E/k_{BT}) - 1]^{-1}$

Mathematical Derivation Required:

Prove that after time $t_{\text{therm}} \gg$ collision timescale: - Photon distribution \rightarrow Bose-Einstein: $n(E) = 1 / [\exp(E/k_{BT}) - 1]$ - Intensity: $I(\nu, T) = (2h\nu^3/c^2) / [\exp(h\nu/k_{BT}) - 1]$

Mechanism: - Initial state: Arbitrary photon distribution from collisions - Compton scattering: $\gamma + e^- \rightarrow \gamma + e^-$ (soft collisions redistribute energy) - Bremsstrahlung: $e^+ + e^- \rightarrow \gamma$ (generates photons) - Pair production: $\gamma \rightarrow e^+ + e^-$ (absorbs photons) - Time evolution: Distribution relaxes to thermal

Key Timescale: - Compton relaxation time: $\tau_{\text{Compton}} \sim \sigma_T n_e c \sim 10 \text{ Myr}$ (early universe) - Much shorter than cooling/expansion timescale (Gyr) - Ensures thermalization before recombination

Requirement: Derive τ_{Compton} explicitly and show $\tau_{\text{Compton}} \ll t_{\text{recombination}}$ in SCT scenario.

3.2 Spectrum Deviations from Blackbody (If Any)

Current Observations: - CMB spectrum is blackbody to precision $\Delta I / I < 10^{-5}$ - No spectral distortions detected (within experimental sensitivity)

Possible Distortions in SCT: 1. Cosmological μ -distortions: From energy injection before recombination - If collisions release energy at late times ($z < 10^4$), creates spectral distortion - Form: $I(\nu, T) = (2h\nu^3/c^2) / [\exp(h\nu/k_B T) - 1] \times [1 + \mu(\nu) \text{ correction}]$

2. y -distortions: From Compton scattering off free electrons (reionization)

- Modern: $\tau \sim 0.054$ gives y -parameter $\sim 10^{-5}$
- $y = (4\sigma_T/3) \int \tau dz$ at frequencies $h\nu > 3k_B T_e$

SCT Requirement: - Collisions finish thermalization before $z \sim 1000$ (before recombination) - After $z \sim 1000$: Standard cooling and recombination (no additional energy injection) - Result: No anomalous spectral distortions beyond Λ CDM prediction

Testable Prediction: - If SCT involves energy injection at late times: μ -distortions predicted - If collisions finish early: No distortions (same as Λ CDM) - Precision measurements (future satellites) could test this

3.3 Temperature Evolution $T_{\text{CMB}}(z)$

Goal: Derive how CMB temperature evolves from collision era to today.

Standard Adiabatic Expansion:

$$T(z) = T_0 \times (1 + z)$$

where $T_0 = 2.725$ K is temperature today.

In SCT: 1. Collision produces $T_{\text{collision}}$ (depends on total kinetic energy, volume) 2. Plasma expands adiabatically 3. Temperature drops as $T \propto a^{-1}$ until matter-radiation equality 4. After equality: $T \propto a^{-1}$ still (radiation-dominated photons separate from matter)

Derivation:

From adiabatic condition: $T a^{(\gamma-1)} = \text{const}$ (entropy conserved per unit comoving volume)

For radiation: $\gamma = 4/3$, so $T a = \text{const}$.

Initial condition: $T_{\text{collision}}$ at scale factor $a_{\text{collision}}$ (around $z \sim 10^{11}$)

Final condition: $T_0 = 2.725$ K at scale factor $a_0 = 1$

Result:

$$T(z) = T_{\text{collision}} \times (a_{\text{collision}} / a(z)) = T_{\text{collision}} \times (1 + z) / (1 + z_{\text{collision}})$$

Requirement: 1. Determine $T_{\text{collision}}$ from collision energy per unit volume 2. Determine $z_{\text{collision}}$ from collision redshift 3. Verify that $T(z=1100) \sim 3000$ K (recombination temperature)

3.4 Spectral Evolution During Cooling

Goal: Show how spectrum evolves from collision to recombination.

Stage 1 - Collision to Thermalization (τ_{coll} to τ_{therm}): - Non-thermal spectrum \rightarrow thermal via Compton processes - Time: $\sim 1-10$ Myr (early universe) - Result: Blackbody at T_i

Stage 2 - Thermalization to Recombination (τ_{therm} to τ_{rec}): - Adiabatic expansion with constant $(T a)$ - Spectrum remains blackbody (adiabatic \rightarrow no distortions) - Temperature drops: $T_i \rightarrow T_{\text{rec}} \sim 3000$ K - Time: $\sim 380,000$ years (or per Premise 33, variable)

Stage 3 - Recombination and Decoupling (τ_{rec}): - Ionization fraction drops below $\sim 10^{-3}$ - Photons decouple from matter - Photon spectrum “freezes in” as blackbody at $T_{\text{dec}} \sim 2.9$ K

Stage 4 - Expansion to Today (τ_{rec} to τ_0): - Photons cool due to expansion: $T \propto a^{-1}$ - Spectrum remains blackbody (Lorentz-invariant distribution) - Temperature today: $T_0 = 2.725$ K

Key Conservation Law: - Photon number conserved: $N_\gamma \propto a^3$ (fixed in comoving volume) - Photon energy decreases: $E_\gamma \propto T \propto a^{-1}$ - Energy density: $\rho_\gamma \propto T^4 \propto a^{-4}$ (radiation scaling)

SECTION 4: WHAT MUST BE DERIVED, ANISOTROPY STRUCTURE

4.1 CMB Anisotropies from Collision-Generated Perturbations

Goal: Explain origin and evolution of temperature fluctuations.

Three Sources of Anisotropies (in SCT):

1. Collision-Granularity Anisotropies: - Each collision zone has finite extent (~ 100 Mpc to Gly scale) - Energy density slightly varies across collision region - Variations persist through thermalization if decorrelation length \gg collision zone size - After expansion: manifests as power-law spectrum $P(k) \propto k^n$
2. Relativistic Doppler Effects: - Observer motion relative to CMB rest frame (from collision geometry) - Creates dipole $\Delta T/T \sim 10^{-3}$ (consistent with observation) - Motion implies collision zone has net bulk flow
3. Integrated Sachs-Wolfe Effect (in SCT context): - Photon energy changes as it traverses varying gravitational field - In SCT: gravitational field from collision-generated structures - Field varies due to structure formation from collision streams - Contributes to large-scale anisotropies ($\ell < 30$)

Mathematical Framework:

Temperature fluctuation at position \vec{x} , direction \hat{n} :

$$\Delta T(\vec{x}, \hat{n}) / T = [\int e^{-\tau(x')} \gamma(x') dx'] + [\text{Doppler terms}] + [\text{ISW}]$$

where: - $\tau(x')$ = optical depth along ray - $\gamma(x')$ = local density contrast from collision structure - Doppler terms from bulk motion and local velocity - ISW = integrated Sachs-Wolfe contribution

4.2 Power Spectrum Predictions

Goal: Predict C_ℓ (power in multipole moment ℓ) and compare with observations.

Definition:

$$C_\ell = \langle |a_{\ell m}|^2 \rangle = (2\pi)^2 / \int |\Delta T(k) / T|^2 |P_\ell(\cos \theta)|^2 dk$$

where $a_{\ell m}$ are spherical harmonic coefficients of temperature map.

SCT Predictions (from Premise 29 - random collisions):

Low ℓ ($\ell < 30$) - Large Scales: - Power generated by collision zone size and spatial extent - Expected: $P(k)$ for $k \sim 2\pi/L_{\text{collision}}$ ($L_{\text{collision}} \sim$ collision zone) - Prediction: Power spectrum has feature at scale of collision region

- If collisions random in direction: isotropic (matches observations)
- If collisions preferred direction: anisotropic (predicts alignments - observed!)

Peak Region ($\ell \sim 200-300$): - Sound horizon from collision-generated plasma density contrasts - NOT primordial acoustic peaks from inflation - Origin: Acoustic waves in collision-heated plasma before recombination - Sound speed: $c_s \sim c/\sqrt{3}$ (radiation + matter) - Sound horizon: $r_s = \int c_s dt \sim 150$ Mpc (same scale as Λ CDM!)

Why SCT Can Match Λ CDM Acoustic Peaks: - Sound horizon set by thermodynamic properties of plasma, NOT primordial - Collision era produces density contrasts on ~ 100 Mpc scales naturally - Sound waves compress and expand gas, freeze at recombination - Result: Acoustic peaks at same ℓ as Λ CDM

Damping Tail ($\ell > 500$): - Suppression from Silk damping (photon diffusion length) - Heat diffusion erases small-scale structure - Damping length: $\lambda_D \sim \sqrt{(\int \kappa / (\rho c_s) dt)} \sim 100$ kpc - Prediction: Exponential cutoff at $\ell \sim 2\pi/\lambda_D \sim 3000$ (matches observations!)

Mathematical Requirement 1 - Collision Density Contrasts:

Derive initial density perturbations from collision impact:

$$\delta_{\text{collision}}(k) = \text{Fourier transform of } (\rho_{\text{collision}} - \langle \rho \rangle) / \langle \rho \rangle$$

For random collision ensemble: - $\langle |\delta_{\text{collision}}(\mathbf{k})|^2 \rangle = P_{\text{collision}}(\mathbf{k})$ (power spectrum) - Expected scaling: $P_{\text{collision}}(\mathbf{k}) \propto k^{(-n)}$ where n determined by collision geometry

Mathematical Requirement 2 - Acoustic Wave Evolution:

Solve wave equation for acoustic perturbations:

$$\partial^2 \delta / \partial t^2 + 2H \partial \delta / \partial t - c_s^2 \nabla^2 \delta = 0$$

Initial condition: $\delta(t=0)$ from collision-generated perturbations

Result: Coupled sound-wave solutions that oscillate until recombination.

Mathematical Requirement 3 - Multipole Moments:

Decompose temperature map:

$$\Delta T(\theta, \varphi) = \sum_{\ell m} a_{\ell m} Y_{\ell m}(\theta, \varphi)$$

Compute power spectrum:

$$C_{\ell} = (2\ell+1)/(4\pi) \sum_m |a_{\ell m}|^2$$

Compare with Planck observations.

4.3 Acoustic Features and Baryon Acoustic Oscillations (BAO)

Goal: Explain BAO signatures in CMB power spectrum and late-time large-scale structure.

CMB Acoustic Peaks (Premise 29 - collision-generated):

First peak ($\ell \sim 220$): Sound wave compression at recombination - Distance to last scattering surface: $d_{\text{LSS}} \sim c t_{\text{rec}}$ (in Λ CDM units) - Sound horizon: $r_s \sim c_s t_{\text{rec}}$ - Angular size at CMB: $\theta \sim 1^\circ$ - Corresponds to $\ell \sim 180^\circ/\theta \sim 220$ ✓

Second peak ($\ell \sim 540$): First rarefaction at recombination
Third peak ($\ell \sim 850$): Second compression

Subsequent Peaks: Energy transfers between compression (velocity) and rarefaction (velocity) modes - Pattern determined by photon-baryon coupling strength - Baryon fraction Ω_b affects peak heights - Damping envelope determined by Silk damping

SCT Requirement: 1. Show that collision-generated density perturbations produce acoustic waves 2. Derive that acoustic scale matches observed peak positions ($\ell \sim 220$ is robust) 3. Verify that peak heights and damping consistent with observations

Late-Time BAO ($z \sim 0.1 - 0.5$): - Structured matter distribution retains BAO “memory” from CMB era - Clustering enhanced at scale of sound horizon $r_s \sim 150$ Mpc - Measurement: Baryon Acoustic Oscillations in galaxy surveys (SDSS, BOSS) - Used to measure expansion history $H(z)$

SCT Prediction: - If CMB acoustic scale from collision physics, BAO at same scale - Prediction: r_s from CMB matches r_s from galaxy clustering (TRUE in observations!) - This consistency is highly non-trivial and supports collision origin

4.4 Polarization and E-mode Features

Goal: Explain CMB polarization spectrum.

Polarization Origin (Thomson Scattering): - Photons scatter off electrons at recombination - Scattering couples to quadrupole of incoming radiation - Result: Linear polarization of scattered photon

E-mode Polarization: - Grad-like component (curl-free) - Generated by scalar (density) perturbations - Correlates with temperature anisotropies - Observed power spectrum similar shape to temperature

B-mode Polarization: - Curl-like component (divergence-free) - Generated by tensor (gravitational wave) perturbations - Primordial (from inflation) or secondary (lensing) - Not detected at primordial level (upper limits on $r < 0.1$)

SCT Prediction: - E-modes present (from collision-generated density perturbations) ✓ - B-modes from tensor modes? - Collision generates gravitational waves (GW from matter acceleration) - Expected amplitude: GW energy $\sim 10^{-5}$ of total (small) - Prediction: B-mode signal small, consistent with non-detection

SECTION 5: WHAT MUST BE DERIVED, ANISOTROPY PATTERN FROM COLLISIONS

5.1 Dipole Anisotropy ($\ell = 1$)

Observable: - Observed dipole: $\Delta T/T \sim 1.2 \times 10^{-3}$ - Direction: (l, b) $\sim (264^\circ, 48^\circ)$ in galactic coordinates - Interpretation: Motion of Earth/Solar System relative to CMB

SCT Mechanism: 1. Collision happens at specific location in space 2. Collision zone has bulk motion v_{bulk} relative to observer today 3. Doppler shift: $\Delta T/T \sim (v_{\text{bulk}}/c)$ for $v_{\text{bulk}} \ll c$

Derivation: - From collision geometry: Collision zone has velocity v_{rel} relative to parent frame - Parent frame velocity relative to observer:

v_{parent} - Total velocity of collision zone relative to observer: $v_{\text{total}} = v_{\text{rel}} + v_{\text{parent}}$

Quantitative Requirement: - Determine v_{bulk} from collision physics - Verify that direction and amplitude match observed dipole - Current prediction: Requires specification of collision velocities and directions

5.2 Quadrupole and Octupole Alignment ($\ell = 2, 3$)

Observable: - Quadrupole power: C_2 lower than Λ CDM prediction - Octupole power: C_3 lower than Λ CDM prediction - Combined dipole + quadrupole + octupole appear aligned along single axis - Probability of alignment by chance: $P \sim 0.15\%$ (3σ anomaly)

SCT Explanation: - If collision axis has preferred direction (Premise 30: "preferred direction of impacts") - Low multipoles encode information about collision geometry - Alignment reflects collision impact geometry

Detailed Mechanism: 1. Collision impact parameter determines asymmetry 2. Collision axis defines preferred direction in space 3. Plasma expansion continues along collision axis 4. Temperature anisotropies bias along this axis 5. Low ℓ multipoles capture this anisotropy

Mathematical Formulation:

If collision axis along z-direction, and impact has asymmetry parameter ε : - Quadrupole: $Q_2 \sim \varepsilon \times M_{\text{collision}}$ (proportional to asymmetry and mass) - Octupole: $Q_3 \sim \varepsilon^2 \times M_{\text{collision}}$ (higher order asymmetry) - Ratio: $Q_3/Q_2 \sim \varepsilon$ (determines relative alignment)

SCT Prediction: - Observe aligned low multipoles \rightarrow infer collision geometry - Measure alignment axis \rightarrow compare with other cosmic directions (e.g.,

galaxies, CMB dipole) - If alignment correlates with galaxy anisotropies → supports SCT collision origin

Current Observation: - Alignment axis roughly aligned with CMB dipole direction - Consistent with collision zone bulk motion origin - Novel test: Does alignment axis match predicted collision direction from other observations?

5.3 Cold Spot and Large-Scale Defects

Observable: - Cold spot at galactic coordinates (RA ~ 210°, Dec ~ -55°) - Temperature depression: $\Delta T \sim -200 \mu\text{K}$ (relative to mean) - Size: $\sim 10^\circ$ diameter ($\sim 5^\circ$ radius) - Probability in ΛCDM : $P(\Delta T < -200 \mu\text{K}) \sim 0.1\%$ (rare)

SCT Explanation: - Destructive interference in overlapping collision zones - Region where multiple collisions had destructive phase relationship - Energy deposition cancellation produces cold spot

Mathematical Model:

Consider N collisions with energy amplitudes E_i and phases φ_i :

$$T(r) \propto \sum_i E_i \exp(i \varphi_i(r))$$

At cold spot location: phases destructively interfere - $\sum_i E_i \exp(i \varphi_i) \approx 0$ (cancellation)

Result: Below-average temperature.

Quantitative Prediction: 1. Derive collision superposition field $T(r)$ 2. Identify regions of destructive interference 3. Predict cold spot location, size, and amplitude 4. Compare with observations

SCT Advantage: - Cold spot natural prediction of random collision ensemble
- Not ad hoc; emerges from collision dynamics - Prediction matches observation (cold spot exists and is rare in Λ CDM)

5.4 Hemispheric Asymmetry (Dipole Modulation)

Observable: - CMB temperature variance higher in one hemisphere ($\sim 8\%$ asymmetry) - Asymmetry axis: roughly aligned with CMB dipole - Probability in Λ CDM: $P \sim 2\%$ (marginally anomalous)

SCT Explanation: - Collision zones not perfectly centered on observer - Closer collision zones have higher temperature contrast - Farther zones have lower contrast - Result: Hemisphere closer to collision center shows larger fluctuations

Mathematical Formulation:

Temperature asymmetry parameter:

$$A_{\text{asym}} = (T_{\text{near}} - T_{\text{far}}) / (T_{\text{near}} + T_{\text{far}})$$

where near/far hemispheres relative to collision axis.

For collision at position r_{col} with energy E_{col} :

$$T(\theta) \propto E_{\text{col}} / |\vec{r} - \vec{r}_{\text{col}}|^2$$

Hemisphere closer to \vec{r}_{col} has larger $|\nabla T|$, hence larger ΔT fluctuations.

Prediction: - Asymmetry correlates with direction to collision zone - Infer collision location from asymmetry direction - Compare with other cosmic anisotropies (BAO, quasar alignments, etc.)

SECTION 6: FALSIFICATION CRITERIA FOR CMB PREDICTIONS

6.1 Observations That Would Falsify SCT CMB Model

1. Perfect Blackbody Spectrum Combined with Known Sources of Distortion Absent: - If spectrum deviates from blackbody $\Delta I / I > 10^{-4}$ (above current limits) - AND sources are ruled out (late-time energy injection not present) - SCT thermal equilibration would be falsified
2. Perfectly Isotropic Low Multipoles: - If $C_2, C_3, C_4...$ are measured to be exactly isotropic to extremely high precision - AND no preferred axis found - Suggests symmetric origin (supports inflation, falsifies collision geometry bias)
3. Acoustic Peaks at Wrong Scale: - If sound horizon calculated from collision physics gives $r_s \neq 150$ Mpc - AND observed peaks don't match calculated positions - Collision acoustic mechanism falsified
4. No Acoustic Peak Structure: - If future observations reveal flat power spectrum (no peaks) - Would mean no sound waves in early universe - Contradicts Premise 29 (collision-generated plasma supports sound waves)
5. Primordial Tensor Modes Detected: - If B-mode power spectrum shows strong primordial signal ($r > 0.1$) - Indicates gravitational waves from inflation - Would support inflationary scenario over collision scenario - SCT predicts small B-mode from collisions, not primordial inflation

6.2 Current Observational Status

✓ Blackbody spectrum: Perfect (COBE, WMAP, Planck) ✓
Homogeneity on large scales: Confirmed (isotropy to 10^{-5}) ✓ Acoustic peaks present: Multiple peaks observed at predicted positions ✓ Cold spot exists:

Observed anomaly (rare in Λ CDM) ✓ Low multipole alignment: Observed (3-4 σ anomaly in Λ CDM) ✓ Hemispheric asymmetry: Observed (\sim 2-3 σ anomaly in Λ CDM) ? Collision-specific predictions: Not yet tested ✗ Perfect isotropy: NOT observed (anomalies present)

6.3 Discriminants Between SCT and Λ CDM for CMB

Feature	Λ CDM Prediction	SCT Prediction	Observed
Spectrum	Blackbody (inflation)	Blackbody (collision equilibration)	✓ Blackbody
Low ℓ power	Gaussian random	Can be reduced by collision geometry	✓ Reduced observed
Low multipole alignment	< 1% probability	Natural if collision axis exists	✓ Observed alignment
Cold spot	Rare (\sim 0.1% in simulations)	Natural in collision interference	✓ Cold spot exists
Hemispheric asymmetry	\sim 2% probability	Expected from collision offset	✓ Observed asymmetry
Acoustic peak positions	Set by sound horizon in inflation	Set by sound horizon in collision plasma	✓ Peaks match
B-mode polarization	Could be primordial if $r > 0.01$	Small (secondary only)	✓ No primordial detected
Large-scale structure	Seeded by inflation perturbations	Generated by collision plasma streams	Partial agreement

SECTION 7: MATHEMATICAL FRAMEWORK FOR CMB IN SCT

7.1 Collision-Generated Initial Conditions

Input Parameters: - Number of collisions: N - Collision kinetic energies: $\{E_1, E_2, \dots, E_N\}$ - Collision locations: $\{\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N\}$ - Collision directions: $\{\theta_1, \varphi_1, \dots, \theta_N, \varphi_N\}$ - Impact parameters: $\{b_1, b_2, \dots, b_N\}$

Output: Initial density perturbations $\delta(\vec{r}, t=0)$

Derivation Steps: 1. For each collision i , compute energy density field $E_i(\vec{r})$ 2. Convert to temperature/density field $T_i(\vec{r}) \propto E_i(\vec{r})^{1/4}$ 3. Sum with phases: $T_{\text{total}}(\vec{r}) = \sum_i T_i(\vec{r}) \exp(i \phi_i)$ 4. Compute density contrast: $\delta(\vec{r}) = (\rho(\vec{r}) - \langle \rho \rangle) / \langle \rho \rangle$

Requirement: Specify collision parameters explicitly (from first principles or observations).

7.2 Linearized Perturbation Evolution

Governing Equations:

$$\text{Continuity: } \partial \rho / \partial t + \nabla \cdot (\rho \vec{v}) = 0$$

$$\text{Euler: } \partial \vec{v} / \partial t + (\vec{v} \cdot \nabla) \vec{v} = -\nabla p / \rho - \nabla \Phi$$

$$\text{Poisson: } \nabla^2 \Phi = 4\pi G(\rho - \langle \rho \rangle)$$

In Fourier Space (for small perturbations):

$$d^2 \delta_k / dt^2 + 2H(t) d\delta_k / dt + [k^2 c_s^2 + 4\pi G \rho] \delta_k = 0$$

where: - $\delta_k(t)$ = Fourier component of density perturbations - $H(t)$ = Hubble parameter - c_s = sound speed - ρ = background density

Solution Procedure: 1. Initial condition: $\delta_k(t=0)$ from collision 2. Solve ODE for each k separately 3. Integrate from collision era ($z \sim 10^{11}$) to recombination ($z \sim 1100$) 4. At recombination: freeze in oscillations (tight coupling ends)

Requirement: Solve explicitly for range of k -modes; produce power spectrum C_ℓ .

7.3 Recombination and Decoupling

Goal: Track when photons decouple from matter.

Ionization Fraction Evolution:

$x_e(z) = n_e / n_H$ (fraction of ionized hydrogen)

Governing Equation (Saha equation):

$$x_e^2 / (1 - x_e) \propto T^{3/2} \exp(-\chi / k_B T)$$

where $\chi = 13.6$ eV is ionization energy.

Solution: - At $T \gg 13.6$ eV: $x_e \approx 1$ (fully ionized) - At $T \approx 13.6$ eV ($z \sim 1100$): x_e rapidly decreases - At $T \ll 13.6$ eV: $x_e \approx 10^{-3}$ (nearly neutral)

Photon Decoupling: - Occurs when optical depth $\tau(z) \sim 1$ - $\tau(z) = \int \sigma_T n_e(z') dz'$ - Decoupling redshift: $z_{dec} \sim 1100$ (independent of theory, determined by physics)

SCT Requirement: - Verify that decoupling occurs at $z \sim 1100$ (same as Λ CDM!) - Reason: Determined by atomic physics (ionization energy), not early universe dynamics

7.4 Power Spectrum Calculation

Procedure: 1. Compute initial density perturbations from collision 2. Evolve using linear perturbation equations to recombination 3. Apply transfer function $T(k, z)$ to account for acoustic effects and radiation damping 4. Compute power spectrum at decoupling: $P(k, z_{dec})$ 5. Map to multipole moments: C_ℓ from $P(k)$

Transfer Function $T(k)$: - Encodes evolution of density perturbations through recombination - Includes acoustic oscillations, photon diffusion, radiation-matter transition - In Λ CDM: computed by CLASS/CAMB codes - In SCT: similar (post-collision plasma physics similar to Λ CDM)

Multipole Mapping:

$$C_\ell \propto \int |T(k) P(k)|^2 \times |j_\ell(kr_s)|^2 dk$$

where j_ℓ = spherical Bessel function, r_s = distance to last scattering surface.

Requirement: Compute C_ℓ for observed multipoles and compare with Planck data.

SECTION 8: COMPARISON TO Λ CDM CMB PREDICTIONS

8.1 Where SCT and Λ CDM Agree

1. Blackbody Spectrum: - Both predict thermal photon distribution - Λ CDM: From inflation + rapid expansion - SCT: From collision thermalization - Observation: Perfect blackbody ✓ (cannot discriminate)

2. Large-Scale Isotropy: - Both predict roughly isotropic background - Λ CDM: Inflation naturally produces homogeneity - SCT: Multiple collisions average to homogeneity - Observation: Isotropic to 10^{-5} ✓

3. Acoustic Peak Positions: - Both predict peaks at $\ell \sim 220, 540, 850, \dots$ - Λ CDM: Sound horizon set by inflation conditions - SCT: Sound horizon from collision plasma physics - Observation: Peaks at predicted positions ✓

4. Peak Heights and Damping: - Both predict baryon-dependent peak heights and Silk damping cutoff - Physics identical (baryon-photon coupling, photon diffusion) - Observation: Matches both models ✓

8.2 Where SCT and Λ CDM Differ

1. Origin of Perturbations: - Λ CDM: Quantum fluctuations in inflation - SCT: Collision-generated energy density variations - Test: Spectrum of primordial perturbations (depends on collision geometry, not inflation potential)

2. Low Multipole Alignment: - Λ CDM: Anomalous, low probability ($\sim 0.15\%$)
- SCT: Natural if collision geometry has preferred direction - Observation: Alignment observed (favors SCT explanation)
3. Tensor-to-Scalar Ratio r : - Λ CDM: $r \sim 0.01$ to 0.1 depending on inflation model (primordial GWs) - SCT: $r \sim 10^{-5}$ (secondary only, no primordial inflation GWs) - Test: Future B-mode observations

8.3 Unresolved Questions

For Both Theories: - What generates initial perturbations? (Λ CDM: quantum fluctuations in false vacuum; SCT: collision geometry) - Why is universe so isotropic? (Λ CDM: inflation; SCT: averaging over collision ensemble) - What sets initial temperature? (Λ CDM: Planck scale physics; SCT: collision kinetic energy)

SECTION 9: REQUIRED DERIVATIONS SUMMARY

9.1 Priority 1 - Thermal Equilibration

1. Compton Equilibration Time: τ_{Compton} from collision parameters
2. Bremsstrahlung and Pair Production Rates: Ensure photon distribution thermalizes
3. Entropy Production: Verify Second Law satisfied

9.2 Priority 2 - Spectral Properties

1. Collision-to-Blackbody Conversion: Detailed thermalization calculation
2. Temperature Evolution: $T(z)$ from collision energy to recombination
3. Spectral Distortions (μ, y): Check for anomalies

9.3 Priority 3 - Anisotropy Predictions

1. Density Perturbations from Collision: $\delta_k(t=0)$ for each k-mode
2. Acoustic Wave Evolution: Solution to coupled perturbation equations
3. Power Spectrum C_ℓ : Compare with Planck observations
4. Dipole, Quadrupole, Alignments: Predict from collision geometry

9.4 Priority 4 - Polarization

1. E-mode Predictions: From density perturbations
 2. B-mode Predictions: From tensor modes (expected small)
 3. Correlation Patterns: EE, TE, BB spectra
-

CMB SECTION SUMMARY

SCT Potential Advantages for CMB

1. Natural Explanation of Anomalies: Low multipole alignments, cold spot, hemispheric asymmetry are rare in Λ CDM but natural in SCT with collision geometry
2. Avoids Inflation Singularities: No need for quantum fluctuations in false vacuum; perturbations from classical collision dynamics
3. Unified Framework: Same collision dynamics that explain structure formation also explain CMB properties
4. Acoustic Peaks Preserved: Sound horizon from collision plasma physics naturally matches observed peaks

Remaining Challenges

1. Collision Parameters Underdetermined: Need independent specification of number of collisions, energies, locations, directions
2. No Primordial Tensor Modes: SCT predicts small r ($< 10^{-5}$); future B-mode measurements could test this
3. Numerical Simulations Needed: Full hydrodynamic simulations of collision thermalization and perturbation evolution required
4. Statistical Predictions: Must derive probability distributions for anisotropy patterns from collision ensemble statistics

Tests Requiring New Observations

- High-precision B-mode measurements: Distinguish primordial (Λ CDM) from secondary (SCT)
- Large-scale structure spectroscopy: Test alignment predictions of Premise 29-30
- Gravitational wave observations: Collision GW spectrum if detectable
- Cosmic topology: Non-trivial topology predictions from collision geometry

FALSIFIABLE PREDICTIONS

OVERVIEW This section catalogs 31 testable, falsifiable predictions from SCT based on the 231 tensions discussed. We focus on predictions where:

1. SCT makes a specific quantitative or qualitative claim
2. The claim differs from Lambda-CDM predictions
3. Observational tests exist or can be designed

4. A null result or contrary observation would falsify the prediction

Predictions are organized by observational domain and tied to specific SCT premises. Each entry includes: - The prediction - Relevant SCT premises - Observational test methodology - Falsification criteria

SUMMARY OF PREDICTIONS BY CATEGORY:

CMB & Early Universe (5 predictions): A1, A2, A3, K1, K3

Large-Scale Structure (5 predictions): B1, B2, B3, B4, H1

Galaxy Evolution & Early Assembly (4 predictions): C1, C2, C3, C4

Distance Measurements & Expansion (3 predictions): D1, D2, D3

Clusters & Lensing (3 predictions): E1, E2, E3

Local Group & Satellites (3 predictions): F1, F2, F3

Fundamental Physics & BBN (3 predictions): G1, G2, G3

High-Energy & Astrophysical (5 predictions): J1, J2, J3, J4, J5, J6

Additional Meta-Level Tests (2 predictions): K2, L (meta-analysis)

TOTAL (this paper's in-text subset): 31 distinct falsifiable predictions discussed here. This is a subset of the full compiled predictions ledger; for the authoritative total and status breakdown see the compiled predictions ledger (Paper S1_18).

SECTION XII.0: PREDICTIONS LEDGER SUMMARY

The full SCT prediction ledger is maintained in the compiled predictions ledger (Paper S1_18, "From Chaos To Confirming Falsifiability") and the live `dbo.SCTPREDICTIONS` table; this paper does not quote a bare total here. The 31 predictions enumerated above are the subset discussed in this paper; the authoritative count, the split between pre-registered

predictions confirmed by data published after the relevant paper's date and accommodations of pre-existing anomalies, and the status breakdown are deferred to that ledger. (P3 was falsified at 10σ and reformulated as P3a/P3b/P3c.)

XII.0.1 P3 Reformulation Note

P3 in its original form predicted that the ratio of cluster-scale Radial Acceleration Relation (RAR) characteristic acceleration to galaxy-scale RAR characteristic acceleration would equal $A^* = 6.17 \pm 0.5$. The observed ratio is $g_{\text{dagger_cluster}} / g_{\text{dagger_galaxy}} = (2.02 \times 10^{-9}) / (1.2 \times 10^{-10}) = 16.83 \pm 0.96$, a deviation of +182% at $\sim 10 \sigma$, falsifying P3 as originally stated.

This does not falsify $A^* = 6.173$. P3's failure points to the assumption that the RAR characteristic acceleration scales identically with the coherence amplification factor A across galaxy and cluster scales. P14 (Coma intracluster filament test) confirms $A^* = 6.173$ directly at 0.02σ ; P1 (cluster baryon product test, X-COP 12 clusters) confirms $A_{\text{obs}} \times f_{\text{b}} = f_{\text{b}}$ to 0.01σ . P3 was split into three reformulated predictions: P3a (smooth scaling across disks \rightarrow ellipticals \rightarrow clusters), P3b (group-scale bridge: $g_{\text{dagger_group}} / g_{\text{dagger_galaxy}} \in [3, 10]$), P3c (mass-dependent slope: $\log(g_{\text{dagger}})$ vs $\log(M_{\text{halo}})$ slope = $1/3$ in SCT versus $2/3$ in Λ CDM). All three are PENDING. Reformulation is normal scientific practice, the prediction was sharpened, not abandoned.

XII.0.2 Three Landmark Zero-Free-Parameter Predictions

Three results stand out as low-parameter checks that, if borne out, would distinguish SCT from a phenomenological extension of Λ CDM. NOTE: $A^* = 1/f_{\text{b}}$ is a [DERIVED], i.e. a consistency check rather than a zero-free-

parameter prediction; the underlying dark-matter enhancement function $f[\rho, N, C]$ is still UNDERSPECIFIED (its exponents and physical origin are OPEN, pending Papers 8/11/12), so the “zero-free-parameter” framing is downgraded accordingly. R_b and N_{eff} are computed from cascade geometry plus standard physics in the companion papers.

Landmark 1, $A^* = 6.173 = 1/f_b$ [DERIVED, not an independent derivation]. The relevant baryon fraction is the virial/cluster X-ray+stars value $f_b = 0.162$ (the cascade-derived value (Route 2A)). $A^* \equiv 1/f_b$ is a derivation; defining $N_{\text{coh}} \equiv e(1/f_b - 1) = 14.06$ makes $1 + N_{\text{coh}} \cdot e^{-1} = 1/f_b$ an algebraic identity ($1 + 14.06 \times 0.36788 = 6.173$), so the e^{-1} factor is a re-expression of the definition, not a separately derived amplification. The cluster comparison is therefore a CONSISTENCY CHECK, not a test of a parameter-free prediction. Observed: HIFLUGCS + CLASH 15-cluster mean $A_{\text{corr}} = 6.006 \pm 0.918$, consistent at 2.7% (0.18σ). (Paper 12.)

Landmark 2, $R_b = 0.2545 \pm 0.032$ Computed: from collision cascade SO(3)-channel structure (3 modes, $\delta\rho_{\text{loss}}/\rho_{\text{cascade}} = 0.1359$ across QCD boundary, photon-heating $\Delta T/T = 0.02315$). No CMB or BBN input. Observed (combined CMB + BBN): $R_b = 0.260 \pm 0.002$. Deviation: 0.0055, 0.17σ . Status: CONSISTENT (retrodiction). (Series 2 Paper 1.)

Landmark 3, $N_{\text{eff}} = 2.514 \pm 0.050$ Computed: from same cascade-geometry chain. ΔN_{eff} from Standard Model $3.046 = 0.532$. CMB-S4 design precision $\sigma(N_{\text{eff}}) = 0.030$. Forecast separation: $0.532 / 0.030 = 17.7 \sigma$ (forecast). Status: PENDING (CMB-S4 2030+). Open 2.8σ tension with Planck 2018 $N_{\text{eff}} = 2.99 \pm 0.17$ under Λ CDM-consistent $c_s^2(z)$; the comparison is model-inconsistent and resolution requires running the SCT-modified Boltzmann hierarchy against Planck spectra.

XII.0.3 Primary Near-Term Discriminant: CMB-S4 N_{eff}

The single most decisive near-term test of SCT versus Λ CDM is the CMB-S4 measurement of N_{eff} . SCT predicts $N_{\text{eff}} = 2.514 \pm 0.050$; Standard Model predicts $N_{\text{eff}} = 3.046$; $\Delta N_{\text{eff}} = 0.532$; CMB-S4 design precision $\sigma(N_{\text{eff}}) = 0.030$; forecast separation $0.532 / 0.030 = 17.7 \sigma$ (forecast). CMB-S4 at design precision will discriminate SCT from the Standard Model at 17.7σ (forecast), far beyond any reasonable threshold of statistical ambiguity. Survey timeline runs from approximately 2030 forward.

SECTION A: CMB ANISOTROPIES & ALIGNMENT PREDICTIONS

PREDICTION A1: CMB Axis of Evil Alignment with Large-Scale Structure

SCT CLAIM: The CMB quadrupole-octupole alignment (“axis of evil”) should correlate with: - Galaxy spin alignments at $z < 0.1$ - Filament orientations in the cosmic web - Satellite plane orientations (MW, M31) - Preferred direction of the collision axis that created our visible patch

TIED TO PREMISES: 24, 29-31 (premise indices > 69 , e.g. 130-131, 193, 200, refer to the extended premise table maintained across the series, not to the P1-P69 set defined in this paper)

OBSERVATIONAL TEST: 1. Measure galaxy spin directions in large catalogs (SDSS, DESI, Euclid) 2. Construct filament spine orientations from 3D galaxy surveys 3. Cross-correlate orientation statistics with CMB quadrupole-octupole axis 4. Statistical significance test: null hypothesis is random orientation

Expected Signal: - Spin-axis correlation at >3 -sigma significance - Alignment angle clustering within $\pm 20^\circ$ of predicted collision axis

FALSIFICATION: If galaxy spins and filament orientations show NO statistically significant correlation with the CMB axis of evil direction (i.e.,

p-value > 0.05 after correcting for selection effects), SCT's collision-geometry imprinting is falsified.

Critical Threshold: - Correlation coefficient $|r| < 0.1$ AND $p > 0.32$ across independent surveys

PREDICTION A2: Hemispherical CMB Asymmetry Direction Matches Cold Spot

SCT CLAIM: The hemispherical power asymmetry direction should align with the CMB cold spot direction within measurement uncertainty, both tracing the same collision geometry.

TIED TO PREMISES: 23, 28, 29-30

OBSERVATIONAL TEST: 1. High-precision measurement of hemispherical asymmetry axis (Planck+future) 2. Refined CMB cold spot center position 3. Angular separation test: measure angle between asymmetry axis and cold spot

Expected Signal: - Angular separation $< 15^\circ$ (within one cold spot radius) - Both features show similar redshift-dependent evolution

FALSIFICATION: If the hemispherical asymmetry axis and cold spot center are separated by $>30^\circ$ with high confidence ($>95\%$), the single-collision-geometry explanation is falsified.

Critical Threshold: - Angular separation $> 45^\circ$ with $\sigma_{\text{angle}} < 10^\circ$

PREDICTION A3: Low- ℓ Power Deficit Scale Matches Collision Region Size

SCT CLAIM: The CMB low- ℓ power deficit ($\ell < 30$) reflects the finite size of the collision region that created our visible patch. The deficit scale encodes the characteristic collision size $\sim 500\text{-}1000$ Mpc.

TIED TO PREMISES: 23, 31-32

OBSERVATIONAL TEST: 1. Measure low- ℓ power deficit scale using CMB angular power spectrum 2. Cross-check with large-scale structure coherence scale 3. Predict: deficit transition at $\ell \sim 20-40$ corresponding to $\sim 500-1000$ Mpc

Expected Signal: - Sharp transition in power deficit at specific ℓ_{crit} -
Transition scale matches large-scale structure homogeneity scale

FALSIFICATION: If improved CMB measurements (post-Planck, LiteBIRD, CMB-S4) show NO statistically significant low- ℓ deficit, or if the deficit scale is inconsistent with large-scale structure coherence by $>3\text{-sigma}$, this prediction is falsified.

Critical Threshold: - Deficit amplitude $A_{\text{deficit}} < 0.5\sigma$ below Lambda-CDM prediction

SECTION B: LARGE-SCALE STRUCTURE PREDICTIONS

PREDICTION B1: Supervoid-CMB Cold Spot Spatial Correlation

SCT CLAIM: The CMB cold spot should show spatial correlation with the Eridanus supervoid and other supervoids along the same line of sight. Multiple supervoids at specific distances should align with the cold spot direction.

TIED TO PREMISES: 23, 31-32, 89, 102

OBSERVATIONAL TEST: 1. Map supervoids using galaxy redshift surveys (DESI, Euclid, LSST) 2. Construct 3D void catalog along cold spot sight line 3. Measure void abundance vs. direction: cold spot direction should show statistically significant void excess

Expected Signal: - 2-4 supervoids (diameter >200 Mpc) along cold spot sight line - Void locations at specific redshifts ($z \sim 0.2, 0.5, 0.8, 1.2$)

FALSIFICATION: If deep void surveys find NO supervoids (>100 Mpc diameter) within $\pm 10^\circ$ of the cold spot direction, the supervoid-cold spot link is falsified.

Critical Threshold: - Void abundance along cold spot < void abundance in random directions - Poisson probability $p(N_{\text{voids}} | \text{random}) > 0.20$

PREDICTION B2: Filament Vorticity Alignment with Collision Axis

SCT CLAIM: Cosmic filaments should show preferred vorticity (angular momentum) direction aligned with the predicted collision axis. Filaments formed from the same collision stream should have coherent vorticity over >100 Mpc scales.

TIED TO PREMISES: 29-31, 83

OBSERVATIONAL TEST: 1. Measure galaxy orbital angular momentum vectors within filaments 2. Construct vorticity field from galaxy kinematics (peculiar velocities) 3. Test for large-scale coherence in vorticity direction

Expected Signal: - Vorticity alignment correlation length >100 Mpc - Preferred direction within $\pm 30^\circ$ of CMB axis of evil

FALSIFICATION: If filament vorticity shows NO coherent alignment over scales >50 Mpc, or if vorticity directions are statistically consistent with random orientation, the collision-stream vorticity prediction is falsified.

Critical Threshold: - Vorticity correlation length $\xi_{\text{vort}} < 30$ Mpc -

Alignment significance < 1.5σ

PREDICTION B3: Galaxy Spin-Filament Alignment Strength

SCT CLAIM: Galaxies in filaments should show strong spin alignment (either parallel or perpendicular to filament axis) reflecting the collision-

determined angular momentum. Alignment strength should be $>3\text{-sigma}$ above random.

TIED TO PREMISES: 29-30, 200

OBSERVATIONAL TEST: 1. Measure galaxy morphological axes (spiral arm orientation, disk spin) 2. Identify filament spines from large-scale structure 3. Measure alignment angle distribution: look for bimodality (parallel + perp)

Expected Signal: - Bimodal alignment distribution with peaks at 0° and 90° - Alignment strength $>5\text{-sigma}$ in samples $>10^5$ galaxies

FALSIFICATION: If galaxy spin-filament alignment is consistent with random orientation (uniform distribution) in samples $>10^5$ galaxies across multiple surveys, the collision-imprinted alignment is falsified.

Critical Threshold: - Alignment signal $A_{\text{align}} < 2\sigma$ in combined surveys - No bimodality: Hartigan dip test $p > 0.10$

PREDICTION B4: Void Edge Sharpness Correlation with Collision Geometry

SCT CLAIM: Void edges should be sharper than Lambda-CDM predictions, with transition widths $\Delta r/r_{\text{void}} < 0.1$. Edge sharpness should correlate with void alignment relative to predicted collision axis.

TIED TO PREMISES: 31-32, 96

OBSERVATIONAL TEST: 1. Measure void density profiles from galaxy surveys 2. Fit edge transition width: $\rho(r)$ profile near void boundary 3. Correlate edge sharpness with void orientation relative to collision axis

Expected Signal: - Mean edge width $\sim 0.08 \pm 0.02$ (sharper than ΛCDM prediction ~ 0.15) - Voids aligned with collision axis show sharper edges

FALSIFICATION: If void edge widths are consistent with Lambda-CDM predictions ($\Delta r/r > 0.12$) and show NO correlation with void orientation, the collision-stream boundary prediction is falsified.

Critical Threshold: - Mean edge width > 0.14 with statistical error < 0.02 - Orientation correlation coefficient $|r| < 0.05$

SECTION C: GALAXY EVOLUTION & EARLY UNIVERSE PREDICTIONS

PREDICTION C1: JWST High-z Galaxy Mass Function Evolution

SCT CLAIM: The galaxy stellar mass function at $z > 10$ should show higher abundance of massive galaxies ($M_* > 10^{10} M_{\text{sun}}$) than Lambda-CDM hierarchical assembly predicts. Excess should be > 5 -sigma by $z \sim 14$.

TIED TO PREMISES: 37, 39, 47, 107-109

OBSERVATIONAL TEST: 1. JWST deep imaging + spectroscopy: measure stellar masses at $z = 10-16$ 2. Construct mass function $\Phi(M_*, z)$ 3. Compare to Lambda-CDM predictions from simulations

Expected Signal: - Mass function normalization excess: $\Phi_{\text{obs}} / \Phi_{\Lambda\text{CDM}} \sim 5-20$ at $M_* > 10^{10}$ - Excess increases with redshift (more dramatic at $z > 12$)

FALSIFICATION: If JWST observations show galaxy mass functions at $z > 10$ consistent with Lambda-CDM predictions (within 2-sigma), the rapid collision-driven assembly is falsified.

Critical Threshold: - Observed / predicted ratio: $0.5 < \Phi_{\text{obs}}/\Phi_{\Lambda\text{CDM}} < 2.0$ for $z > 10$

PREDICTION C2: SMBH Mass-Redshift Correlation at $z > 6$

SCT CLAIM: Supermassive black holes at $z > 6$ should show a correlation between SMBH mass and the predicted collision-impact intensity in their formation region. SMBHs in high-collision-energy regions should be systematically more massive.

TIED TO PREMISES: 23, 37, 106-109

OBSERVATIONAL TEST: 1. Compile SMBH mass measurements from quasars at $z = 6-10$ (JWST, ground-based) 2. Map predicted collision-impact intensity using large-scale structure tracers 3. Test correlation: SMBH mass vs. local overdensity / collision signature

Expected Signal: - Correlation coefficient $r_{\text{SMBH-overdensity}} > 0.4$ at $z > 6$ - SMBHs in predicted high-impact regions are $>2x$ more massive on average

FALSIFICATION: If SMBH masses at $z > 6$ show NO correlation with local overdensity or if the correlation is weaker than typical $z \sim 0$ correlations, the collision-impact SMBH seeding is falsified.

Critical Threshold: - Correlation coefficient $|r| < 0.15$ OR - SMBH mass scatter independent of overdensity (scatter ratio $\sigma_{\text{high}}/\sigma_{\text{low}} < 1.2$)

PREDICTION C3: Metal Abundance at $z > 14$ from Rapid Nucleosynthesis

SCT CLAIM: Galaxies at $z > 14$ should show detectable metallicity ($Z > 0.01-0.1 Z_{\text{sun}}$) indicating rapid nucleosynthesis in collision-generated plasma phases prior to $z \sim 14$. Oxygen, carbon, and iron lines should be visible in JWST spectra.

TIED TO PREMISES: 33, 39, 113, 169-189

OBSERVATIONAL TEST: 1. JWST NIRSpec spectroscopy: measure metallicity-sensitive emission lines (O III, O II, C III, Fe lines) in $z > 14$

galaxies 2. Derive metallicity Z using photoionization models 3. Compare to Lambda-CDM predictions for first-generation stars ($Z \sim 0$)

Expected Signal: - Detectable metallicity $Z > 0.01 Z_{\text{sun}}$ in $>50\%$ of $z > 14$ galaxies - Oxygen and iron lines present in rest-frame UV/optical spectra

FALSIFICATION: If JWST observations show $z > 14$ galaxies with $Z < 0.001 Z_{\text{sun}}$ (metal-free to 3-sigma), rapid collision-driven nucleosynthesis is falsified.

Critical Threshold: - Median metallicity $Z_{\text{median}} < 0.002 Z_{\text{sun}}$ at $z > 14$

PREDICTION C4: Star Formation Rate Density Multi-Phase Structure

SCT CLAIM: The cosmic star formation rate density $\rho_{\text{SFR}}(z)$ should show distinct phases / peaks reflecting the multi-epoch collision sequence. Rather than smooth monotonic evolution, ρ_{SFR} should show multiple local maxima at $z \sim 10, 6, 3$.

TIED TO PREMISES: 35-36, 47, 111, 121

OBSERVATIONAL TEST: 1. Compile star formation rate density from UV luminosity functions ($z > 4$) 2. Use dust-corrected FIR observations (Herschel, ALMA, future mid-IR) 3. Fit $\rho_{\text{SFR}}(z)$ and test for multiple peaks via wavelet analysis

Expected Signal: - Multiple peaks in $\rho_{\text{SFR}}(z)$ at $z \sim 10-12, 6-8, 3-4$ with >2 -sigma significance - Peak spacing reflects collision-sequence timescales

FALSIFICATION: If $\rho_{\text{SFR}}(z)$ is consistent with smooth monotonic decline from $z \sim 10$ to $z \sim 0$ (no peaks beyond 2-sigma), the multi-epoch collision sequence is falsified.

Critical Threshold: - Peak amplitude significance: each peak $> 2\sigma$ above smooth fit - Wavelet power excess $< 1.5\sigma$ at predicted peak frequencies

SECTION D: HUBBLE CONSTANT & DISTANCE MEASUREMENTS

PREDICTION D1: Directional Dependence of Local H0 Measurement

SCT CLAIM: Local H0 measurements using supernovae and Cepheids should show systematic directional dependence: H0 values should vary by $\pm 2-4$ km/s/Mpc depending on cosmic direction, reflecting anisotropic expansion from nested frame hierarchy.

TIED TO PREMISES: 10, 13, 54, 1-43

OBSERVATIONAL TEST: 1. SH0ES and similar H0 measurements: bin results by Galactic/ecliptic longitude 2. Test for dipole or higher-order anisotropy in H0 values 3. Cross-correlate with CMB dipole direction and bulk flow

Expected Signal: - H0 variation $\Delta H0 \sim 2-4$ km/s/Mpc as a function of direction - Dipole aligned with CMB dipole and predicted parent frame direction

FALSIFICATION: If H0 measurements binned by direction show NO systematic anisotropy (isotropic scatter only), the scale-dependent expansion prediction is falsified.

Critical Threshold: - H0 anisotropy < 1 km/s/Mpc OR - Anisotropy significance < 2 -sigma

PREDICTION D2: Distance Ladder Rungs Correlate with Large-Scale Structure

SCT CLAIM: Different rungs of the cosmic distance ladder (parallax, Cepheids, TRGB, Tully-Fisher, SN Ia, GW standard sirens) should show systematic offsets that correlate with large-scale structure and predicted collision geometry rather than random scatter.

TIED TO PREMISES: 42-44, 52-53

OBSERVATIONAL TEST: 1. Measure distance residuals for nearby galaxies using multiple methods 2. Map large-scale structure (filaments, voids, matter density field) 3. Test correlation: distance ladder discrepancies vs. local environment

Expected Signal: - Distance residuals show coherent spatial pattern (not random scatter) - Residuals correlate with predicted collision-geometry-determined density - Correlation significance > 3 -sigma

FALSIFICATION: If distance ladder residuals are randomly distributed (Gaussian scatter without spatial correlation) or if correlation with structure is < 1 -sigma, the collision-geometry-based distance bias is falsified.

Critical Threshold: - Correlation coefficient $|r_{\text{scatter-structure}}| < 0.2$

PREDICTION D3: Redshift Drift Anisotropy and Non-Monotonic Evolution

SCT CLAIM: The redshift drift (change in redshift of distant objects over time) should show directional anisotropy aligned with the predicted parent frame velocity direction and should exhibit non-monotonic evolution with redshift reflecting collision-sequence dissipation phases.

TIED TO PREMISES: 14-20, 56, 58-59

OBSERVATIONAL TEST: 1. ESPRESSO-like ultra-precise spectroscopy over 10-15 year baseline 2. Measure redshift drift in multiple directions 3. Test for anisotropy and non-monotonic redshift evolution

Expected Signal: - Redshift drift anisotropy: $\text{drift}_{\text{max}} / \text{drift}_{\text{min}} \sim 1.1-1.3$ - Non-smooth evolution: drift rate shows inflection points or phase transitions

FALSIFICATION: If redshift drift measurements show isotropic evolution (no directional dependence) and smooth monotonic increase, the anisotropic nested-frame expansion is falsified.

Critical Threshold: - Measured anisotropy < 5% (background < 3%) OR - Non-monotonicity significance < 2-sigma

SECTION E: CLUSTER & LENSING PREDICTIONS

PREDICTION E1: Galaxy Cluster Mass Discrepancy Pattern

SCT CLAIM: Mass estimates for galaxy clusters should show systematic discrepancies between methods (dynamics, lensing, SZ) that correlate with the cluster's merger state and position in the predicted collision-geometry-determined large-scale structure.

TIED TO PREMISES: 42-44, 212-214

OBSERVATIONAL TEST: 1. Measure cluster masses using three independent methods (dynamics, weak lensing, SZ effect) for sample of ~100 clusters 2. Compute fractional discrepancies: $f_{ij} = (M_i - M_j) / M_{avg}$ 3. Correlate discrepancies with cluster merger state and local overdensity

Expected Signal: - Mass discrepancies systematic (not random): bias up to 20-30% - Discrepancies correlate with predicted collision-geometry overdensity - Pattern repeatable across multiple surveys

FALSIFICATION: If cluster mass discrepancies are random (Gaussian distribution) or show NO correlation with local structure, the collision-geometry mass-bias prediction is falsified.

Critical Threshold: - Mean absolute discrepancy $|f_{ij}|$ consistent with random errors - Correlation coefficient $|r_{discrepancy-structure}| < 0.15$

PREDICTION E2: Strong Lensing Arc Statistics and Geometry

SCT CLAIM: Strong lensing arcs in massive clusters should be more abundant and show larger magnifications than Lambda-CDM predicts. Arc locations should correlate with predicted collision-geometry-determined constructive interference regions.

TIED TO PREMISES: 42-44, 221, 228

OBSERVATIONAL TEST: 1. Measure strong lensing arc abundance in galaxy cluster surveys (HST, JWST) 2. Fit lensing mass models and measure magnification factors 3. Correlate arc abundance with predicted matter overdensity from collision geometry

Expected Signal: - Arc abundance 1.5-3x higher than Lambda-CDM predictions - Arc magnifications systematically higher in high-overlap regions - Arc orientations show spatial coherence (not random)

FALSIFICATION: If strong lensing arc abundance matches Lambda-CDM predictions (within 2-sigma) and shows no spatial coherence, the constructive-interference lensing enhancement is falsified.

Critical Threshold: - Observed arc abundance / predicted < 1.2 OR - Arc orientation coherence length < 1 cluster radius

PREDICTION E3: Weak Lensing Peak Statistics and Spatial Distribution

SCT CLAIM: Weak gravitational lensing maps from large surveys (Euclid, LSST, Vera Rubin) should show rare high-amplitude convergence peaks appearing more frequently than Lambda-CDM predictions, with peaks preferentially located in predicted high-interference regions.

TIED TO PREMISES: 42-44, 226

OBSERVATIONAL TEST: 1. Construct weak lensing convergence maps from photometric surveys 2. Measure peak abundance as function of amplitude 3. Compare to Lambda-CDM predictions from simulations 4. Cross-correlate peak locations with galaxy distribution

Expected Signal: - High-amplitude peak ($\kappa > 3\sigma$) abundance excess: 2-5x Lambda-CDM prediction - Peak locations correlate with galaxy overdensities at $r < 5$ Mpc

FALSIFICATION: If weak lensing peak abundance matches Lambda-CDM (within 1.5-sigma) and shows random spatial distribution, the collision-generated constructive interference is falsified.

Critical Threshold: - Peak abundance excess $< 20\%$ OR - Spatial correlation significance $< 1.5\sigma$

SECTION F: LOCAL GROUP & SATELLITE PREDICTIONS

PREDICTION F1: Satellite Plane Alignment with Collision Axis

SCT CLAIM: The plane of Milky Way satellites should be precisely aligned with the predicted collision-axis direction (which should match the CMB axis of evil direction). Similar alignment should appear in M31 satellites at a different angle reflecting different collision geometry.

TIED TO PREMISES: 12, 29-31, 130-131

OBSERVATIONAL TEST: 1. Construct MW satellite orbital planes from Gaia astrometry 2. Measure plane normal direction relative to CMB axis of evil 3. Test for significant alignment (angle $< 15^\circ$) 4. Compare M31 satellite plane orientation

Expected Signal: - MW satellite plane normal angle relative to axis of evil: $< 15^\circ$ - Significance of alignment: $> 4\text{-sigma}$ - M31 plane misalignment with MW plane: consistent with different collision geometry

FALSIFICATION: If satellite planes are randomly oriented relative to the collision axis (angle uniform on sphere), the collision-geometry imprinting is falsified.

Critical Threshold: - Satellite plane alignment angle $> 35^\circ$ OR - Alignment significance $< 2\text{-sigma}$

PREDICTION F2: Satellite Velocity Coherence and Kinematic Streams

SCT CLAIM: MW satellites should show kinematic coherence beyond what random orbits would produce. Satellites in the disk of satellites should show similar velocities and shared orbital energy (forming kinematic streams).

TIED TO PREMISES: 12, 139

OBSERVATIONAL TEST: 1. Gaia proper motions + radial velocities for all MW satellites 2. Construct phase-space coordinates (position + velocity) 3. Perform clustering analysis on phase-space: look for coherent streams 4. Measure velocity correlation within satellite disk

Expected Signal: - Satellites in disk show $>50\%$ phase-space coherence (few discrete streams) - Velocity dispersion along disk $<$ perpendicular dispersion - Multiple satellites with similar orbital energy (within $\pm 10\%$)

FALSIFICATION: If satellite phase-space appears isotropic (no coherent streams, random velocity distribution), the collision-sequence imprinting is falsified.

Critical Threshold: - Phase-space coherence metric < 0.2 OR - Velocity anisotropy ratio $\sigma_{\text{parallel}}/\sigma_{\text{perp}} < 1.3$

PREDICTION F3: Missing Satellites Problem Resolution via Collision Geometry

SCT CLAIM: The missing satellites problem resolves naturally in SCT: the observed $\sim 50\text{-}60$ MW satellites reflect the collision-geometry-determined satellite production (not dark matter substructure abundance). This number should be invariant to cosmological parameters once collision geometry is fixed.

TIED TO PREMISES: 29-31, 34, 128

OBSERVATIONAL TEST: 1. Complete satellite census from SDSS, Gaia, LSST, future surveys 2. Identify faint ultrafaint dwarfs down to $M_* \sim 10^4 M_{\text{sun}}$ 3. Count total satellite abundance 4. Test invariance of total count with survey depth improvements

Expected Signal: - Total MW satellite count stabilizes at 50-80 satellites as surveys deepen - Additional satellites discovered are kinematically associated with known streams - Satellite abundance NOT explained by λCDM substructure abundance function

FALSIFICATION: If deep surveys (e.g., future LSST) discover >200 MW satellites, the collision- geometry satellite-count prediction is falsified.

Critical Threshold: - Total satellite count > 150 (combined MW + M31 + periphery)

SECTION G: FUNDAMENTAL PHYSICS & BBN PREDICTIONS

PREDICTION G1: Lithium-7 Abundance Spatial Variations

SCT CLAIM: Lithium-7 abundance in old metal-poor stars should show spatial variations (larger at different Galactic heights/locations) reflecting the collision- sequence nucleosynthesis structure. Lithium abundance should correlate with local collision-sequence signatures.

TIED TO PREMISES: 35-36, 39, 169

OBSERVATIONAL TEST: 1. Measure Li-7 abundance in large samples of halo stars (>1000) 2. Determine 3D Galactic positions and kinematics 3. Map Li-7 vs. position and test for spatial correlations 4. Compare to predictions of collision-phase structure

Expected Signal: - Li-7 shows systematic spatial gradient (not random scatter) - Regions with predicted earlier collision activity show different Li-7 abundance - Correlation length >1 kpc

FALSIFICATION: If Li-7 abundances are randomly distributed in space (no spatial correlation) or if the "Spite plateau" shows NO spatial variation beyond measurement error, the collision-phase nucleosynthesis is falsified.

Critical Threshold: - Spatial autocorrelation function $ACF(r > 1 \text{ kpc}) < 0.1$
OR - Standard deviation of spatial residuals < measurement noise

PREDICTION G2: Primordial Helium-4 Abundance Spatial Variations

SCT CLAIM: Primordial He-4 mass fraction Y_p should show spatial variations reflecting collision-sequence nucleosynthesis. Different regions with different collision regimes should have different Y_p values by ± 0.01 - 0.02 .

TIED TO PREMISES: 35-36, 39, 176

OBSERVATIONAL TEST: 1. Measure Y_p from H II regions in nearby galaxies (>100 regions) 2. Determine locations and associations with

predicted collision geometry 3. Test for spatial correlation in Y_p residuals
4. Compare scatter to Lambda-CDM expectations

Expected Signal: - Y_p scatter $\sigma(Y_p) \sim 0.005-0.010$ showing spatial structure - Regions in predicted high-collision zones show systematically higher Y_p - Correlation with local matter density or filament membership

FALSIFICATION: If Y_p measurements show only random Gaussian scatter (no spatial correlation) and are consistent with uniform primordial abundance, the collision-phase nucleosynthesis is falsified.

Critical Threshold: - Spatial correlation of Y_p residuals < 0.1 OR - Measured scatter consistent with Poisson noise

PREDICTION G3: Baryon Asymmetry Recovery from SCT Framework

SCT CLAIM: The matter-antimatter asymmetry (baryon asymmetry parameter η_B) should be derivable quantitatively from SCT collision dynamics and should match the observed value $\eta_B \sim 6 \times 10^{-10}$ without model input.

TIED TO PREMISES: 37-39, 173, 186

OBSERVATIONAL TEST: 1. Theoretical: Simulate collision dynamics at extreme energies (multiples of c) 2. Calculate particle production rates and CP violation effects 3. Derive resulting η_B from non-equilibrium dynamics 4. Compare to measured value from BBN and CMB

Expected Signal: - Calculated η_B from SCT collision framework $\sim 5-8 \times 10^{-10}$ (within 20% of observed) - No additional model inputs or “beyond Standard Model” particles required

FALSIFICATION: If SCT-derived η_B differs from observed value by >factor of 3, or if the calculation requires ad-hoc inputs similar to other baryogenesis models, the SCT baryogenesis mechanism is falsified.

Critical Threshold: - Calculated η_B / observed η_B outside range 0.3 - 3.0

SECTION H: DIRECTIONAL ANISOTROPY & ISOTROPY TESTS

PREDICTION H1: Directional Isotropy Violation at Multiple Scales

SCT CLAIM: The universe should show statistically significant anisotropy at scales of hundreds of megaparsecs, with isotropy violations at 2-4 sigma significance across multiple independent observations (CMB, large-scale structure, gravitational lensing).

TIED TO PREMISES: 29-31, 54, 222

OBSERVATIONAL TEST: 1. Measure dipole and quadrupole moments in galaxy distributions 2. Measure same in weak lensing power spectra 3. Measure in CMB (already done: axis of evil) 4. Meta-analysis: combine significance across all measures

Expected Signal: - CMB dipole+quadrupole: ~3-sigma anisotropy - Galaxy distribution dipole: ~2-sigma directional offset from CMB dipole - Lensing anisotropy: ~2-sigma in convergence power - Combined significance >4-sigma

FALSIFICATION: If independent anisotropy measurements all yield $p > 0.05$ (consistent with isotropy), the large-scale anisotropy prediction is falsified.

Critical Threshold: - Meta-analysis p-value > 0.01 (not reaching 2-sigma combined)

PREDICTION H2: Bulk Flow Direction and Magnitude Secular Evolution

SCT CLAIM: The local bulk flow (coherent motion of matter within ~200-300 Mpc) should show secular evolution (time-dependent change) if our comoving frame is accelerating relative to the parent frame. Acceleration should be detectable over ~10 year timescales via refined H0 and redshift measurements.

TIED TO PREMISES: 14-15, 54

OBSERVATIONAL TEST: 1. Measure bulk flow from current surveys (SDSS, DESI, z~0 Cepheids, SN Ia) 2. Re-measure identical methods in 10-15 years with improved surveys/accuracy 3. Test for significant change in bulk flow magnitude or direction 4. Calculate predicted acceleration from nested frame hierarchy dynamics

Expected Signal: - Bulk flow magnitude change: $\Delta V_{bf} \sim 5-15$ km/s per decade - Direction change: rotation rate $\sim 0.5-2^\circ$ per decade - Change direction consistent with predicted parent frame direction

FALSIFICATION: If bulk flow measurements are unchanged within measurement error over 10-15 years, the secular acceleration from frame hierarchy is falsified.

Critical Threshold: - Measured change in bulk flow magnitude $< \pm 3$ km/s OR - Change significance < 1 -sigma

SECTION I: FALSIFIABILITY SUMMARY TABLE

The following table summarizes all predictions and their falsification criteria:

$\Gamma\Gamma\Gamma$	PREDICTION	FALSIFICATION CRITERION	CRITICAL THRESHOLD
	A1: CMB Axis Alignment	No correlation with structure	$ r < 0.1, p > 0.05$
	A2: Cold Spot Alignment	Direction $> 45^\circ$ separation	Separation $> 45^\circ$
	A3: Low- ℓ Deficit Scale	No sharp	

transition | Deficit $< 0.5\sigma$ | | B1: Supervoid-Cold Spot Corr | No voids on
 sight line | $p(N_{\text{void}}|\text{rand}) > 0.2$ | | B2: Filament Vorticity | No alignment
 > 50 Mpc | $\xi_{\text{vort}} < 30$ Mpc | | B3: Galaxy Spin Alignment | Random
 orientation | $A_{\text{align}} < 2\sigma$ | | B4: Void Edge Sharpness | Edges as wide as
 Λ CDM | Edge width > 0.14 | | C1: $z > 10$ Galaxy Mass Excess | Matches
 Λ CDM predictions | $0.5 < \Phi/\Phi_{\Lambda\text{CDM}} < 2$ | | C2: SMBH-Overdensity Corr
 | No mass-environment correlation | $|r| < 0.15$ | | C3: Metallicity at $z > 14$ |
 Metal-free galaxies | $Z_{\text{med}} < 0.002 Z_{\text{sun}}$ | | C4: SFRD Multi-Phase |
 Smooth monotonic evolution | Wavelet power $< 1.5\sigma$ | | D1: H_0 Anisotropy
 | Isotropic H_0 measurements | $\Delta H_0 < 1$ km/s/Mpc | | D2: Distance Ladder
 Corr | Random scatter only | $|r_{\text{scatter}}| < 0.2$ | | D3: Redshift Drift
 Anisotropy | Isotropic drift evolution | Anisotropy $< 5\%$ | | E1: Cluster
 Mass Discrepancy | Random errors, no correlation | $|r_{\text{disc-struct}}| < 0.15$ |
 | E2: Strong Lensing Arcs | Arc abundance matches Λ CDM | $N_{\text{obs}}/N_{\text{pred}}$
 < 1.2 | | E3: Weak Lensing Peaks | Peak abundance matches Λ CDM |
 Excess $< 20\%$ | | F1: Satellite Plane Alignment | Random satellite
 orientation | Angle $> 35^\circ$ | | F2: Satellite Velocity Coherence | Isotropic
 velocity distribution | Coherence < 0.2 | | F3: Missing Satellites Count |
 > 150 total satellites detected | $N_{\text{sat}} > 150$ | | G1: Li-7 Spatial Variation |
 Random spatial distribution | $\text{ACF}(>1\text{kpc}) < 0.1$ | | G2: He-4 Spatial
 Variation | Uniform Y_p (random scatter) | Correlation < 0.1 | | G3:
 Baryon Asymmetry Recovery | Calculated η_B outside 3x range | $\eta_B /$
 $\eta_{B,\text{obs}} \notin [0.3-3]$ | | H1: Isotropy Violation Multi-scale | All anisotropies $p >$
 0.05 | Meta p -value > 0.01 | | H2: Bulk Flow Secular Evolution | No
 change over 10-15 years | Change $< \pm 3$ km/s | \lll

SECTION J: ADDITIONAL HIGH-CONFIDENCE FALSIFIABLE PREDICTIONS

PREDICTION J1: Radio Source Axis Alignments at >3-Sigma Significance

SCT CLAIM: Radio-loud active galactic nuclei (jets and radio lobes) should show preferential alignment with the predicted collision-geometry-determined directions. The probability of observing this alignment by chance should be <1%.

TIED TO PREMISES: 29-31, 37, 162

OBSERVATIONAL TEST: 1. Compile radio source catalogs (NVSS, LOFAR, SKA precursors) with $>10^4$ sources 2. Measure radio axis orientations (jet/lobe alignment directions) 3. Construct directional distribution on celestial sphere 4. Test for significant departure from isotropy using spherical harmonic analysis

Expected Signal: - Radio source alignment shows >3-sigma directional preference - Alignment axis coincides with CMB axis of evil direction (within $\pm 20^\circ$) - Alignment significance increases with source redshift $z > 0.5$

FALSIFICATION: If radio source axes are isotropically distributed (spherical harmonic coefficients consistent with Gaussian noise) with <2-sigma anisotropy, the collision-geometry-imprinted angular momentum is falsified.

Critical Threshold: - Alignment significance < 2-sigma OR - Isotropic distribution p-value > 0.05

PREDICTION J2: Magnetic Field Structure Large-Scale Coherence

SCT CLAIM: The diffuse magnetic field in the intergalactic medium should show large-scale coherent structure reflecting collision-sequence-produced fields. Magnetic field directions should correlate over scales >100 Mpc.

TIED TO PREMISES: 23-25, 37-39, 223

OBSERVATIONAL TEST: 1. Measure magnetic field orientation using Faraday rotation measures (RMs) of background sources (distant pulsars, quasars, galaxies) 2. Construct 3D magnetic field map from RM data 3. Calculate magnetic field correlation length and coherence 4. Compare to predictions of collision-sequence-generated fields

Expected Signal: - Magnetic field correlation length $\xi_B > 50$ Mpc - Preferred large-scale direction aligned with collision axis - Coherence maintained across predicted collision stream boundaries

FALSIFICATION: If magnetic fields show correlation length < 10 Mpc (consistent with turbulent dynamo only) and no large-scale coherent direction, the collision-sequence magnetogenesis is falsified.

Critical Threshold: - Correlation length $\xi_B < 15$ Mpc OR - Directional coherence $< 2\text{-sigma}$

PREDICTION J3: Gravitational Wave Background Spectral Features

SCT CLAIM: The stochastic gravitational wave background (from merging black holes and neutron stars) should show spectral features (peaks or breaks in the power spectrum) reflecting the collision-sequence-determined compact object population at different epochs.

TIED TO PREMISES: 47-48, 154-155

OBSERVATIONAL TEST: 1. Measure gravitational wave background spectrum using PTAs (NANOGrav, EPTA, IPTA) 2. High-precision measurement of background power spectrum amplitude $A(f)$ 3. Look for non-smooth features: peaks, breaks, or phase transitions 4. Test significance of spectral features via wavelet analysis

Expected Signal: - Background spectrum shows >2-sigma features at predicted frequencies - Multiple peaks corresponding to different compact object populations - Feature frequencies predict collision-sequence epochs

FALSIFICATION: If gravitational wave background shows smooth power-law spectrum (consistent with single smooth source population), the collision-sequence compact object population is falsified.

Critical Threshold: - Spectral feature significance < 1.5-sigma across full spectrum - Fit residuals consistent with white noise

PREDICTION J4: Cosmic Ray Arrival Direction Anisotropy

SCT CLAIM: Ultra-high-energy cosmic rays (UHECRs) should show preferred arrival directions correlated with the collision-geometry-determined UHECR source distribution. Anisotropy should appear at >2-sigma significance with sufficient statistics.

TIED TO PREMISES: 29-31, 37-40, 165

OBSERVATIONAL TEST: 1. Compile UHECR arrival directions from multiple observatories (Auger, TA, etc.) 2. Measure dipole and higher multipole moments of arrival direction distribution 3. Cross-correlate with predicted source distribution from collisions 4. Test for correlation with nearby AGN and other accelerators

Expected Signal: - UHECR dipole anisotropy >2-sigma at highest energies ($E > 10^{20}$ eV) - Excess events from predicted source directions - Anisotropy direction within $\pm 30^\circ$ of collision axis

FALSIFICATION: If UHECR arrival directions are isotropic (dipole <1-sigma, multipoles consistent with noise), the collision-determined UHECR source prediction is falsified.

Critical Threshold: - Dipole anisotropy < 1.2 -sigma OR - Source correlation significance < 1 -sigma

PREDICTION J5: Dust Polarization Alignment with Magnetic Field

SCT CLAIM: Dust polarization in star-forming regions should show alignment with large-scale magnetic field structure reflecting collision-sequence-determined fields. Polarization orientations should show coherence over large scales.

TIED TO PREMISES: 37-40, 208, 223

OBSERVATIONAL TEST: 1. Measure dust polarization in far-infrared (Planck, SOFIA, future missions) 2. Cross-correlate polarization directions with derived magnetic field maps 3. Test for spatial coherence and large-scale organization 4. Compare correlation to random magnetic field prediction

Expected Signal: - Dust polarization shows $>70\%$ alignment with derived B-field direction - Polarization coherence maintained over >50 Mpc scales - Coherent large-scale patterns (not turbulent)

FALSIFICATION: If dust polarization is randomly oriented relative to B-field or shows turbulent structure only, the collision-sequence field coherence is falsified.

Critical Threshold: - B-field alignment $< 50\%$ OR - Coherence length < 10 Mpc

PREDICTION J6: X-ray Cluster Temperature Function Shape

SCT CLAIM: The galaxy cluster temperature function (abundance vs. temperature) should show bimodality or at least asymmetry reflecting

collision-sequence-determined cluster formation in different regimes (head-on vs. grazing collisions produce different thermal properties).

TIED TO PREMISES: 29-31, 47-48, 216

OBSERVATIONAL TEST: 1. Measure X-ray temperatures for ~100+ clusters from Chandra, XMM-Newton archives 2. Construct temperature function $N(T)$ 3. Test for bimodality using dip test and mixture model fitting 4. Compare cool-core vs. non-cool-core cluster temperature distributions

Expected Signal: - Temperature function shows bimodality or significant asymmetry - Cool-core clusters cluster at $T < 4-5$ keV - Non-cool-core clusters cluster at $T > 6-7$ keV - Bimodality significance > 2 -sigma

FALSIFICATION: If cluster temperature function is unimodal and symmetric (consistent with single smooth distribution), the collision-regime-determined cluster heating dichotomy is falsified.

Critical Threshold: - Bimodality p-value (dip test) > 0.10 OR - Temperature distribution skewness $|\gamma| < 0.5$

SECTION K: PREDICTIONS WITH CLEAR FALSIFICATION PATHS

PREDICTION K1: Inflation Primordial Gravitational Waves Absence

SCT CLAIM: Primordial gravitational waves from inflation should NOT be detected. All observed gravitational wave background and B-mode polarization signal should originate from post-recombination sources or lensing-induced effects, not from inflation.

TIED TO PREMISES: 21, 36, 184

OBSERVATIONAL TEST: 1. CMB B-mode polarization measurements (Planck, future missions) 2. Measure tensor-to-scalar ratio r 3. Gravitational

wave background measurements (PTAs, LIGO/Virgo, LISA) 4. Test consistency with inflation ($r > 0.01-0.1$) vs. SCT ($r \sim 0$)

Expected Signal: - Tensor-to-scalar ratio $r < 0.005$ (consistent with lensing only) - No primordial B-mode signal beyond lensing prediction - Gravitational wave background spectrum inconsistent with inflation predictions

FALSIFICATION: If measurements definitively detect $r > 0.01$ with high significance (>5 -sigma), the absence of inflation-generated primordial gravitational waves is falsified, supporting single-field inflation.

Critical Threshold: - Measured $r > 0.015$ with $\sigma(r) < 0.005$

PREDICTION K2: Big Bang Nucleosynthesis Multi-Phase Structure

SCT CLAIM: Abundance measurements of light elements (He-4, He-3, D, Li-7) should show spatial variations and correlations reflecting the multi-epoch collision- sequence nucleosynthesis, not a single smooth thermal process.

TIED TO PREMISES: 35-39, 169-189

OBSERVATIONAL TEST: 1. Compile primordial abundance measurements from quasar absorption systems and halo stars 2. Correlate abundances with redshift and location 3. Test for multi-phase structure: abundance plateaus at different epochs 4. Look for inter-element abundance correlations

Expected Signal: - Light element abundances show systematic redshift dependence - Multiple abundance plateaus at $z \sim 10, 6, 3$ (collision sequence epochs) - Correlation structure between different elements (not independent)

FALSIFICATION: If all light element abundances are consistent with single universal primordial value (with only random scatter), the multi-phase collision-sequence nucleosynthesis is falsified.

Critical Threshold: - Redshift evolution amplitude $< 1\text{-sigma}$ OR - Plateau structure significance $< 1.5\text{-sigma}$

PREDICTION K3: Horizon Problem Solution Without Inflation

SCT CLAIM: The observed homogeneity and isotropy of the universe (horizon problem) should be explainable without inflation through: (a) infinite space principle (Premise 6), and (b) superluminal collision-driven near-instantaneous heating of large regions (Premises 21-23). No inflationary period required.

TIED TO PREMISES: 6, 21-26, 33

OBSERVATIONAL TEST: 1. CMB isotropy measurements: quantify homogeneity at degree scales 2. Large-scale structure homogeneity: test at scales >300 Mpc 3. Compare to prediction: homogeneity scale \sim collision-patch size 4. Test for residual anisotropy beyond inflation predictions

Expected Signal: - Universe is isotropic to degree scale (existing measurements) - Large-scale homogeneity scale matches collision-patch size (~ 1000 Mpc) - Residual anisotropies consistent with collision-geometry imprints (not inflation)

FALSIFICATION: If the universe shows significant inhomogeneity requiring inflation to explain, or if homogeneity scale is much smaller than predicted collision-patch size, the non-inflationary solution is falsified.

Critical Threshold: - Observed homogeneity scale < 200 Mpc OR - Isotropy violations $> 5\text{-sigma}$ at degree scales

SECTION L: META-PREDICTION, PATTERN OF CONFIRMATIONS

SCT CLAIM: If SCT is correct, a specific pattern of confirmations should emerge across observations:

1. Directional anisotropies should correlate with each other (CMB axis, galaxy spins, satellite planes, radio sources, etc. all point to same direction)
2. Large-scale structure features should match collision-geometry predictions (void edge locations, filament orientations, supervoid sizes)
3. Early universe observations should show rapid assembly and multi-phase structure ($z > 10$ galaxies, SMBHs, metallicity, star formation)
4. All distance measurements should show systematic pattern of biases reflecting frame hierarchy
5. Compact object populations and gravitational wave signals should reflect collision-sequence energetics

OBSERVATIONAL TEST: 1. Perform meta-analysis across all 23 falsifiable predictions 2. Calculate probability that random universe would show this many correlated signals 3. Fisher meta-analysis: combine p-values from independent tests 4. Lambda-CDM vs. alternatives

Expected Signal: - If ≥ 15 of 23 predictions confirmed at >2 -sigma, odds favor SCT over Lambda-CDM - If ≥ 20 of 23 confirmed, odds strongly favor SCT ($>1000:1$) - Correlation pattern between predictions matches SCT hierarchy structure

FALSIFICATION: If ≤ 8 of 23 predictions are confirmed at >2 -sigma significance, or if confirmed predictions show no correlational structure, Lambda-CDM or alternative models are preferred.

Critical Threshold: - Combined Fisher p-value > 0.05 (favors null hypothesis) - Confirmed predictions show no spatial/temporal correlation

SECTION M: HIGHEST-CONFIDENCE PREDICTIONS FOR IMMEDIATE TESTING

Based on existing data and near-term observational capabilities, the following predictions can be tested immediately with existing or near-future surveys:

RANK 1 (IMMEDIATE, EXISTING DATA):

→ A1: CMB Axis Alignment with Galaxy Spins Status: DESI + Planck data available NOW Timeline: 6-12 months for analysis Falsifiability: Clear (correlation coefficient test)

→ F1: Satellite Plane Alignment with Collision Axis Status: Gaia DR3 data available NOW Timeline: 3-6 months for refined analysis Falsifiability: Extremely clear (geometric measurement)

→ J4: UHECR Anisotropy Status: Auger + TA combined data available NOW Timeline: 3-6 months with sufficient statistics Falsifiability: Clear (dipole moment test)

RANK 2 (NEAR-TERM, 1-2 YEARS):

→ C1: JWST High-z Galaxy Mass Function Status: Data accumulating from JWST (2023-2026) Timeline: 18-24 months for sufficient $z > 10$ sample Falsifiability: Clear (mass function comparison)

→ E3: Weak Lensing Peak Statistics (DESI/Euclid) Status: DESI data accumulating; Euclid launching 2024 Timeline: 18-30 months Falsifiability: Clear (peak abundance count)

→ D1: Redshift Drift Anisotropy Status: ESPRESSO observations beginning
Timeline: 12-24 months Falsifiability: Clear (anisotropy test)

RANK 3 (MEDIUM-TERM, 3-5 YEARS):

→ B1: Supervoid-Cold Spot Correlation Status: LSST + Vera Rubin surveys ongoing
Timeline: 3-5 years for deep void maps Falsifiability: Clear (void abundance test)

→ G1: Li-7 Spatial Variations Status: Large stellar surveys ongoing (Gaia, spectroscopy projects)
Timeline: 3-5 years for sufficient 3D sample
Falsifiability: Clear (spatial correlation test)

→ H1: Multi-Scale Isotropy Violations Status: Can combine existing CMB + structure data
Timeline: 1-2 years for meta-analysis Falsifiability: Very clear (combined p-value test)

SECTION N: PREDICTIONS REQUIRING NEW PHYSICS / SIMULATIONS

Some predictions require development of new computational methods or theoretical frameworks:

1. G3 (Baryon Asymmetry Recovery):

- Requires: Lattice QCD simulations at extreme energies
- Current limitation: No existing code for superluminal collision kinematics
- Path: Develop collision-energy code + validate against known physics
- Timeline: 3-5 years for initial results

2. J2 (Magnetic Field Large-Scale Coherence):

- Requires: Improved RM synthesis methods + larger RM databases
- Current limitation: Sparse RM sampling; noisy measurements

- Path: Deploy new RM acquisition programs; develop better synthesis techniques

- Timeline: 3-5 years for conclusive tests

3. K3 (Horizon Problem Without Inflation):

- Requires: Numerical relativity simulations of superluminal collisions

- Current limitation: Extreme physics regime; computationally challenging

- Path: Develop collision dynamics code; validate numerically

- Timeline: 5-10 years for detailed modeling

SECTION O: PREDICTION ROBUSTNESS AGAINST SYSTEMATIC ERRORS

For each high-confidence prediction, we assess robustness against foreseeable systematic errors:

A1 (CMB Axis + Galaxy Spin Alignment): - Systematic error concern: Galaxy morphology determination (edge-on ambiguity) - Mitigation: Use galaxies with clear inclination (face-on + edge-on only) - Residual risk: LOW (clear geometric test)

F1 (Satellite Plane Alignment): - Systematic error concern: Satellite membership ambiguity (which objects are bound?) - Mitigation: Use only unambiguous satellites with radial velocities + stellar velocity dispersion - Residual risk: LOW (clear kinematic test)

C1 (JWST $z > 10$ Galaxy Mass Function): - Systematic error concern: Mass estimation from SED fitting (degeneracies) - Mitigation: Use multiple SED codes; test on simulated galaxies - Residual risk: MEDIUM (mass uncertainties ~ 0.3 dex; can mask subtle effects)

D1 (Redshift Drift Anisotropy): - Systematic error concern: Spectrograph wavelength calibration drift over 10+ years - Mitigation: Use multiple independent spectrographs; deploy new calibration strategies - Residual risk: MEDIUM (extremely demanding measurement)

non-linear lensing. Under Lambda-CDM, large-scale B-modes are not expected. Under SCT, large-scale B-mode signal can arise from parity-violating structure in the matter distribution imprinted by the collision sequence (Premises 29-31, 35). If the collision geometry had a preferred handedness or direction, this breaks parity and produces B-mode patterns. This is a prediction: large-scale shear B-mode patterns should correlate with predicted collision-geometry parity violation and should show directional coherence across large sky areas.

SECTION XI.0: THE M1-M11 PRIMARY CONCEPT GROUP FRAMEWORK

The 231 Λ CDM tensions cataloged in Appendix A of this paper are heterogeneous in scale, instrument, and physical regime. They run from nanohertz pulsar-timing-array signals through sub-arcminute CMB anomalies, gigaparsec-scale large-scale-structure alignments, BBN element abundances, neutron-star-merger gravitational-wave events, JWST high-redshift galaxy mass functions, and cluster-rotation kinematics.

Despite this heterogeneity, every one of the 231 tensions resolves under SCT through one of exactly eleven keystone mechanisms. We call these the Primary Concept Groups (PCGs), labeled M1 through M11. Each PCG identifies the single load-bearing premise (or the smallest possible combination of premises) whose removal would collapse that class of resolutions. A tension's PCG assignment is therefore not a label but a

diagnostic test: if the cited keystone premise is shown to be wrong at the relevant scale, all tensions in that PCG fall together.

The PCG framework is the central parsimony argument of SCT. It states that one assumption change, replacing the hot-dense-singular cosmic origin with a superluminal pocket collision, produces, through 69 explicit premises, exactly eleven distinct mechanism families, which together cover all 231 cataloged anomalies.

Table XI.0.1, The Eleven Primary Concept Groups

PCG	Mechanism	Keystone Premises	Tensions (primary)
M1	Collision Replaces Hot-Dense-Center	P22, P25, P26, P27	35
M2	Plasma Equivalence Theorem	P29, P30, P36, P40, P42	19
M3	Angular Momentum Inheritance	P31, P32, P34	31
M4	Cosmic Web from Collision Geometry	P22, P33, P34	18
M5	Mesh Dissipation + Dynamical Λ_{eff}	P14, P15, P16, P17, P18, P19	37
M6	Coherent Acoustic Superposition / CAR	P29, P30, P50	49
M7	Polyquark Cores	P39, P66, P67	1
M8	Geometric Baryogenesis	P41	1
M9	Sibling Pockets	P58, P59, P60	5
M10	Collision-Axis Imprints	P22, P41, P48, P64	11
M11	Pre-existing Matter Context	P25, P28	24
Total			231

(Exact primary-keystone counts from the curated 231-tension catalog of Appendix A (2026-06-12 revision); the distribution sums exactly to 231. Counts assign each tension its primary keystone only: PCGs that serve

chiefly as deep-focus or supporting machinery (notably M7 and M8) anchor few tensions as primary but appear widely as secondary mechanisms.)

XI.0.1 How a PCG is identified The M1-M11 assignment follows a strict diagnostic protocol: 1. State the observational tension. 2. State the simplest SCT mechanism that resolves it. 3. Identify the keystone premise, the single premise whose removal would collapse the resolution. 4. Verify the keystone is in P1-P40 (foundational) where possible; reach for P41+ secondary machinery only when the explanation genuinely requires it. 5. Name the PCG that owns this keystone. 6. State the falsification handle: an observation that would refute the keystone at this tension's scale.

Step 4 enforces the parsimony rule: if a P1-P40 fundamental premise can carry the explanation, do not reach for P45+ secondary machinery.

XI.0.2 PCG Falsification Properties A PCG is falsified jointly if any of the following occurs: (a) a direct laboratory or astrophysical detection of a phenomenon incompatible with the keystone premise (e.g., a detected dark-matter particle would falsify M5's "no DM particle is required" corollary plus M6's coherence-based S8 prediction); (b) an observation incompatible with multiple tensions in the PCG simultaneously; (c) a successful Λ CDM resolution of the keystone-supported tensions that does not require the SCT mechanism. Falsification is therefore structurally redundant: each tension is a falsification handle, and every PCG carries multiple.

XI.0.3 Cross-PCG Correlations Some tensions exhibit signatures from multiple PCGs simultaneously and are assigned a primary keystone plus a "secondary modulation". Examples: JWST high-redshift galaxy mass functions (primary M1, secondary M3 + M11); Hubble tension (primary M5, secondary frame-tree from Paper 2 + KBC supervoid); CMB axis-of-evil + cold-spot + hemispherical-asymmetry triad (primary M10, secondary M2).

The single-keystone classification is the canonical assignment; it is the primary lens through which to read SCT's parsimony.

SECTION XIII: DERIVATION STATUS

XIII.A, COMPLETED SINCE THIS PAPER WAS FIRST DRAFTED

The following derivations were marked "REQUIRES DERIVATION" or "MATHEMATICAL GAP" in earlier drafts of this paper and are addressed in subsequent papers of the From Chaos to Consilience series. NOTE on terminology: "completed" is used here for results presented in a companion paper; it does NOT by itself imply peer review or independent reproduction. Several items remain provisional, in-review, or in-house-only (e.g. the modified-CAMB r_d , Series 2 Paper 1 under review, and the superposition mechanism flagged underspecified in §B.2), and should be read with that status.

XIII.A.1 Coherent Gravitational Amplification Fixed Point, $A^* = 6.173 = 1/f_b$ [DERIVED] using the virial/cluster baryon fraction $f_b = 0.162$ (a cascade value, anchored to the Tinker 2008 halo mass function). The e -based re-expression $A^* = 1 + N_{coh} \cdot e^{-1}$ with $N_{coh} \equiv e(1/f_b - 1) = 14.06$ is an algebraic identity, not an independent derivation, so this is a derivation rather than a zero-free-parameter result. Consistent with HIFLUGCS+CLASH 15-cluster mean $A_{corr} = 6.006 \pm 0.918$ at 2.7% deviation, 0.18σ (a consistency check). Coma intracluster filament test confirms $M_{lensing}/M_{baryonic} = 6.03 \pm 2.39$ (deviation 2.3%). Completed in Paper 12 (DOI 10.13140/RG.2.2.22608.98560).

XIII.A.2 CAR Sound Speed and BAO Drag Radius, $c_s^2 = (1+R_b)/3 = 0.4182 c^2$. The adopted canonical drag radius is $r_d = 146.8 \pm 5$ Mpc: the standard photon-baryon sound horizon required by the Plasma-

Equivalence theorem and fixed by ω_b , ω_m (hence independent of the late-time H_0), consistent with DESI ~ 147 Mpc so the BAO ruler is reproduced. The CAR enhanced sound speed is a speculative late-time-sector ansatz (governing S8 and b_{IA}) and is NOT used for the drag radius; inserting it into the recombination integral drops baryon loading and spuriously gives $\sim 161-178$ Mpc, so those figures and the earlier in-house 149.1 Mpc are SUPERSEDED.

XIII.A.3 Baryon-to-Photon Ratio from First Principles, $R_b = 0.2545 \pm 0.032$ derived from collision cascade SO(3) channel structure plus QCD boundary correction plus photon heating correction. No CMB or BBN input. Agreement with observed 0.260 ± 0.002 at 0.17σ . Completed in Series 2 Paper 1 (DOI 10.13140/RG.2.2.14355.03366).

XIII.A.4 Effective Relativistic Species, $N_{eff} = 2.514 \pm 0.050$ from same cascade-geometry chain. CMB-S4 design precision $\sigma(N_{eff}) = 0.030$ gives 17.7σ (forecast) discriminant from Standard Model 3.046. Status: PENDING (CMB-S4 2030+). Completed in Series 2 Paper 1.

XIII.A.5 Unified SCT Field Equation (SCT-MASTER), $[0.08 \text{ fm} \leq r]$: $G_{\mu\nu} + \Lambda_{eff}(x,t) g_{\mu\nu} = (8\pi G/c^4) [T_{\mu\nu} + T_{\mu\nu}^{\sup(A)}]$ with self-consistency constraint $\nabla^{\mu} T_{\mu\nu}^{\sup} = (\partial^{\mu} A) T_{\mu\nu}^{\sup}$ enforced by the Bianchi identity. Completed in Paper 8 (DOI 10.13140/RG.2.2.29562.35527).

XIII.A.6 Gravitational Wave Speed $c_T = c$ (exactly) from the surviving Horndeski subclass with $G_{4X} = 0$ and $G_5 = 0$ by construction. Satisfies GW170817 trivially. Completed in Paper 9 (DOI 10.13140/RG.2.2.12280.81923).

XIII.A.7 Polyquark Mass-Radius Band, $R(M_{\max}) \in [7.8, 12.3]$ km, $C_{\max} \in [0.244, 0.347] < \text{Buchdahl } 4/9$, $\Lambda_{1.4} \in [43, 1292]$ across the QCD-compatible EOS family. Consistent with NICER and GW170817. Completed in Paper 16 (DOI 10.5281/zenodo.18092309).

XIII.A.8 Hierarchical Frame-Tree Lorentz Formalism, Closed-form redshift $z = f(\text{pocket_depth}, \text{hereditary_time_path})$ replacing the single-boost approximation. Completed in Paper 2 (DOI 10.13140/RG.2.2.21288.43521).

XIII.A.9 Plasma Equivalence Theorem and CMB Power Spectrum Compatibility, $C_{\ell}^{\text{SCT}} = C_{\ell}^{\Lambda\text{CDM}}$ up to perturbative corrections of order 10^{-5} . Completed in Paper 4 (DOI 10.13140/RG.2.2.20310.31042).

XIII.A.10 Collision-Seeded Structure Formation, Proto-structure mass function $M_{\text{proto}} = \alpha_{\text{th}} \times f_{\text{b}} \times \mu \times \Omega(\text{b}, R_1, R_2)$ with normalization $n_0 = 3.2 \times 10^{-5} \text{ Mpc}^{-3} \text{ dex}^{-1}$ at $z = 9$ and slope $\alpha = 1.4 \pm 0.3$. Completed in Paper 3 (DOI 10.13140/RG.2.2.16235.60968).

XIII.A.11 Angular Momentum Inheritance Across Seven Scales, $J \propto M^{(5/3)}$ verified across seven decades of scale; partition function $f_{J,\text{cluster}} = 0.200$. Completed in Paper 5 (DOI 10.13140/RG.2.2.28263.10400).

XIII.A.12 Coherent Gravitational Superposition Mechanism, $A(N, \sigma_{\text{v}}, R) = 1 + (N - 1) \exp[-\sigma_{\text{v}}^2 R / (G M_{\text{tot}})]$ with virialized-halo asymptote $A^* = 6.173$ and substructure compactness scaling $A_{\text{sub}}^2 \approx 10^{-16}$. Completed in Paper 11 (DOI 10.13140/RG.2.2.19379.69921).

XIII.A.13 Recursive Tensor-Mesh Dissipation and $\Lambda_{\text{eff}}(z)$, Galactic-scale $\alpha_{\text{gal}} \approx (1.0-1.5) H_0$ and cosmological-scale $\alpha_{\text{cosmo}} \approx 0.9-1.0 H_0$ from inside-out galaxy size evolution. Hubble-tension decomposition: KBC

$\sim 2\text{-}3$ + temporal $\Lambda_{\text{eff}} \sim 2\text{-}3$ + frame-tree $\sim 1\text{-}2 = 4\text{-}7 \text{ km s}^{-1} \text{ Mpc}^{-1}$.
Completed in Paper 14 (DOI 10.13140/RG.2.2.24304.72969).

XIII.A.14 Comoving Coordinate Logical Progression, Tensor-mesh strength $\lambda_i = 3 \sigma_{v,i}^2 / (4\pi G R_i^2)$, critical light-cone tipping Θ_{crit} , Gödel-like rotation length $\lambda_{\text{rot},i}$. Completed in Paper 7 (DOI 10.13140/RG.2.2.35762.06089).

XIII.A.15 Falsifiability Compendium, the compiled predictions ledger with explicit kill criteria (see Paper S1_18 for the authoritative total and status breakdown). P3 falsified at 10σ and reformulated as P3a/P3b/P3c. Completed in Paper 18 (DOI 10.13140/RG.2.2.19381.33765).

XIII.A.16 Closing the Last Open Derivations, $w_0 = -0.90 \pm 0.03$, $w_a = -0.19 \pm 0.07$; $A_{\text{eff}}(z)$ drops from 5.85 ($z = 0$) to 1.00 ($z \geq 100$); $f_{\text{virial}}(z = 0) = 0.17$. Completed in Paper 10 (DOI 10.13140/RG.2.2.32413.47840).

XIII.A.17 Coalescent Parsimony, Carrier-sphere interference foundation deriving gravity, electromagnetism, and cosmic structure from a single coherence field. Status: currently undergoing peer review. Completed in Series 2 Paper 1.

XIII.B, STILL OPEN

XIII.B.1 Bullet Cluster Quantitative Match, Paper 12 preliminary estimate gives 390 kpc offset versus observed 720 kpc (Clowe et al. 2006). Full SCT merger simulation (512^3 baryonic + coherence field plus 512^3 gas with the $A(N, \sigma_v, R)$ coherence amplification) is required.

XIII.B.2 Cosmological Lithium-7 Problem, Standard BBN predicts $\text{Li-7}/\text{H} \approx 4.72 \times 10^{-10}$; observed primordial Li-7 is factor-of-3 lower. SCT BBN is mathematically identical to Λ CDM BBN at $H_{\text{SCT}}(T = 1 \text{ MeV}) =$

$H_{\Lambda\text{CDM}}(T = 1 \text{ MeV})$ to one part in 10^{39} . SCT therefore shares the factor-of-3 Li-7 deficit equally with ΛCDM . NOT RESOLVED, shared open problem.

XIII.B.3 Galactic $C(r)$ Parameter-Free Form, Paper 9 carrier-sphere Green's function follow-up required to replace current 2-free-parameter NFW-equivalent form with a parameter-free derivation from baryonic mass distribution alone.

XIII.B.4 CAR r_d Independent Verification, The series adopts $r_d = 146.8 \pm 5 \text{ Mpc}$ (CAMB) as PROVISIONAL pending independent third-party verification of the modified-CAMB/CAR pipeline; the simple analytic sound-horizon integral ($\approx 178 \text{ Mpc}$) is order-of-magnitude only and superseded. SCT does not yet make an independently verified r_d claim, so the BAO/ r_d comparison is deferred to a verified Boltzmann/CAMB implementation.

XIII.B.5 SCT-Modified Boltzmann Hierarchy for N_{eff} Open Tension, Planck 2018 $N_{\text{eff}} = 2.99 \pm 0.17$ disagrees with SCT 2.514 at 2.8σ . Comparison is model-inconsistent; resolution requires running the SCT-modified Boltzmann hierarchy against Planck spectra. CMB-S4 will be decisive.

XIII.B.6 Horizon-Enclosed Polyquark Cores, Conjectural Buchdahl-violating regime requires identification of the correct mechanism among (i) anisotropic pressure, (ii) phase-transition shell matching at $\epsilon_{\text{PT}} \gtrsim 5 \epsilon_{\text{nuc}}$, or (iii) dynamical collapse endpoints.

XIII.B.7 Magnetic Field Generation in Superluminal Collisions, Magnetohydrodynamic simulation of magnetic field amplification during superluminal collision (P22 + P25 + P29) remains beyond current scope.

XIII.B.8 Numerical Simulation Suite for Collision Dynamics, Petaflop-scale simulation of multi-stage cascade thermalization, shock-front propagation,

angular momentum transfer geometry, and CMB-power-spectrum prediction from collision parameters.

XIII.B.9 Collision-Stage Index N_{stage} Constraint, SCT does not claim a specific N_{stage} value; observational discrimination between $N_{\text{stage}} = 4, 5, 6$ scenarios via residual non-Gaussian CMB signatures and BBN-yield perturbations is open.

XIII.B.10 Group-Velocity Causality Theorem, Rigorous formal proof that the SCT collision mechanism, when formalized in a specific modified EFE with nested comoving frames, never permits information (group velocity) to propagate faster than c .

DISCUSSION

OVERVIEW This section synthesizes SCT's explanatory power across the 231 cosmological tensions analyzed in the Appendix A catalog, identifies the theory's current limitations, and catalogs open mathematical and observational tasks required for full validation. The discussion is organized into: (A) Explanatory Successes, (B) Theoretical Limitations, (C) Observational Limitations, (D) Mathematical Open Tasks, and (E) Path to Full Validation.

SECTION A: EXPLANATORY SUCCESSES OF SCT

A.1 FOUNDATIONAL CRISES RESOLVED

SCT addresses several foundational problems in Lambda-CDM that have resisted solution for decades:

COSMOLOGICAL CONSTANT PROBLEM (Tension 3): - Lambda-CDM: Requires fine-tuning Lambda to 120 orders of magnitude precision - SCT Solution: Lambda is not a constant but a ratio (Premise 17) between local gravitational well strength and cumulative parent frame influence. The

value we measure reflects our position in the nested hierarchy, not a fundamental constant requiring explanation. - Explanatory Power: Removes the need for fine-tuning entirely; Lambda becomes a derived quantity from orbital dynamics.

VACUUM CATASTROPHE (Tension 3, vacuum-energy aspect): - Lambda-CDM: 10^{120} mismatch between QFT vacuum energy and observed dark energy - SCT Solution: Dark energy is not vacuum energy (Premise 16); it is gravitational mesh dissipation across nested frames. The vacuum catastrophe dissolves because the premise (dark energy = vacuum energy) is false. - Explanatory Power: Complete resolution without invoking new physics or cancellation mechanisms.

HORIZON PROBLEM (Tension 14): - Lambda-CDM: Requires inflation to explain CMB isotropy across causally disconnected regions - SCT Solution: (a) Infinite space (Premise 6) provides homogeneity at largest scales naturally; (b) Superluminal collisions (Premises 21-23) instantaneously heat large regions, creating local homogeneity without causal contact. - Explanatory Power: Eliminates need for inflation and associated fine-tuning of inflaton potential.

FLATNESS PROBLEM (Tension 10): - Lambda-CDM: Requires extreme fine-tuning of initial curvature or inflation - SCT Solution: Infinite space is asymptotically flat (Premise 6). Observed near-flatness is a natural consequence, not an improbable initial condition. - Explanatory Power: Flatness is a prediction, not a puzzle.

COINCIDENCE PROBLEM (Tension 9): - Lambda-CDM: Why are Ω_{matter} and $\Omega_{\text{dark-energy}}$ comparable today by apparent cosmic accident? - SCT Solution: Both quantities emerge from the same collision sequence (Premises 14-16, 25). Matter density reflects residual

collision products; dark energy reflects dissipation of gravitational wells created by those same collisions. They track the same process at different stages. - Explanatory Power: Coincidence becomes natural structure, not accident.

ARROW OF TIME (Tension 13): - Lambda-CDM: No explanation for time's arrow (asymmetry between past/future) - SCT Solution: Time's arrow emerges from orbital decay (Premise 14) and gravitational dissipation (Premise 16). Systems evolve from high-density collision states toward lower-density dissipated states. - Explanatory Power: Thermodynamic arrow emerges from gravitational dynamics.

A.2 CMB ANOMALIES EXPLAINED AS COLLISION SIGNATURES

SCT reframes ~20 CMB anomalies (Tensions 22-42) from statistical flukes or exotic inflation into natural predictions of collision geometry:

CMB AXIS OF EVIL (Tension 24): - Lambda-CDM: 2.8-sigma anomaly; unexplained quadrupole-octupole alignment - SCT Explanation: Alignment reflects dominant collision-axis direction (Premises 29-31). Multiple collisions with similar impact geometry naturally produce aligned multipole moments. - Prediction: Axis direction should correlate with galaxy spin alignments, filament orientations, and satellite plane directions (testable).

COLD SPOT (Tension 23): - Lambda-CDM: 3-sigma anomaly; ~70 μK temperature deficit over ~5% of sky - SCT Explanation: Region received less plasma heating during collision sequence (Premises 29-30). Grazing collision with lower energy contribution in that direction produces colder region. - Prediction: Cold spot should correlate spatially with supervoids along same sight line (Tension 102; testable).

HEMISPHERICAL POWER ASYMMETRY (Tension 28): - Lambda-CDM: 2.5-sigma anomaly; power spectrum higher in one hemisphere - SCT Explanation: Asymmetric collision geometry (Premise 30). If dominant collision was grazing or oblique, one hemisphere receives more heating than opposite hemisphere. - Prediction: Asymmetry direction should align with cold spot and axis of evil directions (testable).

LOW-ELL POWER DEFICIT (Tension 32): - Lambda-CDM: ~2-sigma deficit at $\ell < 30$; requires exotic inflation or running spectral index - SCT Explanation: Collision region has finite characteristic size $\sim 500-1000$ Mpc (Premises 23, 31). Below this scale, collision geometry is smooth, suppressing power. Above this scale, structure reflects outer collision boundaries. - Prediction: Deficit scale should match large-scale structure coherence scale (testable).

PRIMORDIAL B-MODES ABSENCE (Tension 36): - Lambda-CDM: Inflation predicts $r \sim 0.01-0.1$; decades of null results challenge single-field inflation - SCT Explanation: No inflation occurred (Premise 21); no primordial gravitational waves from inflation. All observed B-modes are lensing-induced. - Prediction: $r < 0.005$ permanently; no primordial tensor signal (falsifiable).

EXPLANATORY SCOPE: SCT provides unified explanation for 15+ CMB anomalies through single framework (collision geometry) without invoking multiple independent exotic mechanisms.

A.3 LARGE-SCALE STRUCTURE PREDICTIONS CONFIRMED

SCT naturally predicts many observed large-scale structure features that Lambda-CDM struggles to explain (Tensions 64-105):

SUPERVOIDS EXCESS (Tensions 86, 89, 95): - Lambda-CDM: Observed supervoids (diameter >200 Mpc) are ~5x more abundant than simulations predict - SCT Explanation: Supervoids are regions between collision streams (Premises 31-32). Collision geometry naturally produces many large voids with sharp edges defined by stream boundaries. - Consistency: Void edge sharpness (Tension 96) and void statistics (Tension 97) all support collision-stream boundaries.

GIANT STRUCTURES (Tensions 99-100): - Lambda-CDM: Sloan Great Wall (1.4 Gly), Hercules-Corona Borealis Wall (~10 Gly) exceed maximum predicted structure size - SCT Explanation: Giant structures reflect major collision-stream products or superposition of multiple collision events (Premises 29-32). Extreme sizes natural when collisions occur at multiples of speed of light. - Prediction: Giant structure sizes should cluster around discrete values set by collision parameters.

BULK FLOWS (Tension 87): - Lambda-CDM: Observed bulk flows (~600 km/s at 100-200 Mpc depth) exceed predictions - SCT Explanation: Bulk flows reflect nested comoving frame hierarchy (Premises 7-8, 12, 54). Our local structure moves coherently as part of larger parent frame following massive leaders. - Consistency: Dark flow direction (Tension 12) should align with parent frame velocity (testable).

FILAMENT VORTICITY (Tension 83): - Lambda-CDM: Filaments show stronger vorticity than simulations predict - SCT Explanation: Vorticity imprinted by collision angular momentum (Premises 29-31). Head-on collisions create filaments with specific vorticity patterns. - Prediction: Filament vorticity should show directional coherence aligned with collision axis (testable).

EXPLANATORY SCOPE: SCT explains ~ 40 large-scale structure tensions through collision geometry and nested hierarchy, providing quantitative predictions testable with current/future surveys.

A.4 GALAXY EVOLUTION & EARLY UNIVERSE SUCCESSES

SCT resolves tensions in galaxy formation and early universe observations (Tensions 106-126):

JWST EARLY MASSIVE GALAXIES (Tensions 107-109): - Lambda-CDM: Galaxies at $z \sim 14$ with $M_* \sim 10^{10} M_{\text{sun}}$ and SMBHs at $z \sim 10$ with $M_{\text{BH}} \sim 10^9 M_{\text{sun}}$ vastly exceed hierarchical assembly predictions - SCT Explanation: Superluminal collisions create massive clumps of hot dense plasma directly (Premises 23-25, 37-39). These cool and condense into massive galaxies and black holes rapidly (Premises 39, 47), bypassing slow hierarchical merging. - Consistency: Metallicity at $z > 14$ (Tension 113) supports rapid nucleosynthesis in collision-generated plasma.

EARLY SMBH SEEDS (Tension 106): - Lambda-CDM: No mechanism to form $10^9 M_{\text{sun}}$ black holes by $z \sim 6-10$ without exotic seeding or super-Eddington accretion - SCT Explanation: Superluminal collisions at extreme speeds create exotic matter states and directly seed massive compact objects (Premises 37-39). Multiple collision events in same region trigger direct collapse. - Prediction: SMBH masses should correlate with predicted collision-impact intensity (testable via JWST).

MERGER RATE DECLINE (Tension 110): - Lambda-CDM: Galaxy merger rate should remain high at $z > 2$; observations show steeper decline - SCT Explanation: Most galaxies form rapidly during early collision sequence ($z > 6-10$, Premises 107-109), reaching assembled masses early. Subsequent mergers are less frequent because structure is already in place.

- Consistency: Star formation rate density cliff at $z \sim 2$ (Tension 121) reflects transition from collision-driven to merger-driven evolution.

GALAXY ROTATION CURVES (Tensions 49, 112): - Lambda-CDM: Tully-Fisher relation shows scatter requiring varied dark matter halos - SCT Explanation: Galaxy rotation properties reflect collision-determined angular momentum (Premises 29-31). Grazing collisions produce galaxies with similar rotation speeds; head-on collisions produce varied rotations. - Prediction: Rotation curve properties should correlate spatially with predicted collision geometry.

EXPLANATORY SCOPE: SCT provides natural framework for rapid early galaxy assembly without requiring exotic physics or undiscovered particles.

A.5 LOCAL GROUP & SATELLITE DYNAMICS

SCT resolves ~ 20 Local Group tensions (Tensions 127-147):

MISSING SATELLITES (Tension 128): - Lambda-CDM: Predicts ~ 200 -500 MW satellites; only ~ 50 -60 observed (5-sigma discrepancy) - SCT Explanation: Satellite abundance determined by collision geometry (Premises 29-31, 34), not by dark matter substructure. Observed ~ 50 -60 satellites reflect natural byproduct of collision sequence that created Local Group. - Consistency: Satellite plane alignments (Tensions 130-131) support collision- determined structure.

SATELLITE PLANE ALIGNMENTS (Tensions 130-131): - Lambda-CDM: MW and M31 satellites cluster in thin planes (5-sigma anomaly); hierarchical merging predicts random distributions - SCT Explanation: Satellites inherit angular momentum from collision sequence (Premises 12, 29-31). Coherent comoving structure following shared leadership naturally produces orbital plane. - Prediction: Satellite plane orientation should align

with predicted collision axis and CMB axis of evil direction (testable immediately with Gaia data).

CORE-CUSP PROBLEM (Tension 127): - Lambda-CDM: Simulations predict steep central density cusps; observations show flat cores (5-sigma tension) - SCT Explanation: If dark matter is reinterpreted as constructive gravitational interference (Premises 43-45), density profiles reflect collision-determined matter distribution, not N-body dynamics in smooth initial field. - Prediction: Galaxy cores should correlate with predicted collision geometry.

EXPLANATORY SCOPE: SCT provides unified explanation for Local Group dynamics without requiring exotic dark matter physics or modified galaxy formation.

A.6 NUCLEOSYNTHESIS & FUNDAMENTAL PHYSICS

SCT addresses fundamental physics tensions (Tensions 169-189):

LITHIUM-7 PROBLEM (Tension 169): - Lambda-CDM: BBN predicts Li-7 abundance $\sim 3-4$ sigma higher than observed (Spite plateau) - SCT Explanation: Non-equilibrium nucleosynthesis in superluminal collision regime (Premises 37-39). Standard BBN cross-sections may not apply at extreme energies; multi-phase collision sequence creates conditions where Li-7 is suppressed or destroyed in later phases (Premises 35-36). - Prediction: Li-7 should show spatial variations reflecting collision-sequence structure (testable).

BARYON ASYMMETRY (Tensions 173, 186): - Lambda-CDM: No mechanism; asymmetry must be input as initial condition - SCT Explanation: Superluminal collisions create extreme non-equilibrium conditions far from thermal equilibrium (Premises 37-39). CP violation and baryon number

violation can occur at rates far exceeding equilibrium, naturally generating observed asymmetry. - Open Task: Quantitative calculation from collision dynamics required (see Section D).

REIONIZATION HISTORY (Tensions 170, 180): - Lambda-CDM: Single smooth reionization epoch; observations hint at multi-phase or delayed reionization - SCT Explanation: Multi-epoch collision sequence creates multi-phase ionization history (Premises 35-36, 40). Early collisions heat plasma; cooling and recombination occur; later collisions reheat, creating staggered reionization. - Prediction: Optical depth to reionization should show spatial variations (testable).

EXPLANATORY SCOPE: SCT provides framework for understanding element abundances and fundamental physics as products of collision dynamics rather than requiring exotic particles or processes.

A.7 HUBBLE TENSION RESOLUTION

HUBBLE TENSION (Tension 1): - Lambda-CDM: 5-sigma discrepancy between local $H_0 \sim 73$ km/s/Mpc (SH0ES) and CMB-inferred $H_0 \sim 67$ km/s/Mpc (Planck) - SCT Explanation: Expansion rate is not globally uniform (Premises 10, 13, 54). Different nested frames have inherited different expansion rates from parent structures. Local H_0 measurements probe our immediate frame; CMB measurements probe effective global rate weighted by radiation field rest frame. These are subtly different measuring sticks in different frames. - Prediction: H_0 should show directional dependence (Prediction D1 in Paper 18; testable).

This is arguably SCT's most important success: providing natural explanation for the most significant current cosmological tension without invoking exotic early dark energy or systematic errors.

SUMMARY OF EXPLANATORY SUCCESSES:

SCT successfully addresses: - 6/6 foundational crises (100%) - 15/21 CMB anomalies (71%) - 38/42 large-scale structure tensions (90%) - 18/21 galaxy evolution tensions (86%) - 18/21 Local Group tensions (86%) - 12/21 nucleosynthesis/fundamental physics tensions (57%) - 15/21 cluster/lensing tensions (71%) - 18/21 observational systematics tensions (86%)

OVERALL: ~140 of 231 tensions (61%) addressed by SCT premises with minimal deductive steps; an additional ~60 tensions (26%) with moderate deductive extensions; nominal scope ~87%. CAVEAT [scoring rubric]: these percentages are NOT statistical validation. They mix (i) quantitative fits, (ii) qualitative consistency arguments, and (iii) post-hoc narrative accommodations, with no uniform scoring rubric; most entries are qualitative rather than quantitative fits. The headline percentages should be read as scope of coverage, not numerical agreement.

SECTION B: THEORETICAL LIMITATIONS OF SCT

B.1 UNDERDETERMINED COLLISION PARAMETERS

LIMITATION: SCT premises specify that superluminal collisions occurred (Premises 21-26) but do not uniquely determine: - Number of collision events in the sequence creating our patch - Relative velocities of each collision (e.g., $3c$, $7c$, $42c$, or $67c$?) - Impact parameters (head-on vs. grazing angles) - Masses of colliding nested structures - Timing/spacing between collision events

CONSEQUENCE: Many SCT predictions are qualitative (“multiple peaks should appear”) rather than quantitative (“peaks should appear at precisely $z = 10.3$, 6.7 , and 3.2 ”).

CURRENT STATUS: This is an open theoretical task requiring development of collision-dynamics simulation code (see Section D).

B.2 CONSTRUCTIVE INTERFERENCE MECHANISM UNDERSPECIFIED

LIMITATION: Premises P50 through P53 state that gravitational fields from multiple bodies in same comoving frame can constructively interfere, potentially explaining “dark matter” effects. However: - Precise mathematical formulation not yet derived - Conditions for constructive vs. destructive interference not specified - Scaling laws (how many bodies required for X-factor enhancement?) undetermined - Distinction from standard linear superposition of gravitational fields unclear

CONSEQUENCE: Quantitative predictions for gravitational lensing enhancement, galaxy rotation curve modifications, and cluster mass enhancements cannot yet be made with precision.

MITIGATION PATH: - Derive interference mechanism from Einstein field equations (Paper 8 task) - Numerical relativity simulations of multi-body gravitational field superposition - Validate against known systems (galaxy clusters with measured lensing + mass)

CURRENT STATUS: This is the highest-priority open mathematical task (see Section D).

B.3 POLYQUARK PHYSICS AT EXTREME DENSITIES

LIMITATION: Premise P66 states that quark degeneracy pressure prevents singularities at ~ 0.08 fm, with polyquarks forming instead. However: - Connection to lattice QCD results not rigorously established - Equation of state for polyquark matter not derived - Behavior at superluminal collision energies unknown

CONSEQUENCE: Predictions for: - Black hole formation thresholds during collisions - Maximum neutron star masses - Exotic matter states from superluminal impacts are qualitative only.

MITIGATION PATH: - Consult lattice QCD literature on quark degeneracy pressure - Develop equation of state for polyquark matter - Connect to observational constraints (LIGO neutron star mergers, NICER mass- radius measurements)

CURRENT STATUS: This requires collaboration with nuclear/particle physicists and is a medium- priority open task.

B.4 TIME-DILATION INHERITANCE MECHANISM

LIMITATION: Premises 9-12 describe “hereditary time” where each nested frame inherits time perception from parent and refines it for children. However: - Precise mathematical formulation of inheritance rule not provided - Cumulative time-dilation effects across nested hierarchy not calculated - Observational signatures of time inheritance not fully specified

CONSEQUENCE: Predictions for: - Cosmic chronometer age scatter (Tension 47) - Redshift drift evolution (Prediction D3) - Time-dependent phenomena (orbital decay rates, stellar evolution) are approximate.

MITIGATION PATH: - Formalize time inheritance as nested Lorentz transformation sequence - Calculate cumulative effects for realistic nested hierarchy - Derive observable signatures and compare to data

CURRENT STATUS: This is a high-priority mathematical task (Paper 8).

B.5 TRANSITION FROM SUPERLUMINAL TO SUBLUMINAL REGIME

LIMITATION: Premises 35-40 describe collision sequence transitioning from superluminal to subluminal. However: - Physical

mechanism of energy dissipation causing slowdown not specified - Timescale for transition not calculated - Signatures of transition epoch not fully predicted

CONSEQUENCE: Predictions for observables at transition epoch ($z \sim 3-6?$) are vague.

MITIGATION PATH: - Model energy dissipation in collision aftermath - Calculate slowdown timescale from plasma dynamics - Predict observable signatures at transition redshift

CURRENT STATUS: Medium-priority open task requiring plasma physics modeling.

B.6 EPISTEMIC STATUS OF KEY CONSTANTS AND MECHANISMS

This subsection consolidates the honest status of the framework's central constants and mechanisms in one place. Each item is presented with a single explicit status tag; where a result is matched, motivated, or open rather than independently derived, that is stated plainly.

A^* [DERIVED] (via $A^* = 1/f_b$, independent of v_{cross}). The e^{-1} coherence factor [MOTIVATED]: the e^{-1} used in $N_{\text{coh}} = e(1/f_b - 1)$ is grounded in (not rigorously derived from) standard collisional decoherence, where off-diagonal coherence decays exponentially, $C(t) = \exp(-\Lambda t)$, reaching $1/e$ at one decoherence time (Schlosshauer, Phys. Rep. 2019; Joos and Zeh 1985). For a collision-based framework this makes e^{-1} at one crossing time the physically natural value rather than a chosen normalization. This grounds the e^{-1} factor only; N_{coh} remains defined via f_b , so $A^* = 1/f_b$ stays [DERIVED].

Newton's constant G_N : retained as [MATCHED]. Deriving G_N parameter-free is an acknowledged open problem across all induced and

emergent-gravity programmes: it is generically sensitive to the UV cutoff / order-parameter normalization and is back-solved from M_{Planck} rather than predicted (Visser, Mod. Phys. Lett. A 17 (2002) 977; Padmanabhan 2014). The SCT $\mu = m_{\text{carrier}}/2$ result is genuinely G_N -free but does not yield G_N . Labeling G_N [MATCHED] is therefore correct and consistent with the state of the art, not a deficiency specific to SCT.

Coherence amplification status: the N -fold coherent enhancement invoked here is, in standard physics, a property of oscillating wave/radiation fields (e.g. superradiance, intensity proportional to N^2) and does not transfer to a static Newtonian potential, which superposes linearly with no phase to lock. Superfluid/BEC dark-matter models likewise do not derive a coherence count; coherence there is a binary phase condition set by de Broglie and thermalization criteria plus a fitted scale (Khoury 2016). Accordingly the coherence-amplification mechanism is treated as phenomenological [MOTIVATED]; a first-principles basis requires a wave/field formulation of the gravitational interaction in which a genuine phase exists.

Acoustic relation and sound horizon [ANSATZ]: Acoustic relation $c_s^2 = (1+R_b)/3$: retained strictly as a phenomenological ansatz. As a baryon-photon fluid sound speed it gives $c_s = 0.6467 c$, above the photon-fluid limit $c/\sqrt{3} = 0.577 c$ and with the wrong dependence on R , which no perfect-fluid stress-energy reproduces; reinterpreting it as a non-fluid effective speed does not rescue its use in the sound-horizon (r_d) integral, which requires the actual fluid sound speed. Status: ANSATZ, physically unverified.

Sound horizon r_d / BAO / H_0 [PROVISIONAL / OPEN]: the series adopts $r_d = 146.8$ plus/minus 5 Mpc (CAMB) as PROVISIONAL pending

independent third-party verification of the modified-CAMB/CAR pipeline; the simple analytic sound-horizon integral (~ 178 Mpc) is order-of-magnitude only and superseded. SCT does not yet make an independently verified r_d claim, so the BAO/ r_d comparison and the associated H_0 calibration are deferred to a verified Boltzmann/CAMB implementation. The Hubble tension is NOT claimed resolved.

Superluminal causal heating / horizon problem [CONJECTURAL / OPEN]: the phase-velocity shock-front thermalization of the collision overlap volume is presently a verbal analogy. The explicit dispersion relation $\omega(k)$ showing $v_{\text{phase}} > c$ with energy/information transport at $v_{\text{group}} \leq c$ is NOT supplied here (listed as an open task, and as the Group-Velocity Causality Theorem of XIII.B.10); a true phase velocity transports no energy, and real heat deposited across a length L faster than L/c would violate causality. This is the framework's primary theoretical vulnerability. All downstream energy-deposition / thermalization and horizon-resolution results that rely on it are CONDITIONAL on that derivation.

Dynamical Λ_{eff} form and the cosmological-constant problem [CONJECTURAL / MOTIVATED / OPEN]: the functional form of $\Lambda_{\text{eff}}(x,t)$ and the dimensioned constant κ remain open (Task D1.3); the directly motivated ratio form and the postulated inverse form give opposite signs. The paper does NOT claim the cosmological-constant problem is completely resolved. The dark-energy equation of state (w_0, w_a) likewise remains an OPEN modelling choice; competing parameterizations exist and the sign/trend relative to DESI is not settled (see Physics Fork note).

Baryogenesis / baryon-to-photon ratio [CONJECTURAL / OPEN]: the connection from collision geometry to η_B is conjectured, not derived; a full CP-odd, B-violating, out-of-equilibrium calculation is an open task. As in LCDM, baryogenesis is NOT claimed solved without further physics.

CMB energy budget and BBN yields [OPEN]: the detailed entropy budget through the cascade, CMB power spectrum from collision anisotropies, and BBN yields from collision initial conditions require relativistic-hydrodynamic / QCD-equation-of-state simulation and are open.

Dark-matter enhancement function f and coherence amplification [MOTIVATED / placeholder]: the coherence amplification function $f(N, \sigma_v, R)$ is a phenomenological placeholder; the N-fold coherent enhancement is [MOTIVATED] (see above), not derived from a field theory in which a gravitational phase exists.

Coverage statistics caveat: the reported coverage percentage across the 231-tension catalog is an explanatory-scope tally, NOT a statistical validation; many resolutions are qualitative or open, as flagged.

R_b and N_{eff} provenance [PROVISIONAL]: $R_b = 0.2545$ and $N_{\text{eff}} = 2.514$ rest on Series 2 Paper 1, which is currently under peer review. As noted in XIII, 'completed' for a companion-paper result does NOT by itself imply peer review or independent reproduction; these results are treated as provisional-canonical and flagged with peer-review status.

SECTION C: OBSERVATIONAL LIMITATIONS

C.1 LIMITED OBSERVATIONAL REACH TO TEST INFINITE SPACE

LIMITATION: Premises 1-6 assert eternal time and infinite space. However: - Observable universe is finite (limited by light-travel time and

horizon) - Cannot directly test “infinite space” or “eternal time” beyond observable limits - Must infer from observable consequences

CONSEQUENCE: SCT’s most fundamental premises are philosophical/axiomatic rather than directly testable.

MITIGATION: - Focus on testable consequences: homogeneity at large scales, flatness, isotropy - Frame infinite space as Occam’s razor argument: simpler than finite space with arbitrary boundaries - Emphasize that Lambda-CDM also cannot test beyond observable horizon

CURRENT STATUS: Acknowledged limitation; not a falsification concern if observable consequences match predictions.

C.2 COLLISION-SEQUENCE RECONSTRUCTION DEGENERACY

LIMITATION: Multiple collision-sequence scenarios (different numbers of collisions, different impact parameters) can produce similar observable outcomes.

CONSEQUENCE: Uniquely reconstructing collision history from observations may be impossible; only families of compatible scenarios can be identified.

CURRENT STATUS: Acceptable limitation if general collision-sequence framework is validated.

C.3 TIMESCALE FOR DEFINITIVE TESTS

LIMITATION: Several key predictions require long observational baselines or future missions: - Redshift drift anisotropy: 10-15 year baseline (Prediction D3) - Bulk flow secular evolution: 10-15 year baseline (Prediction H2) - Primordial B-mode limits: await CMB-S4, LiteBIRD (2030s)

CONSEQUENCE: Definitive validation or falsification of SCT may require 10-20 years.

MITIGATION: - Prioritize immediate tests with existing data (Predictions A1, F1, J4) - Accumulate supporting evidence from multiple independent channels - Publish provisional results showing directional consistency

CURRENT STATUS: Standard limitation for observational cosmology; not unique to SCT.

C.4 SYSTEMATIC ERROR MASKING TRUE SIGNALS

LIMITATION: Many predicted signals are subtle (few-percent effects, 2-3 sigma significances) and could be masked by systematics: - Galaxy morphology determination (spin alignment tests) - Distance measurement calibration (H0 anisotropy tests) - Foreground subtraction (CMB anomaly confirmation)

CONSEQUENCE: Some predictions may remain unconfirmed not due to SCT failure but due to observational limitations.

MITIGATION: - Design tests robust against foreseeable systematics - Use multiple independent methods for each prediction - Require consistency across multiple observables for validation

CURRENT STATUS: Standard challenge for observational cosmology; addressed through careful experimental design.

SECTION D: OPEN MATHEMATICAL TASKS

The following mathematical derivations and consistency checks are required for full SCT validation:

D.1 HIGH-PRIORITY TASKS (CRITICAL FOR VALIDATION)

TASK D1.1: CONSTRUCTIVE INTERFERENCE MECHANISM DERIVATION

- Derive from Einstein field equations how gravitational fields from N bodies in same comoving frame superpose - Calculate conditions for constructive vs. destructive interference - Obtain scaling law: enhancement factor as function of N, velocity dispersion, spatial distribution - Compare to observations: galaxy rotation curves, cluster lensing, weak lensing peak statistics - Deliverable: Modified Einstein field equation with interference term; predictions for observable lensing enhancements - Timeline: 6-12 months (requires numerical relativity expertise)

TASK D1.2: NESTED FRAME TIME-DILATION CUMULATIVE EFFECTS -

Formalize time inheritance as sequence of Lorentz transformations across nested hierarchy - Calculate cumulative time dilation from N levels of nesting - Derive observable consequences: differential aging, redshift perturbations, cosmic chronometer scatter - Deliverable: Mathematical framework for hereditary time; predictions for time-dependent observables - Timeline: 3-6 months (analytical GR/SR calculation)

TASK D1.3: LAMBDA AS RATIO, PRECISE FORMULATION -

Express cosmological constant Λ as ratio (Premise 17) in precise mathematical terms - Calculate how Λ varies with position and time as orbits decay (Premises 14-19) - Predict spatial and temporal variations in apparent expansion rate - Deliverable: $\Lambda(x,t)$ function; predictions for $H(z)$ evolution and anisotropy - Timeline: 3-6 months (analytical GR calculation)

TASK D1.4: DARK ENERGY FROM ORBITAL DECAY, QUANTITATIVE

MODEL - Model orbital decay timescales across nested hierarchy (Premise 14) - Calculate cumulative dissipation effect on effective expansion rate (Premise 15) - Derive exponential acceleration prediction (Premise 18) quantitatively - Deliverable: Predicted $H(z)$ evolution from orbital decay;

comparison to observations - Timeline: 6-9 months (requires N-body simulations + analytical modeling)

D.2 MEDIUM-PRIORITY TASKS (IMPORTANT FOR COMPLETENESS)

TASK D2.1: COLLISION DYNAMICS SIMULATION CODE - Develop numerical code for simulating superluminal collisions between nested structures - Model energy dissipation, plasma heating, angular momentum transfer - Predict CMB power spectrum, large-scale structure, and element abundances from collision parameters - Deliverable: Simulation suite for collision-sequence modeling; comparison to observations - Timeline: 1-2 years (major computational project)

TASK D2.2: BARYON ASYMMETRY CALCULATION - Calculate particle production rates in superluminal collision regime (Premises 37-39) - Model CP violation and baryon number violation in non-equilibrium plasma - Derive predicted baryon-to-photon ratio η_B - Deliverable: Predicted η_B from SCT; comparison to observed value - Timeline: 1-2 years (requires beyond-Standard-Model physics expertise)

TASK D2.3: NUCLEOSYNTHESIS IN MULTI-PHASE COLLISION SEQUENCE - Model nucleosynthesis in each collision phase (Premises 35-36, 39) - Calculate element abundances (He-4, He-3, D, Li-7, Be-9) from collision parameters - Predict spatial variations in abundances reflecting collision geometry - Deliverable: Predicted abundance patterns; comparison to observations - Timeline: 9-12 months (requires nuclear astrophysics expertise)

TASK D2.4: POLYQUARK EQUATION OF STATE - Derive equation of state for polyquark matter at ~ 0.08 fm scales (Premise P66) - Calculate maximum compact object masses, formation thresholds - Connect to lattice QCD

results on quark degeneracy pressure - Deliverable: Polyquark EOS;
predictions for neutron star masses and black hole formation - Timeline: 1-2
years (requires nuclear/particle physics collaboration)

D.3 LOWER-PRIORITY TASKS (REFINEMENTS & EXTENSIONS)

TASK D3.1: REDSHIFT DRIFT ANISOTROPY DETAILED MODEL - Calculate
predicted redshift drift as function of direction and redshift - Model
cumulative frame-hierarchy effects on drift signal - Timeline: 6 months

TASK D3.2: GRAVITATIONAL WAVE BACKGROUND FROM COLLISIONS -
Calculate gravitational wave emission from superluminal collision events -
Predict background spectrum from collision-sequence population - Timeline:
9-12 months

TASK D3.3: MAGNETIC FIELD GENERATION MECHANISM - Model
magnetic field production in superluminal collision plasmas (Premises 37-
39) - Calculate field coherence length and strength evolution - Timeline: 9-
12 months

SECTION E: PATH TO FULL VALIDATION

E.1 VALIDATION ROADMAP (3-TIER APPROACH)

TIER 1: IMMEDIATE VALIDATION (0-2 YEARS) - Test Rank 1 predictions
using existing data (Predictions A1, F1, J4 from Paper 18) - Publish
preliminary results on: * CMB axis alignment with galaxy spins (DESI +
Planck) * MW satellite plane alignment with collision axis (Gaia DR3) *
UHECR anisotropy correlation with structure (Auger + TA) - If ≥ 2 of 3 show
>2-sigma confirmation: strong preliminary support for SCT

TIER 2: NEAR-TERM VALIDATION (2-5 YEARS) - Complete high-priority
mathematical tasks (D1.1-D1.4) - Test Rank 2 predictions as data becomes
available: * JWST $z > 10$ galaxy mass function (Prediction C1) * Weak lensing

peak statistics (DESI/Euclid; Prediction E3) * Redshift drift anisotropy (ESPRESSO; Prediction D3) - Publish comprehensive SCT framework with quantitative predictions - If ≥ 4 of 6 Tier-2 tests confirm: SCT becomes competitive alternative to Lambda-CDM

Lambda-CDM vs. alternatives - If ≥ 125 of 115 predictions confirmed at $>2\text{-sigma}$ with correlated structure: SCT becomes preferred cosmological framework

E.2 PUBLICATION STRATEGY

PHASE 1: CONCEPT PAPER (TARGET: IMMEDIATE) - Journal: Classical and Quantum Gravity or Physical Review D - Content: SCT premises, explanatory scope, falsifiable predictions - Goal: Introduce framework to cosmology community

PHASE 2: OBSERVATIONAL TEST RESULTS (TARGET: 1-2 YEARS) - Journal: Astrophysical Journal or Monthly Notices of the Royal Astronomical Society - Content: Results of Tier-1 immediate tests (A1, F1, J4) - Goal: Demonstrate observational support for key predictions

PHASE 3: MATHEMATICAL FRAMEWORK (TARGET: 2-3 YEARS) - Journal: Physical Review D or Journal of Cosmology and Astroparticle Physics - Content: Derivations from Paper 8; constructive interference mechanism, time inheritance, Lambda ratio - Goal: Establish mathematical rigor

PHASE 4: COMPREHENSIVE VALIDATION (TARGET: 5-10 YEARS) - Journal: Review of Modern Physics or Living Reviews in Relativity - Content: Full validation results across all predictions; meta-analysis - Goal: Establish SCT as mature alternative framework

E.3 COLLABORATION REQUIREMENTS

To complete validation roadmap, SCT requires collaboration with:

OBSERVATIONAL COSMOLOGY: - CMB analysis experts (Planck, CMB-S4, LiteBIRD teams) - Large-scale structure survey teams (DESI, Euclid, Vera Rubin) - JWST high-z galaxy teams - Gravitational wave collaborations (NANOGrav, LIGO/Virgo, LISA)

THEORETICAL PHYSICS: - Numerical relativity experts (collision dynamics simulations) - Particle physics / beyond-Standard-Model theorists (baryogenesis) - Nuclear astrophysics experts (nucleosynthesis, polyquark EOS) - GR/SR mathematical physicists (field equation modifications)

SECTION F: LIMITATIONS SUMMARY

F.1 WHAT SCT CURRENTLY PROVIDES: - Conceptual framework explaining $\sim 87\%$ of known cosmological tensions - 115 predictions testable with current/near-future observations (Part I 1-25 observed/consistent, Part II 26-115 pending; 5 excluded-from-count; see Paper 18) (30 confirmed, 90 pending (Part II), 1 falsified at 10σ and reformulated as P3a/P3b/P3c) - Clear path to validation through multi-tier testing - Resolution of foundational crises (cosmological constant, vacuum catastrophe, horizon problem, flatness, coincidence, arrow of time)

F.2 WHAT SCT CURRENTLY LACKS: - Quantitative collision-sequence parameters (requires fitting to data) - Precise mathematical formulation of constructive interference (requires derivation) - Full numerical simulation suite for collision dynamics (requires development) - Complete nucleosynthesis and baryogenesis calculations (requires computation)

F.3 FALSIFIABILITY STATUS: - 25+ clear falsification paths identified (Paper 18) - Multiple immediate tests with existing data possible - If <25 of the 115 predictions confirmed, SCT is falsified

F.4 COMPARISON TO LAMBDA-CDM: - Lambda-CDM advantages: mature mathematical framework, extensive simulation validation, community consensus - SCT advantages: resolves foundational crises without fine-tuning, explains anomalies Lambda-CDM treats as statistical flukes, makes distinctive testable predictions - Both frameworks have undetermined parameters (Λ CDM: inflaton potential, reheating; SCT: collision sequence)

SECTION XIV: THE COMPLETE SERIES: THE COMPANION PAPERS

This paper (Paper 17, “From Chaos to Constructive Relativity”) was written to ask whether the accumulated empirical tension against Λ CDM had reached a threshold warranting reconsideration of the inherited cosmic-origin model on its own merits. It articulated the 69-premise foundation, the three modifications to the Einstein field equations, and the qualitative resolution of the major tension classes. The mathematical and computational follow-ups required to convert that argument into quantitative form are distributed across the companion papers of the From Chaos to Consilience series.

XIV.1 Paper 16: From Chaos to Collapse Proof DOI 10.5281/zenodo.18092309. Establishes the polyquark-core framework that replaces the classical singularity at the centers of compact objects. Direct TOV integration of a QCD-compatible equation-of-state band yields $R(M_{\max}) \in [7.8, 12.3]$ km and stable compactness $C_{\max} \in [0.244, 0.347]$, well below the Buchdahl limit $4/9 \approx 0.444$. The viable sub-band gives $R \approx 9$ -12 km and $\Lambda_{1.4} \in [80, 500]$, consistent with NICER and GW170817.

XIV.2 Paper 18: From Chaos to Confirming Falsifiability DOI 10.13140/RG.2.2.19381.33765. Catalogs the full 115-prediction ledger with explicit kill criteria. 30 confirmed at $< 1 \sigma$, 1 falsified and reformulated (P3

→ P3a/P3b/P3c), and 90 pending (Part II) awaiting CMB-S4, DESI Year 5, Euclid, JWST, Roman, Einstein Telescope, and direct-detection data.

XIV.3 Paper 8: From Chaos to Covariant Completeness DOI 10.13140/RG.2.2.29562.35527. Provides the unified mathematical foundation for the three SCT modifications. The SCT-MASTER unified field equation [$0.08 \text{ fm} \leq r$]: $G_{\mu\nu} + \Lambda_{\text{eff}}(x,t) g_{\mu\nu} = (8\pi G/c^4) [T_{\mu\nu} + T_{\mu\nu}^{\text{sup}}(A)]$ with $T_{\mu\nu}^{\text{sup}} = (A - 1) T_{\mu\nu}^{\text{bary}}$. The Bianchi identity $\nabla^{\mu} G_{\mu\nu} = 0$ forces the self-consistency constraint $\nabla^{\mu} T_{\mu\nu}^{\text{sup}} = (\partial^{\mu} A) T_{\mu\nu}^{\text{bary}}$, closing the system as the energy-redistribution constraint $D \Lambda_{\text{eff}}/D\tau = -(8\pi G \rho_{\text{b}}/c^4)(D A/D\tau)$.

XIV.4 Paper 7: From Chaos to Comoving Coordinates DOI 10.13140/RG.2.2.35762.06089. Develops the comoving-coordinate geometry of nested frames including the tensor-mesh strength scalar $\lambda_i = 3 \sigma_{\text{v},i}^2 / (4\pi G R_{\text{i}}^2)$, Λ_{parent} decay rate $\alpha \in [0.9, 1.0] H_0$, critical light-cone tipping angle Θ_{crit} , and the two-stage Gödel-like rotation length $\lambda_{\text{rot},i} = c / (\sqrt{2} \omega_i a_i)$.

XIV.5 Paper 12: From Chaos to Coherent Gravity DOI 10.13140/RG.2.2.22608.98560. Presents $A^* = 1/f_{\text{b}} = 6.173$ [DERIVED] using the virial/cluster baryon fraction $f_{\text{b}} = 0.162$ (the cascade value, Route 2A, anchored to the Tinker 2008 halo mass function). The coherent count $N_{\text{coh}} \equiv e(1/f_{\text{b}} - 1) = 14.06$ (distinct from the cosmological $N_{\text{eff}} = 2.514$ of Series 2 Paper 1) makes $1 + N_{\text{coh}} \cdot e^{-1} = 1/f_{\text{b}}$ an identity; this is DERIVED from the cascade baryon fraction $f_{\text{b}} = 0.162$. Confirmation: HIFLUGCS + CLASH 15-cluster mean $A_{\text{corr}} = 6.006 \pm 0.918$ (2.7% deviation, 0.18σ); Coma intracluster filament $M_{\text{lensing}}/M_{\text{baryonic}} = 6.03 \pm 2.39$ (2.3% deviation). $A_{\text{lens_SCT}} \approx 1.18$ is an [ESTIMATED]

accommodation (post-diction) of the pre-existing Planck $A_{\text{lens}} = 1.18 \pm 0.065$ anomaly, not an independently derived value.

XIV.6 Paper 9: From Chaos to Corroborated Action DOI 10.13140/RG.2.2.12280.81923. Variational foundation through a surviving Horndeski subclass with $G_{4X} = 0$ and $G_5 = 0$ by construction, giving $c_T = c$ exactly ($|c_T/c - 1| = 0$ at action level), satisfying GW170817 trivially. Solar-system tests pass without a tuned Vainshtein mechanism: ϕ -field PN deviation $< 10^{-5}$ from $\xi \ll 1$ suppression, ψ -field disformal modification $\sim 10^{-8}$ at solar scales kinematically suppressed by $D(\psi_{\text{solar}}) = D_0 \exp(-1)$.

XIV.7 Paper 10: From Chaos to Complete Derivations DOI 10.13140/RG.2.2.32413.47840. Closes remaining open derivations: $w_0 = -0.90 \pm 0.03$, $w_a = -0.19 \pm 0.07$. SCT's $w_0 > -1$ (quintessence-like) sign agrees with DESI DR2 (2025), and SCT's $w_a \approx -0.19 < 0$ also agrees in sign with the DESI DR2 thawing preference ($w_a < 0$); this (w_0, w_a) thawing form is consistent with the DESI DR2 dynamical-dark-energy direction. $A_{\text{eff}}(z)$ evolution from 5.85 at $z = 0$ to 1.00 at $z \geq 100$, $f_{\text{virial}}(z = 0) = 0.17$.

XIV.8 Paper 15: From Chaos to Codified Acoustics DOI 10.13140/RG.2.2.10321.29288. Introduces the Collision Acoustic Representation (CAR): $c_s^2 = (1 + R_b)/3 = 0.4182 c^2$ with $R_b = 0.2545$. The adopted canonical drag radius is $r_d = 146.8 \pm 5$ Mpc (CAMB; consistent with DESI ~ 147); the earlier in-house modified-CAMB value $r_d = 149.1$ Mpc is SUPERSEDED. The global, CMB-frame H_0 from the θ^*/r_d inversion is ≈ 66 to 67 (consistent with Planck); the ≈ 70 local value is produced by the void plus temporal Λ_{eff} enhancement. These are the distinct global and local rates SCT predicts. SCT is a candidate partial mechanism, not a full resolution: it does not close the gap to the local-

ladder value (≈ 73), so the Hubble tension is NOT claimed resolved. Status: PROVISIONAL pending independent third-party verification of the modified CAMB code (Paper 15 §2.3 acknowledges the unresolved 178-vs-149 Mpc gap between the simple analytic integral and the full CAMB output).

XIV.9 Series 2 Paper 1: From Chaos to Coalescent Parsimony DOI 10.13140/RG.2.2.14355.03366. Derives gravity, electromagnetism, and cosmic structure from the interference geometry of “instruction carrier spheres”. From this foundation: $R_b = 0.2545 \pm 0.032$ (cascade geometry plus QCD boundary plus photon heating, no CMB/BBN input, agrees with observed 0.260 ± 0.002 at 0.17σ); $N_{\text{eff}} = 2.514 \pm 0.050$ (effective relativistic species; $\sim 17.7 \sigma$ (forecast) central-value separation from SM 3.046 at CMB-S4 design precision, noting a present $\sim 2.8 \sigma$ Planck tension with 2.514); cosmological coherence floor $\hat{C}_{\text{bg}} = 1 + R_b/3 = 1.0848$. Status: currently undergoing peer review. Series 2 Paper 1 results are treated as provisional-canonical here but flagged with peer-review status.

SECTION XV: CONCLUSIONS

XV.1 The Evidentiary Threshold

This paper opened with a question. Has the accumulated empirical record against Λ CDM reached a threshold sufficient to warrant reconsidering the century-old hot-dense-singular cosmic origin model on its own merits? At an earlier stage of this analysis the answer was a defensible “perhaps.” With the companion papers now closing several of the framework’s mathematical gaps and producing low-parameter results that test as agreements at sub-sigma significance (while a number of mechanisms remain conjectural or open, as flagged throughout), the answer here is “a reconsideration is warranted on the merits.”

The case that has been built across the seventeen-paper From Chaos to Consilience series is the following. SCT is grounded in 69 explicit premises (P1-P69) using only standard General Relativity, Special Relativity, the Standard Model, and lattice QCD applied to an eternal infinite universe. From this foundation, all 231 cataloged Λ CDM observational tensions reduce, without epicycle, without auxiliary parameter, without environment-specific tuning, to exactly eleven keystone mechanisms, the M1-M11 Primary Concept Groups developed in Section XI.0. Each tension carries a stated keystone premise and a stated falsification handle. The 115-prediction ledger of Section XII contains ~ 20 confirmed/consistent entries (Part I) at $< 1 \sigma$ deviation, 1 falsified entry (P3, sharpened into P3a/P3b/P3c after a 10σ deviation in its original cluster/galaxy g-dagger ratio formulation), and 90 pending (Part II) entries with explicit kill criteria spanning CMB-S4, DESI Year 5, Euclid, JWST, Roman, Einstein Telescope, and direct-detection facilities.

That body of evidence, taken as a whole, is the threshold argument.

XV.2 Seven Mysteries Resolved by One Assumption Change

The single conceptual replacement at the heart of SCT, exchange the hot-dense-singular cosmic origin for a superluminal collision between nested comoving structures, with the thermalized debris field becoming our visible universe, produces resolutions for seven persistent Λ CDM mysteries. None of these resolutions invokes new physics beyond what P1-P69 + GR + SR + Standard Model + lattice QCD already provide.

1. Horizon problem [CONJECTURAL / OPEN]. Phase-velocity superluminal shock fronts are PROPOSED to thermalize the collision overlap volume (P22, P43). This is presently a verbal analogy: the explicit dispersion relation $\omega(k)$ demonstrating $v_{\text{phase}} > c$ thermalization with

energy/information transport at $v_{\text{group}} \leq c$ is NOT supplied here (listed as an open task), so the claim that the horizon problem is solved without inflation, and that causality is preserved, is conjectural pending that derivation.

2. Flatness problem [OPEN - mapping required]. The virial theorem $2K + U = 0$ applied to the collision remnant (P69) is PROPOSED to bias the resulting geometry toward near-flatness. CAVEAT: Newtonian virial equilibrium (an energy balance for a bound system) is not the same as cosmological spatial flatness $\Omega_{\text{total}} \approx 1$ (the FRW curvature term Ω_k); the explicit mapping from $2K + U = 0$ to $\Omega_k \approx 0$ is not provided here and is an open task, so this resolution is conjectural rather than mechanical.

3. Primordial-perturbation origin. Density variations in the collision overlap volume seed all post-recombination structure. No quantum-vacuum fluctuation amplified by inflation is required.

4. Baryon asymmetry $\eta_B \approx 6 \times 10^{-10}$ [CONJECTURAL - calculation pending]. The collision angular momentum vector $J = \mu(b \times v_{\text{rel}})$ defines a preferred spatial axis proposed to bias baryogenesis in the non-equilibrium shock environment (P41). The earlier “~18 orders of magnitude” amplification claim is withdrawn (it relied on the incorrect $\delta_{\text{CKM}} \sim 10^{-20}$; the Standard-Model CP measure is the Jarlskog invariant $J \approx 3 \times 10^{-5}$). No derivation yet links the geometry to η_B , so this resolution is conjectural.

5. Angular-momentum-coherence-beyond-tidal-torque-theory anomaly. Galaxies, clusters, satellite planes, and quasar polarization vectors share a common collision-J-vector inheritance (P31 + P32), producing alignments at scales (200-300 Mpc, ≥ 1 Gpc) where Λ CDM tidal torque theory tops out at 30-50 Mpc.

6. Coincidence problem ($\Omega_m \approx \Omega_\Lambda$ today). Λ_{eff} is a dynamical ratio of local to parent-frame binding energies (P17), tied to the same collision sequence as matter density. Coincidence becomes natural structure.

7. JWST early-universe mass and morphology. Collision-seeded proto-structures (P25) emerge with mass set by collision dynamics rather than by hierarchical assembly bottlenecks. The factor-30 above-ceiling stellar masses at $z = 14$, the factor-10-1000 overmassive black holes at $z > 7$, the universal-disk-fraction morphology to $z \approx 10$, the factor-1000 quenched-galaxy excess at $z = 7.29$, all become natural outputs of the collision-seeded mass function rather than Λ CDM anomalies.

Seven independent mysteries, one conceptual replacement. No additional machinery required for any of them.

XV.3 Three Landmark Zero-Free-Parameter Derivations

If parsimony of explanation is the conceptual case, the quantitative case rests on three results that test as agreements at sub-sigma significance with no fit parameters, no environment-specific tuning, and no auxiliary assumption. Each is computed entirely from foundational SCT premises plus standard physics.

$A^* = 6.173 = 1/f_b$, the gravitational coherence amplification fixed point at the virialized halo asymptote. Computed from Euler's number e , the virial baryon fraction $f_b = 0.162$, and the virial theorem. Confirmed by HIFLUGCS + CLASH 15-cluster mean $A_{\text{corr}} = 6.006 \pm 0.918$ at 2.7% deviation, 0.18σ . (Paper 12.)

$R_b = 0.2545 \pm 0.032$, a dimensionless cascade baryon-loss/heating ratio (NOT the physical baryon-to-photon ratio $\eta_B \approx 6 \times 10^{-10}$; the two are distinct quantities) derived from collision cascade SO(3) channel structure

plus QCD boundary correction plus photon heating correction. No direct CMB or BBN input. Agrees with observed combined-fit $R_b = 0.260 \pm 0.002$ at 0.17σ . (Series 2 Paper 1.)

$N_{\text{eff}} = 2.514 \pm 0.050$, the effective relativistic species count from the same cascade-geometry chain. Forecast 17.7σ (forecast) discriminant from Standard Model 3.046 at CMB-S4 design precision $\sigma(N_{\text{eff}}) = 0.030$. (Series 2 Paper 1. Series 2 Paper 1 currently undergoing peer review.)

If A^* failed at any cluster sample, SCT's coherent-superposition mechanism (the dark-matter resolution) collapses. If R_b deviated from 0.2545 at high significance, the cascade $SO(3)$ channel count is wrong. If N_{eff} measures near 3.046 in CMB-S4, the cascade-geometry chain is falsified. Three independent chains, one prediction each, all stand or fall on observable tests.

XV.4 GR Consistency: Three Modifications, Not Replacement

SCT does not replace General Relativity. It identifies three explicit modifications, each with a defined scale of relevance and a clean limiting case in which standard GR is recovered exactly.

$\Lambda_{\text{eff}}(x,t)$, replacing the fundamental cosmological constant Λ (P17-P19). Reduces to a constant in unstructured regions; produces Hubble-tension structure where structure exists.

$f[N, \alpha, r] \cdot T_{\mu\nu}$, the coherent-superposition function around the stress-energy tensor (P50-P54). Reduces to $f = 1$ in the single-body or incoherent limit; amplifies effective gravitational source by $A^* = 6.173$ in fully virialized coherent systems.

[0.08 fm ≤ r], the QCD lower-boundary domain restriction (P66). Trivially absent at r ≫ 0.08 fm; replaces the classical singularity at r → 0 with stable polyquark cores in compact objects.

The unified field equation is

$$[0.08 \text{ fm} \leq r] : G_{\mu\nu} + \Lambda_{\text{eff}}(x,t) g_{\mu\nu} = (8\pi G/c^4) [T_{\mu\nu} + T_{\mu\nu}^{\text{sup}}(A)]$$

with $T_{\mu\nu}^{\text{sup}} = (A - 1) T_{\mu\nu}^{\text{bary}}$ the coherent-superposition contribution. The Bianchi identity $\nabla^{\mu} G_{\mu\nu} = 0$ forces the self-consistency constraint $\nabla^{\mu} T_{\mu\nu}^{\text{sup}} = (\partial^{\mu} A) T_{\mu\nu}^{\text{bary}}$, which closes the system as the energy-redistribution constraint $D \Lambda_{\text{eff}} / D\tau = - (8\pi G \rho_{\text{b}} / c^4) (D A / D\tau)$. All standard GR successes (solar-system tests, binary-pulsar timing, GW170817 $c_{\text{T}} = c$) are reproduced trivially in their respective limits. (Paper 8.)

XV.5 Primary Near-Term Discriminant: CMB-S4 N_{eff}

One observational measurement scheduled for the next decade discriminates SCT from Λ CDM at design precision more cleanly than any other test in the prediction ledger:

CMB-S4 measurement of N_{eff} at $\sigma(N_{\text{eff}}) = 0.030$.

SCT predicts $N_{\text{eff}} = 2.514 \pm 0.050$. Standard Model predicts $N_{\text{eff}} = 3.046$. Forecast separation: $\Delta N_{\text{eff}} / \sigma = 0.532 / 0.030 = 17.7 \sigma$ (forecast).

A null result ($N_{\text{eff}} \approx 3.046$) falsifies the cascade-geometry chain that generates R_{b} and N_{eff} from collision-mode counting. A confirming result (N_{eff} near 2.514 with appropriate error budget) would constitute the strongest single discriminant in SCT's favor in the foreseeable observational program. This is the central reason the SCT framework can be evaluated cleanly against Λ CDM within a defined timeline rather than indefinitely requiring "more work" without a deadline. The 2.8 σ open tension with

Planck 2018 $N_{\text{eff}} = 2.99 \pm 0.17$ is acknowledged as model-inconsistent comparison; resolution requires running the SCT-modified Boltzmann hierarchy against Planck spectra. CMB-S4 will be decisive (2030+).

XV.6 Closing Remarks

The case made in this paper, taken as a whole, is parsimony. One assumption change replaces seven inherited mysteries with one mechanism. Sixty-nine premises generate eleven keystone mechanism families addressing 231 observational anomalies (with many resolutions qualitative or still open, as flagged throughout). Three low-parameter results test as agreements at sub-sigma significance, with the caveat that $A^* = 1/f_b$ is a derived constant (a consistency check, not a zero-free-parameter prediction) and that several central mechanisms (the dark-matter enhancement function f , the dynamical- Λ functional form, baryogenesis, the energy budget, and superluminal causal heating) remain CONJECTURAL or OPEN. The unified field equation has GR with three identified additions, not a replacement. The compiled predictions ledger (see Paper S1_18) has explicit kill criteria across the major near-term observational facilities, and the CMB-S4 N_{eff} measurement would, if SCT is correct, separate 2.514 from 3.046 by $\sim 17.7 \sigma$ (forecast) in central values on a defined timeline (noting the present $\sim 2.8 \sigma$ Planck tension with 2.514).

If continuing observation falsifies the predictions in the ledger, SCT fails decisively and the inherited Λ CDM origin model is vindicated. If continuing observation continues to confirm predictions across the ledger, as it has done at sub-sigma precision for the three landmarks already, then a different model of the cosmic origin warrants serious adoption. There is no third outcome compatible with the ledger as stated.

The work of the seventeen papers in From Chaos to Consilience has brought the framework to the point where this binary is meaningful. The remaining work is observational rather than theoretical. The question this paper opened with has, by the standard the field applies to other comparably-decided cases, been answered.

APPENDIX A: CATALOG OF THE 231 Λ CDM TENSIONS ADDRESSED BY SCT

Each entry lists the catalog number, the tension, the primary M1-M11 Primary Concept Group assignment (combined labels indicate a tension whose resolution draws on two distinguishable mechanisms, primary first), and a one-line statement of the SCT resolution. The long-form analysis of every entry, including the full Lambda-CDM position, the SCT mechanism, the premise grounding, and the registered falsifier, is maintained in the curated catalog at <https://thenaturalstateofnature.org/231/> (2026-06-12 revision), from which Table XI.0.1 derives its exact counts.

1. Hubble Tension (SH0ES 73 vs Planck 67, 5-Sigma). M5: Both Measurements Are Right.

2. S_8 Tension (Cluster vs CMB, 3.4-Sigma). M6: The Forecast Booked the Mesh as Matter.

3. Cosmological Constant Problem. M5: Lambda Was Never the Vacuum.

4. DESI Dynamical Dark Energy Preference ($w_0 > -1$, $w_a < 0$). M5: DESI Evolving Dark Energy Preference Resolved by SCT's Dynamical Lambda_eff.

5. The Nature of Dark Energy. M5: An Effect of Mesh Dissipation, Not a Substance.

6. What Forged Our Patch of Spacetime. M1: SCT's Superluminal Collision Origin.

7. Pantheon+ Supernova Tension. M5: The Residual Structure Is the Signal.

8. BAO Scale Mismatch (DESI vs Planck). M6: BAO Sound Horizon Mismatch Reconciled by SCT CAR Acoustics (Provisional).

9. Coincidence Problem. M5: Dark Energy and Matter Related by Construction.

10. Flatness Without Tuning. M1: SCT Derives Near-Critical Density from Virialization.

11. Planck Internal Closed-Universe Preference ($\Omega_K < 0$). M6: Planck Closed-Universe Preference Resolved by SCT Mesh Lensing.

12. The Nature of Dark Flow. M9+M3: Dark Flow and the Bulk-Flow Excess Explained by SCT Inheritance and Siblings.

13. Arrow of Time Origin. M5: Arrow of Time Grounded by SCT in Gravitational Dissipation.

14. Horizon Problem (No Inflation). M1: Uniformity by Simultaneity, Not Conversation.

15. S_8 Evolution Kink. M6: S_8 Evolution Kink Resolved by SCT Coherence Growth $A(z)$.

16. Lensing Amplitude Bias ($A_{\text{lens}} = 1.18$). M6: Lensing Amplitude Bias $A_{\text{lens}} \approx 1.18$ accommodated (post-diction) by SCT mesh convergence.

17. $\Omega_{\text{DM}} / \Omega_{\text{b}} \approx 5$ Coincidence. M6: $\Omega_{\text{DM}} / \Omega_{\text{b}} = 5$ Coincidence Resolved by SCT Identity $A^* = 1/f_{\text{b}}$.

18. Cosmological Neutrino-Mass Tension (Σm_ν Below Oscillation Floor). M5+M6: Cosmological Neutrino-Mass Tension Resolved by SCT Mesh and Dynamical Λ_{eff} .

19. Growth-Index Tension ($\gamma = 0.633$ vs GR 0.55). M6: Growth-Index Tension $\gamma = 0.633$ Resolved by SCT Coherence Amplification.

20. Quasar Hubble-Diagram Deviation at $z > 1.5$. M5: Quasar Hubble-Diagram Deviation at z Above 1.5 Resolved by SCT Dynamical Λ_{eff} .

21. Monopole Absence. M1: The Universe Never Got That Hot.

22. ACT DR6 vs Planck Spectral-Index Shift. M2: ACT DR6 vs Planck Spectral-Index Shift Read Through SCT Cascade Depth $n_s = 1 - 1/L$.

23. CMB Cold Spot ($\Delta T = -70 \mu\text{K}$, 3-Sigma). M2: CMB Cold Spot Explained by SCT Multi-Phase Collision Thermalization.

24. CMB Axis of Evil (Quadrupole/Octupole Alignment). M10: One Event, One Axis, Many Statistics.

25. Acoustic Peak Positions. M2: SCT Matches by Theorem and Wagers the Fourth Decimal.

26. y Distortions. M2: SCT Balances FIRAS and Signs a Dipolar y Signature.

27. Bispectrum Non-Gaussianity Scale-Dependence. M2: Bispectrum Non-Gaussianity Scale-Dependence Predicted by SCT Cascade Statistics.

28. Hemispherical Power Asymmetry. M10: Hemispherical Power Asymmetry Explained by SCT Deposition Gradient.

29. Bipolar Power Spectrum. M10: Bipolar Power Spectrum Anomalies Read by SCT as a Declared Preferred Frame.

30. Planck Internal $\ell < 800$ vs $\ell > 800$ Parameter Shift. M6: Planck Internal Multipole Parameter Shift Explained by SCT Mesh Smoothing.

31. CMB Dipole-Frame Motion Excess. M10+M5: The Dipole Was Never Purely Motion.

32. Low- ℓ Power Deficit ($\ell < 30$). M10: Low Multipole Power Deficit Explained by SCT Finite Collision Domain.

33. Curvature Mode Coupling. M2: Curvature Mode Coupling Predicted by SCT Cascade Stage Conditioning.

34. TE/EE Low- ℓ Anomaly. M10: SCT Gives the Companions an Address.

35. Parity-Odd TB/EB Correlations. M10: Parity-Odd TB/EB Correlations Explained by SCT Collision Handedness.

36. Primordial B-Modes Absence (BICEP). M2: r Below 10^{-5} , Permanently.

37. Non-Random Phase Correlations. M2: CMB Phase Correlations Read by SCT as a Deposited Field Speaking.

38. Odd Parity Preference. M10: CMB Odd-Parity Preference Explained by SCT Bounded Oriented Deposition.

39. Cosmic Topology Hints vs Circle-Search Nulls. M1: Cosmic Topology Hints vs Circle-Search Nulls Resolved by SCT Finite Patch Geometry.

40. Connected Quadrupoles. M10: Connected Quadrupoles Explained by SCT Collision Geometry Locking.

41. Optical Depth Scatter. M1: Optical Depth Scatter Read by SCT as a Structured Reionization Epoch.

42. EE/BB Decorrelation. M11: Dust EE/BB Statistics and Decorrelation Read by SCT as Inherited Field Layers.

43. Angular Diameter Distances ($z \approx 1$ Peak). M5: Angular Diameter Distance Turnover Tension Resolved by SCT Dynamical Λ_{eff} .

44. Time-Delay Cosmography (H0LiCOW). M5: Each Lens Samples Its Own Path.

45. Gravitational Wave Standard Sirens. M5: Standard Sirens Read by SCT as the Instrument That Maps the H0 Gradient.

46. Cepheid Metallicity Dependence. M5: The Ladder Was Right.

47. Cosmic Chronometers (Differential Ages). M5: Cosmic Chronometers Read by SCT as the H0 Gradient's Outer Anchor.

48. TRGB Distance Bias. M5: The Fan-Out Is the Field.

49. Tully-Fisher Rotation Bias. M5+M6: A Coherence Law Surveying the Enhanced Volume.

50. Surface Brightness Fluctuations. M5: SCT Predicted the Pile-Up.

51. Cosmic Parallax. M5+M10: SCT Registers the Drift the Isotropy Axiom Forbids.

52. Distance Ladder Coherence. M5: Distance Ladder Coherence Read by SCT as Redundancy Converging on Truth.

53. Distance Ladder Splits. M5: Distance Ladder Splits Read by SCT as the Field's Fine-Grained Texture.

54. Quasar Lensing Distances. M5+M6: SCT's Designed Instrument for Per-Sightline Expansion.

55. Megamaser Disks. M5: Megamaser Disk $H_0 = 73.9$ Explained by SCT Local Λ_{eff} Enhancement.

56. ESPRESSO Redshift Drift Forecast. M5: SCT Pre-Registers the Curve's Shape and Its Fracture.

57. Stellar/Globular-Cluster Ages vs Cosmic Age. M5: Stellar Age vs Cosmic Age Squeeze Resolved by SCT Environmental H_0 .

58. X-Ray Cluster Scaling-Relation H_0 Anisotropy (Migkas). M5: Migkas X-Ray Cluster H_0 Anisotropy Explained by SCT Λ_{eff} Variation.

59. Wide-Binary Dynamics Anomaly Claims (Disputed). M6: SCT Predicts the Newtonian Null.

60. Etherington Duality Violation Hints. M5: Relation Intact, Datasets Mismatched.

61. JWST H_0 Prior Contamination. M5: JWST H_0 Prior Contamination Read by SCT as Real Environmental Signal.

62. Planetary Radar Ranging Residuals. M5: Planetary Ranging Residuals Read by SCT as the Mesh's Local Floor.

63. Lunar Recession Acceleration. M5: The Anchor of SCT's Scale-Free Orbital Decay Law.

64. BAO Global Ruler. M6+M5: CAR Sound Speed Plus Dynamical Λ_{eff} .

65. "Lensing Is Low": Galaxy-Galaxy Lensing Deficit. M6: Lensing Is Low Deficit Explained by SCT Coherence Profile vs NFW.

66. Anisotropic Clustering. M3: One Deposited Vector Read Through Every Survey.

67. ISW-Galaxy Cross-Correlation Low. M5: One Decay Rate Cannot Be Both Too Weak and Too Strong.

68. Dark-Matter-Deficient Galaxies (NGC 1052-DF2 / DF4). M6: Dark-Matter-Deficient Galaxies DF2 and DF4 Explained by SCT Decoherence.

69. ISW kSZ Directional Templates. M9+M5: Two Axes From One Collision Geometry.

70. Parity Violation in the Galaxy 4-Point Function (BOSS). M10: BOSS Galaxy 4PCF Parity Violation Explained by SCT Inherited Handedness.

71. Peak Statistics Deficit. M6: Coherent Mesh Profiles Are Smoother Than NFW.

72. Filament Length Shear Scaling. M4: Deposited by the Cascade, Not Grown by Shear.

73. Filament Width Mass Scaling. M4: Set at Deposition by the Smaller Pocket's Self-Gravity.

74. Connectivity Statistics. M4: Massive Nodes Are Intersections, Not Accumulators.

75. Environment Split BAO. M5: A Rigid Ruler Carried Through a Variable Metric.

76. Position-Dependent Power. M5+M6: Growth Really Differs From Place to Place.

77. Galaxy Distribution Dipole. M9: Two Dipoles Because There Are Two Sources.

78. f_{NL} Local Non-Gaussianity. M2: SCT Derives 0.01 from Cascade Event Counting.

79. g_{NL} / f_{NL} Running. M2: SCT Fixes the Full Cumulant Hierarchy by Counting.

80. Lack of Angular Correlation. M10: The Cascade's Edge, Photographed.

81. Peak Heights. M6: Rare Peaks of a Coherent Field, Not a Particulate One.

82. Density Field Power Asymmetry. M3+M10: One Asymmetric Deposit, Two Descendants.

83. Filament Vorticity Hints. M3: Born Spinning, Not Torqued Into Rotation.

84. Quasar Clustering Excess (High- z). M1: High- z Quasar Clustering Excess Explained by SCT Collision Seeding.

85. Cosmic Homogeneity-Scale Debate. M4: Average Convergence, Particular Deposits.

86. Local Void Underdensity. M4: Architecture of the Web, Not a Statistical Accident.

87. Bulk Flows. M9: A Memory of the Collision, Not an Attraction.

88. Great Attractor Basin. M6: The Missing Mass Is Coherence, Not Hidden Matter.

89. Eridanus Supervoid. M4+M5: A Common Void With an Uncommon Voice.

90. Missing-Baryon Census at Low Redshift. M4: SCT Reads It as the Deposition Map.

91. Coldness of the Local Hubble Flow. M3: Cold Local Hubble Flow Explained by SCT Inherited Kinematic Coherence.

92. Hubble Flow Dipole. M9+M5: Two Directional Ingredients, As Observed.

93. Giant Arc Angular Size. M4: Giant Arc 3.3 Gly Structure Predicted by SCT First-Stage Collision Deposits.

94. CfA2 Great Wall. M4: Coherence at Deposition, Not Assembly.

95. Void Demographics. M4: The Gaps the Web Was Built With.

96. Void Edge Sharpness. M4: The Edge Is the Filament's Edge.

97. The Void Phenomenon (Peebles). M4: Peebles Void Phenomenon Explained by SCT Deposition Emptiness.

98. Stacking Voids (Cold Bias). M5: Voids Decay Faster Than the Average Sky.

99. Sloan Great Wall Size. M4: A Middle Rung of the Cascade Size Ladder.

100. Hercules-Corona Borealis Wall. M4: The First Rung, Inside the 5 Gpc Cap.

101. Spin-Filament Alignment. M3: Spin-Filament Alignment Explained by SCT Angular Momentum Inheritance.

102. Supervoid CMB Cold Spot Link. M2+M5: Shared Origin, Stacked Contributions.

103. Large Quasar Groups. M4+M11: Rare Lights on a Deposited Scaffold.

104. Ring Peculiar Galaxies. M3: Born This Way, Not Struck This Way.

105. The Big Ring (Ultra-Large Annulus at $z \approx 0.8$). M4: The Big Ring Annulus Predicted by SCT Shock-Shell Ring Morphology.

106. Early Supermassive Black Holes. M1: Early Supermassive Black Holes Born Big via SCT Collision Seeding.

107. Little Red Dots (JWST). M1: Little Red Dots Read by SCT as the Black Hole Seeding Epoch Made Visible.

108. JWST Early Massive Galaxies ($z = 14$, $M^* \approx 10^{10} M_{\odot}$). M1: JWST Early Massive Galaxies Explained by SCT Collision-Seeded Proto-Structures.

109. Overmassive High- z Black Holes (JWST). M1: Overmassive High- z Black Holes Explained by SCT Seeds-Before-Hosts.

110. Merger Rate Decline ($z > 2$). M1: High- z Merger Rate Puzzle Resolved by SCT Seeded Assembly.

111. Star Formation Rate Density. M1+M5: Fuel Delivered Early, Throttle Closed Late.

112. Galaxy Rotation Curve Universality Failure. M6: The Parameter That Varies Is Coherence.

113. JADES-GS-z14-0 Oxygen Abundance at $z = 14.18$. M1: JADES-GS-z14-0 Oxygen at $z = 14.18$ Explained by SCT Early Seeding Clock.

114. Bulge-Disk Decoupling. M3: Two Angular Momentum Settings of One Deposit.

115. Stellar to Halo Mass Relation. M6: Grading Against the Wrong Denominator.

116. Star Formation Efficiency. M1: The 50 Percent Was Never Real.

117. Galaxy Morphology Evolution Stall. M3: An Initial Condition Being Preserved.

118. Bar Fraction Evolution Anomaly. M3+M6: Early Disks to Host Them, No Live Halo to Brake Them.

119. Galaxy Size Evolution. M1+M5: Born Compact, Grown by a Loosening Mesh.

120. Compact Red Nuggets. M1+M5: Deposited Dense, Quenched Fast, Loosened Slowly.

121. Star Formation Cliff ($z \approx 2$). M5: Cosmic Noon Star Formation Cliff Explained by SCT Mesh Clock.

122. Dynamically Cold Rotating Disks at $z = 4-5$. M3: Cold Rotating Disks at $z = 4-5$ Explained by SCT Inherited Rotation.

123. Thick Disks and Warps. M3+M6: Inherited Components, Standing Tides.

124. Triangulum Warp. M6: Held by a Standing Tide, Not Excited by an Encounter.

125. Metallicity Floor. M11+M1: A First Generation That Was Fast and Everywhere.

126. UDGs with Excess Globular Clusters (DF44-Class, Disputed). M6: DF44-Class UDG Disputes Resolved by SCT Coherence Spectrum.

127. Core-Cusp Problem (5-Sigma). M6: No Particle, No Cusp, Nothing to Repair.

128. Missing Satellites (~ 50 Expected vs 20). M3: Missing Satellites Resolved by SCT Debris Condensation.

129. Centaurus A Satellite-Plane Corotation. M3: Centaurus A Satellite Plane Corotation Explained by SCT Debris Inheritance.

130. Satellite Plane Alignments (MW/LMC). M3: One Debris Field, One Plane.

131. M31 Satellite Plane. M3: The Same Stamp Twice, Because the Origin Was Shared.

132. Rarity of Magellanic-Cloud-Analog Pairs. M3: Magellanic Cloud Pair Rarity Explained by SCT Debris Fragmentation.

133. Carina Velocity Gradient. M3: The Rotation Came With the Galaxy.

134. Centaurus A Streams. M3: One Inheritance, Worn as Streams and as a Satellite Plane.

135. Radial Acceleration Relation / BTFR Tightness. M6: RAR and BTFR Tightness Explained by SCT Deterministic Coherence Law.

136. Field Halo Streams (GD-1, Pal 5). M6: The Prediction Is Silence.

137. Orphan Stream Plane. M6: One Potential, Because There Is Only One Thing There.

138. Sagittarius Stream Warp. M6: The Registered Wrap-Spacing Test.

139. Too-Big-To-Fail Problem. M6: No Subhalos to Fail.

140. SMC Bridge. M3: Born Together, Tethered Since.

141. Leo T Offset. M6+M11: Bound by Coherence, Fueled at Birth.

142. NGC 6822 Offset. M3: Two Axes From One Deposit, No Collider Required.

143. Sculptor dSph Flattening. M3+M6: Flattened by Inheritance, Cored by Coherence.

144. Segue 1 Velocity Dispersion. M6: Darkness That Scales With Coherence.

145. LMC Disk Integrity. M3+M6: A Cold Disk With Less to Fear.
146. Fornax 3 Core. M6: The Medium Was Never There.
147. Ursa Minor Structure. M6: A Quiet Harbor by Construction.
148. Pulsar Timing Array Nanohertz Excess. M1: PTA Nanohertz Background Excess Explained by SCT Seeded Binary Demographics.
149. Blue Transient Shocks. M1+M11: An Inventory the Cascade Stocked.
150. 3.5 keV X-Ray Line (Disputed). M6: No Particle to Decay.
151. The Final Parsec Problem. M5: The Universal Decay Channel Was Already Running.
152. Quasar Variability. M5: SCT Confirms the Clock and Predicts Its Residuals.
153. Fast Radio Bursts. M11+M1: Two Populations Because There Are Two Pedigrees.
154. Galactic Center GeV Excess. M6+M11: SCT Mandates the Astrophysical Origin.
155. Galactic 511 keV Positron-Annihilation Excess. M11: Galactic 511 keV Positron Excess Read by SCT as Seeded Bulge Demographics.
156. CIB-Galaxy Cross-Correlation. M1: Glow of an Early, Clustered Start.
157. Diffuse Gamma-Ray Background. M11: Diffuse Gamma-Ray Background Attributed by SCT to a Seeded Source Ledger.
158. Far-IR Background Excess. M1+M11: The Dust Was on Schedule, the Schedule Was Earlier.

159. Extragalactic Radio Background. M11+M1: The Demography Was Undercounted at the Source.

160. High Energy Neutrino Anisotropy. M1: IceCube Neutrino Attribution Crisis Solved by SCT Cocooned Seeds.

161. Fermi Bubbles Asymmetry. M1: Fermi Bubbles Asymmetry Explained by SCT Seeded Engine and Structured Halo.

162. Radio Axis Alignments. M3: One Vector, Worn by the Largest Things That Spin.

163. TeV-Blazar Transparency Anomaly (Disputed). M4: TeV Blazar Transparency Anomaly Explained by SCT Inhomogeneous EBL.

164. AMS-02 / PAMELA Positron Excess. M6+M11: SCT Requires the Pulsar Attribution.

165. UHECR Origins. M11: Acceleration the Census Never Counted.

166. IGM Magnetogenesis. M11: Void Magnetic Fields Explained by SCT Inherited Flux.

167. Cosmic Optical Background Excess (New Horizons). M11: New Horizons Optical Background Excess Read by SCT as the Deeper Ledger.

168. Quasar Optical-Polarization Alignments (Gpc Scales). M3: Gpc Quasar Polarization Alignments Explained by SCT J Inheritance.

169. Lithium-7 Plateau Discrepancy (Spite). M2: An Honest Non-Resolution.

170. Delta Reion Timing. M2: The Epoch Was Always Plural.

171. Ω_b from BBN vs CMB. M6: Ω_b BBN versus CMB Gap Read by SCT as Acoustic Chain Bias.

172. 21 cm Global Signal. M11: The Backlight Was Already Predicted Brighter.

173. Baryon-to-Photon Ratio (Emergent). M2: Baryon-to-Photon Ratio Derived by SCT Cascade Geometry at 0.17 Sigma.

174. Helium-4 Plateau Shift. M2: SCT Shares the Jeopardy Honestly.

175. Strong CP Non-Solving. M11: A Parameter With a History.

176. He-4 Abundance. M2: SCT Commits to the Standard Plateau with Declared Stakes.

177. Fine-Structure-Constant Spatial Dipole (Disputed). M6: SCT Predicts Structured Variation with the Coherence Field.

178. D/H Ratio. M2+M11: One Ratio, Many Histories.

179. He-3 Constraints. M11: The Helium-3 Problem Reframed by SCT Baseline Reservoir Dominance.

180. Reionization Photon-Budget Crisis (JWST). M4: JWST Photon Budget Crisis Resolved by SCT Deposited IGM Clumping.

181. Deuterium Scatter. M11: The Texture Is the Web's Processing History.

182. Lyman-Alpha Forest Power Deficit. M6: Suppression Is Registered, Not Optional.

183. Cosmic Neutrino Background. M2: Same Neutrinos, Different Bookkeeping.

184. Trans-Planckian Problem of Inflation. M1: Trans-Planckian Problem Unmasked by SCT Macroscopic Structure Generation.

185. DAMA/LIBRA Modulation vs Null Replications. M6: SCT Registered the Silence.

186. The Baryon Asymmetry Problem. M8: The Collision Supplies What the Smoothness Starved.

187. Quasar Luminosity Function. M1: The Census Was Stocked at Deposition.

188. Hierarchy Problem (Planck vs Weak). M7: No Planck Wall to Fall From.

189. EDGES vs SARAS-3 Global 21 cm Contradiction. M1: SCT Covers Both Branches.

190. Anomalous Microwave Excess (AME). M3: Rotation Inherited All the Way Down.

191. Spinning Dust Peak Shift. M3: The Missing Spin-Up Force Is History.

192. Halo Concentration Diversity. M6: The Third Dial Is Coherence.

193. Cluster Orientation Alignments. M3: A Boundary Condition, Not an Achievement.

194. Canis Major Overdensity. M3: Canis Major Overdensity Resolved by SCT as Deposited Disk-Edge Structure.

195. Point Source Foreground Overestimate. M11: CMB Point-Source Foreground Bias Predicted by SCT Deep Source Ledger.

196. Monoceros Ring. M3: Monoceros Ring Explained by SCT as Native Deposited Disk Structure.

197. CO Contamination 1-10%. M11: CO Line Contamination of CMB Maps Read by SCT as a Fuller Molecular Sky.

198. Radio Loop Foregrounds. M11: Radio Loop Foregrounds Read by SCT as Inherited Magnetic Architecture.

199. JWST $z > 10$ UV Luminosity-Function Excess. M1: JWST UV Luminosity Function Excess Predicted by SCT Power-Law Census.

200. Laniakea Strands. M3: Laniakea Flow Strands Explained by SCT Inherited Stream Kinematics.

201. Non-Thermal Pressure Fractions. M1: Low Non-Thermal Pressure Fractions Explained by SCT Settled-at-Birth Clusters.

202. Galactic Plane Emission Model Fail. M11: The Medium Is Diverse by Inheritance.

203. Synchrotron Depolarization. M11: A Field With a Pedigree.

204. Eastern Bubbles. M11: A Symmetric Outflow Met a Structured Medium.

205. Free-Free Optical Depth. M11: More Than One Kind of Ionized Gas.

206. Dust-to-Gas Ratios. M1: Early Dust Budget Crisis Resolved by SCT Factory-Count Head Start.

207. Shear B Modes. M6: Cosmic Shear B-Modes Read by SCT as the Mesh Vector Sector.

208. Dust Polarization Fraction. M11: Two Stubborn Ends, One Inherited Term.

209. Dust Lane Asymmetries. M3+M6: Held Off-Axis, Not Caught in Transit.

210. HI Mass Function High-Mass Excess. M11+M6: Reservoirs Delivered, Then Held Gently.

211. Radio Relics and Halos. M1+M6: Radio Relic Efficiency Crisis Resolved by SCT Deposited Fossil Floor.

212. Bullet Cluster Shocks. M6: The Offset Without the Particle.

213. Lensed-Quasar Flux-Ratio Anomalies. M6: Flux-Ratio Anomalies Explained by SCT Amplified Visible Substructure.

214. Cluster Mass Bias. M6: Two Methods Reading Two Different Masses.

215. Cool Core Dichotomy. M2: The Two Families Were Born, Not Made.

216. Thermal SZ Power Spectrum. M6: The Registered Suppression Exponent.

217. Planck SZ Cluster Counts vs Primary CMB. M6: Planck SZ Count Tension Resolved by SCT Two-Mass Structure.

218. Cluster Cooling-Flow Problem. M1: Cooling-Flow Problem Resolved by SCT Relic Entropy Floor.

219. Splashback Radius Discrepancies. M6: Splashback Displacement Explained by SCT Coherence Boundary.

220. El Gordo (ACT-CL J0102-4915). M1: El Gordo's Forbidden Parameters Predicted by SCT Cascade Kinematics.

221. Massive High-z Protoclusters Too Early. M1: Born-Hot Protoclusters Predicted by SCT Deposition Thermalization.

222. Gas-DM Offsets. M6: Gas-Lensing Offsets Explained by SCT Coherence Riding the Galaxies.

223. Kinetic SZ Null Hint. M5: Kinetic SZ Low Amplitudes Read by SCT as Apparent-Velocity Contamination.

224. ICM Metallicity Gradients. M11: Enriched Before Assembly, Everywhere at Once.

225. X-Ray Cavities Mismatch. M11: More Engines on the Books.

226. Cluster-Scale Core-Cusp (BCG/Strong-Lensing Cores). M6: Cluster Core-Cusp Problem Resolved by SCT Inward-Retiring Amplification.

227. Concentration-Mass Relation. M6: Concentration-Mass Relation Read by SCT as Coherence in an NFW Template.

228. Strong Lensing Arcs. M6: A Lens With Coherent Texture.

229. Entropy Floors. M1: ICM Entropy Floors Explained by SCT Relic Thermalization.

230. Cluster Substructure Excess. M6: Cluster GGSL Substructure Excess Explained by SCT Amplification Squared.

231. Pressure Turbulence Profiles. M1: Quiescent Cluster Turbulence Explained by SCT Settled Deposition.

REFERENCES

A. SCT Series Papers (From Chaos to Consilience). All by DR JM NIPOK (ORCID 0009-0006-3940-4450), CC BY-NC-SA 4.0.

NIPOK, DR JM (2026a). From Chaos to Convergent Foundations: The Foundational Premises of Successive Collision Theory (Paper 1). DOI: 10.13140/RG.2.2.19171.62243.

NIPOK, DR JM (2026b). From Chaos To Common Ancestry: A Hierarchical Frame-Tree Lorentzian Approach for High-Precision Cosmology (Paper 2). DOI: 10.13140/RG.2.2.21288.43521.

NIPOK, DR JM (2026c). From Chaos To Concordance Spectra: Demonstrating CMB Power Spectrum Compatibility with SCT (Paper 4). DOI: 10.13140/RG.2.2.20310.31042.

NIPOK, DR JM (2026d). From Chaos To Collisothermal Cosmogenesis: Early Structure Formation Under Successive Collision Theory (Paper 3). DOI: 10.13140/RG.2.2.16235.60968.

NIPOK, DR JM (2026e). From Chaos To Corotating Hierarchies: Angular Momentum Inheritance Across Seven Scales (Paper 5). DOI: 10.13140/RG.2.2.28263.10400.

NIPOK, DR JM (2026f). From Chaos to Cosmic Collisions: Changing One Lambda-CDM Assumption Brings Dark Matter Into The Light (Paper 11). DOI: 10.13140/RG.2.2.19379.69921.

NIPOK, DR JM (2026g). From Chaos To Cosmic Expansion: Recursive Tensor-Mesh Dissipation (Paper 14). DOI: 10.13140/RG.2.2.24304.72969.

NIPOK, DR JM (2026h). From Chaos To Constructive Relativity (this paper) (Paper 17). DOI: 10.13140/RG.2.2.23479.79528.

NIPOK, DR JM (2026i). From Chaos To Collapse Proof: An End to Black Hole Singularities (Paper 16). DOI: 10.5281/zenodo.18092309.

NIPOK, DR JM (2026j). From Chaos To Confirming Falsifiability: Predictions, Falsification Criteria, and Evidence (Paper 18). DOI: 10.13140/RG.2.2.19381.33765.

NIPOK, DR JM (2026k). From Chaos to Covariant Completeness: A Unified Mathematical Foundation (Paper 8). DOI: 10.13140/RG.2.2.29562.35527.

NIPOK, DR JM (2026l). From Chaos To Comoving Coordinates: The Logical Progression of Spacetime Geometry (Paper 7). DOI: 10.13140/RG.2.2.35762.06089.

NIPOK, DR JM (2026m). From Chaos to Coherent Gravity: The SCT Formalism That Solves the Dark Matter Problem (Paper 12). DOI: 10.13140/RG.2.2.22608.98560.

NIPOK, DR JM (2026n). From Chaos to Corroborated Action: The Unified Variational Foundation (Paper 9). DOI: 10.13140/RG.2.2.12280.81923.

NIPOK, DR JM (2026o). From Chaos To Complete Derivations: Closing the Remaining Derivations (Paper 10). DOI: 10.13140/RG.2.2.32413.47840.

NIPOK, DR JM (2026p). From Chaos To Codified Acoustics: A Low-Parameter Collision Geometry Unifying DESI-DR2, DES-Y6, HSC-Y3, and KiDS-DR5 (Paper 15). DOI: 10.13140/RG.2.2.10321.29288.

NIPOK, DR JM (2026q). From Chaos To Coalescent Parsimony: Carrier-Sphere Interference Foundation (currently under peer review) (Series 2 Paper 1). DOI: 10.13140/RG.2.2.14355.03366.

B. Selected Observational Papers Cited

Riess A. G. et al. (2022). "Hubble Constant from SH0ES." *Astrophys. J. Lett.* 934, L7., $H_0 = 73.0 \pm 1.0$. Planck Collaboration (2020). "Planck 2018 results. VI." *Astron. Astrophys.* 641, A6., $H_0 = 67.4 \pm 0.5$; $A_{\text{lens}} = 1.18 \pm 0.065$; $n_s = 0.9649 \pm 0.0042$. DESI Collaboration (2025). "DESI DR2 Results II: Cosmological Constraints from BAO." arXiv:2503.14738 (superseding DESI DR1 2024, arXiv:2404.03002)., dynamical-dark-energy preference strengthened to $\sim 2.8\text{-}4.2 \sigma$ with $w_0 > -1$ (quintessence-like) at low z and $w_a \approx -1.05$; $r_d = 147 \pm 1$ Mpc. Carniani S. et al. (2024). "JADES-GS-z14-0 spectroscopic confirmation at $z = 14.18$." Witstok J. et

al. (2024). “MoM-z14: Strong N/C enrichment at $z = 14.44$.” Xiao M. et al. (2024). “ $\epsilon_* \approx 47\text{-}52\%$ requirement at $z = 7\text{-}9$.” Weibel A. et al. (2025). “Quenched massive galaxies at $z = 7.29$: 100-1000 \times above Λ CDM simulations.” Nature. Zhou R. et al. (2025). “tSZ thermal energy excess in SPT2349-56 at $z = 4.3$: 6.4 σ above TNG-Cluster.” Nature 536, 1226. Tang Q. et al. (2025). “Cluster rotation velocity scaling: 360 km/s at $10^{14} M_{\odot}$ rising to 693 km/s at $10^{15} M_{\odot}$.” Tudorache C. et al. (2025). “First detection of bulk angular momentum in an individual cosmic filament via MeerKAT 21 cm HI.” West M. J. et al. (2025). “Cluster major-axis correlated alignment over 200-300 comoving Mpc.” Smith R. J. et al. (2023). “BCG-cluster shape alignment in place at $z > 1.3$ at one-in-a-million significance.” Mandarakas N. et al. (2021). “VLBI 3D jet alignment at 400-900 Mpc with $> 99.5\%$ significance.” Kroupa P. et al. (2024). “Joint probability of co-rotating satellite plane alignments across six independent host systems.” Meneghetti M. et al. (2020). “GGSL excess in 11 Hubble Frontier Fields clusters.” Science 369, 1347. Migkas K. et al. (2021). “ $\sim 3 \sigma$ directional cosmological-parameter variations in Planck data.” Miller M. C. et al. (2021). “PSR J0740+6620 NICER radius: $13.7 +1.9/-1.5$ km.” Riley T. E. et al. (2019). “PSR J0030+0451 NICER radius: $13.02 +1.24/-1.06$ km.” LIGO/Virgo Collaboration (2017). “GW170817 binary neutron star merger.”, $\Lambda_{1.4} < 800$. Spite F. & Spite M. (1982); Sbordone L. et al. (2010). “Primordial Li-7 abundance: factor-3 deficit.”

C. Foundational Physics References

Penrose R. (1965). “Gravitational collapse and space-time singularities.” Phys. Rev. Lett. 14, 57. Hawking S. W. & Penrose R. (1970). “The singularities of gravitational collapse and cosmology.” Proc. Roy. Soc. A 314, 529. Buchdahl H. A. (1959). “General relativistic fluid spheres.” Phys.

Rev. 116, 1027. Hinderer T. (2008). "Tidal Love numbers of neutron stars." *Astrophys. J.* 677, 1216. Damour T. & Nagar A. (2009). "Surface energy-density correction to the tidal Love number of self-bound stars."

DR JM NIPOK ORCID 0009-0006-3940-4450 <https://www.linkedin.com/in/dr-jm-nipok-15084a3a1/> <https://linktr.ee/thenaturalstateofnature/>
<https://thenaturalstateofnature.org/231/>